GEVINUS TANCOLO, Gould.

GEVINUS TANCOLO, *Gould.*

Formosan Green Woodpecker.

Gecinus tancolo, Gould in Proc. of Zool. Soc. 1862, p. 283.

——— *tancola*, Swinh. in Ibis, 1863, p. 389.

Tancolo, aborigines of Foochow.

THE present bird, the *Gecinus occipitalis* and *G. squamatus* of the Himalayas, and the *G. viridis* and *G. canus* of Europe, all belong to the same genus, and there are other Indian species which might have been mentioned as pertaining to this very distinct form of Woodpeckers. All these birds evince a partiality for the ground, where they search for the pupæ of ants, terrestrial insects, and their larvæ. When characterizing this bird in the 'Proceedings of the Zoological Society of London' above referred to, I mentioned that the *G. tancolo* is most nearly allied to, but quite distinct from, the *G. occipitalis*; and in confirmation of this view I may quote Mr. Swinhoe's remarks on the subject from the 'Ibis' for 1863, p. 389 :—

"The Formosan Green Woodpecker is a local representative of the larger Himalayan *G. occipitalis*, which is, however, at once to be distinguished from it by its greater dimensions, by its large and entirely black bill, by the sides of its neck being yellowish green instead of grey, by the brighter yellowish green of the breast, belly, and back, by its lateral rectrices being wholly brown instead of partially brownish white, and by its primary coverts being margined on the outer web with golden green instead of being barred with brown. The wing is shorter in the Formosan bird; and the primaries have fewer whitish spots, and only indications of some on the outer edge of the first primary, instead of distinct spots; and a grey eye-streak divides the black on the lores from the red frontal crest."

Lores, a narrow band across the forehead, back part of the head, nape, and a stripe down the cheeks black; centre of the forehead blood-red; back dull green, passing into greenish yellow on the rump; shoulders and upper part of the wings dull wax-yellow; primaries olive-brown, with small elongated marks of buff on their external margins; internal webs of the greater coverts and primaries crossed with distinct bars of greyish white; throat and cheeks grey; under parts of the shoulders and axillaries alternately barred with greenish white and blackish brown; chest and under surface sordid green; "bill blackish grey on the upper and nearly the whole apical half of the lower mandible, the basal edge of the former and the rest of the latter being greenish yellow; legs deep leaden, with a tinge of olive-green; sole-pads brownish; claws leaden black; irides pearly white." (Swinhoe.)

The female differs in having the crown grey, broadly streaked with black.

The Plate represents the two sexes, of the size of life.

Gould, John. 1864. "Formosan Green Woodpecker, Gecinus tancol [Pl. 35]."
The Birds of Asia 6(XVI), -. <https://doi.org/10.5962/p.323340>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122488>

DOI: <https://doi.org/10.5962/p.323340>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323340>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.