IYNGIPICUS AURANTIIVENTRIS.

IYNGIPICUS AURANTIIVENTRIS.

Orange-breasted Pygmy Woodpecker.

Picus (Bæopipo) aurantiiventris, Salvad. Atti R. Accad. Sci. Torino, iii. p. 524 (1868).

Iyngipicus aurantiiventris, Salvad. Uccelli di Borneo, p. 41, tav. iv. fig. 2 (1874).—Sharpe, Proc. Zool. Soc. 1881, p. 792.

Iyngipicus aurantiiventris, Sharpe, Ibis, 1879, p. 240.

As far as we know at present, this species is strictly confined to the island of Borneo, where it is especially found in the Sarawak district. It was here that it was first discovered by the Marquis Doria and Dr. Beccari. It appears to me to be a very distinct species, belonging to the section of the genus *Iyngipicus* which contains the species with the occipital tuft on each side of the head. The orange colouring on the belly also appears to be present in every specimen, whether young or old, though, as might be expected, it is more developed in the fully adult birds. The four central tail-feathers are black in both sexes, while the conspicuous black striping of the upper tail-coverts likewise appears to me to be a striking character. The British Museum has received several specimens from the Messrs. Everett in Borneo. I have in my own collection a skin obtained by Mr. Alfred Everett at Mateng, while the Museum possesses examples shot by Mr. Henry Everett at Paku. Mr. Lowe met with the bird in Lumbidan, where it was also procured by Governor Ussher and Mr. Treacher. Lastly, Mr. W. B. Pryer has recently obtained the species in the district of Sandakan, in North-eastern Borneo.

As regards the habits and mode of life we know absolutely nothing, Mr. Treacher merely stating that in Lumbidan the native name is "Burong anie putie."

The colour of the iris, according to the Marquis Doria, is whitish blue.

For the benefit of ornithologists I translate the description given by Count Salvadori:—

"*Adult male*. Above black, varied with white; head dusky grey, surrounded by black behind; the occiput ornamented with a lateral streak of red; back banded with white and black; sides of head and of neck white; a broad band behind the eyes blackish dusky; throat and neck whitish; an obscure malar band of dusky; underneath lurid white, streaked with dusky black; the breast and the abdomen tinged with orange-yellow; wings black, spotted with white on the inner margin; three median tail-feathers black, the lateral ones banded with white on both webs; bill dusky horn-colour; feet lead-colour; iris whitish blue.

"*Adult female*. Very similar to the male, but distinguished by the absence of the red streak on the sides of the occiput, and by the paler orange-yellow colour of the abdomen."

The Plate represents a pair of birds of the natural size, the female being in my own collection, whilst the male is from a skin in the British Museum.

[R. B. S.]

Gould, John. 1882. "Orange-breasted Pygmy Woodpecker, *Iyngipicus aurantiiventris* [Pl. 23]." *The Birds of Asia* 6(XXXIII), –.
<https://doi.org/10.5962/p.323371>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122488>

DOI: <https://doi.org/10.5962/p.323371>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323371>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.