MYCEROBAS MELANOXANTHUS.

J. Gould and H.C. Richter del. et lith.

Hillman and L. Waller. Imp.

MYCEROBAS MELANOXANTHUS.

Black and Yellow Grosbeak.

- Coccothraustes melanozanthos*, Hodgs. Asiat. Research., vol. xix. p. 150.—Ib. Journ. Asiat. Soc. Beng., vol. xiii. part ii. p. 950. pl. fig. 3, bill.
— *melanozanthus*, Blyth, Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 125.—Cat. of Spec. and Draw. of Mamm. and Birds presented to Brit. Mus. by B. H. Hodgson, Esq., p. 105.—Gray and Mitch. Gen. of Birds, vol. ii. p. 358, *Coccothraustes*, sp. 5.
— *fortirostris*, Lafr.
Mycerobas melanozanthus, Bonap. Consp. Gen. Av., p. 505.
-

Mr. HODGSON states that the true habitat of the Black and Yellow Grosbeak is the northern regions of India, but that in summer it wanders from thence into its central portions in search of the ripe stony fruits and berries which constitute its principal food, and for crushing which its enormously developed and powerful mandibles seem admirably adapted. A female or young bird is figured among the drawings of the late Hon. F. J. Shore, from a specimen killed by him on the 17th of March 1828, a little above Najpore.

The male has the head, neck, throat, breast, upper surface, wings and tail sooty black; the third, fourth, fifth and sixth primaries with a mark of snow-white at the base, the scapularies with a large oblong mark of yellowish white near the tip of their outer webs, the secondaries and all but the first four primaries with a small narrow mark of the same colour near the tip of their outer webs; abdomen and under tail-coverts yellow; thighs mottled black and yellow; bill purplish horn-colour; feet purplish brown.

The female has the upper surface dull black, mottled with yellow over the head, back of the neck and down the centre of the back; wings similar to those of the male, but the coverts also bordered with yellowish white; rump and upper tail-coverts narrowly bordered with olive; under surface yellow, each feather with a tear-shaped mark of black down the centre, these marks becoming longer and more conspicuous on the flanks.

I am not quite certain that the fully adult female does not lose the spotted markings of the breast; the changes probably resembling those which take place in the Greenfinch (*Fringilla chloris*, Linn.) of Europe.

In some specimens the yellow is so pale as to be almost white.

The young are said to resemble the female until the second moult, and then gradually to assume the adult plumage.

The figures represent birds in the three states above described of the natural size.

Gould, John. 1851. "Black and Yellow Grosbeak, Myserobas melanoxanthus [Pl. 20]." *The Birds of Asia* 5(III), -. <https://doi.org/10.5962/p.323402>.

View This Item Online: <https://www.biodiversitylibrary.org/item/121124>

DOI: <https://doi.org/10.5962/p.323402>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323402>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.