



PALEORNIS MALACCENSIS.



## PALÆORNIS MALACCENSIS.

### Malacca Parrakeet.

- Psittacus longicauda*, Bodd. Tabl. des Pl. Enl. Daub., p. 53.  
*Palæornis longicauda*, Gray and Mitch. Gen. of Birds, vol. ii. p. 410, *Palæornis*, sp. 10.—Horsf. Cat. of Birds in Mus. East Ind. Comp., vol. ii. p. 618.  
*Psittacus Malaccensis*, Gmel. Edit. Linn. Syst. Nat., tom. i. p. 325.  
*Le Grand Perruche à longs brins de Malac*, Buff. Hist. des Ois., tom. vi. p. 155.—Pl. Enl. 887.  
*La Perruche à nuque et joues rouges*, LeVaill. Hist. des Perr., pl. 72.—Bourj. St. Hil. Coll. des Perr., pl. 1.  
*Belurus malaccensis*, Bonap. Rev. Zool. 1854, p. 152.—De Souancé, Rev. Zool. 1856, p. 208.  
*Palæornis malaccensis*, Vig. Zool. Journ., vol. ii. p. 52.—Selby in Jard. Nat. Lib. Parrots, p. 75. pl. 3.—Blyth, Journ. Asiat. Soc. Beng., vol. xi. p. 788, and xix. p. 233.—Ib. Cat. of Birds in Mus. Asiat. Soc. Beng., p. 5.—Mottl. and Dill. Nat. Hist. of Lab., p. 26.  
*Psittacus erubescens*, Shaw, Gen. Zool., vol. viii. p. 437.—Raffles, Trans. Linn. Soc., vol. xiii. p. 231.  
——— *ginginianus*, var. C., Lath. Ind. Orn., vol. i. p. 99.—Ib. Gen. Hist. of Birds, vol. ii. p. 165.  
——— *barbatulatus*, Bechst.  
*Conurus barbatulatus*, Kuhl, Mon. Psitt. in Abhand., No. 38.  
*Belocercus barbatulatus*, Müll. et Schleg.  
*Palæornis erythrogenys*, Less. Traité d'Orn., p. 215.  
*Malacca Parrakeet*, Lath. Gen. Syn., vol. i. p. 241. no. 39 C, and Supp., p. 60.  
*Blossom-cheeked Parrakeet*, Shaw, Gen. Zool., vol. viii. p. 437.  
*Psittacus erythrocephalus*, var. *δ. malaccensis*, Gmel. Edit. Linn. Syst. Nat., tom. i. p. 325.  
*Burong Bayau*, Sumatrans (Raffles).  
*Madna Bhola* of the Calcutta Bird-dealers (Blyth).

As its specific name implies, this truly elegant species is a native of Malacca; it also occurs in Sumatra, Borneo, and Labuan, but I have never yet seen it from Java.

Sir Stamford Raffles states that "it is this Parrot that figures in the fables and poetry of the Malays, in which it is represented to be endowed with a supernatural degree of intelligence."

Messrs. Mottley and Dillwyn inform us in their "Natural History of Labuan," that "these handsome Parrakeets are not uncommon, and are to be seen in the early morning flying about above the tops of the trees in small flocks of six or eight, uttering in their flight a loud quick scream, very much like the note of the Common Swift. They are particularly fond of the fruit of the *Dryabalanops Camphora*, which they split open, and eat the curious crumpled cotyledons, in spite of their pungent taste and smell of turpentine. A specimen was shot while feeding upon the seeds of *Dillenia speciosa*, a shrub about ten or fifteen feet high; and it is the only instance in which we have known them venture so near the ground. When first seen, he was busy opening the capsules of the plant, and scraping out the seeds with his beak, never omitting to clip off at a single bite every one he emptied; having done this, he dropped himself under the twig he sat on, swinging by one leg to watch its fall; when it reached the ground he testified his satisfaction by a low chirp, and giving himself a vigorous swing, caught the perch with his other foot, and walked gravely along to another capsule, not hopping, but placing one foot before the other in a most old-fashioned way. Another of these Parrakeets, which had been pinioned by a shot without being otherwise injured, was placed in a cage, where, soon finding his two long tail-feathers to be an incumbrance, he deliberately turned round, pulled them out, and then walked round the cage, evidently to try the effect of his contrivance."

So far as I am aware, no difference occurs in the colouring of the sexes; but Mr. Moore has given the following description of what he considers the youthful plumage:—

"The young has the plumage yellowish green, darkest on the crown and sides of the throat, and palest beneath; wings above the same, and having the primaries, secondaries, a portion of the tertiaries and speculars bluish on their outer webs, and the three former narrowly edged with yellowish; under wing-coverts and axillaries green; rump more bluish green, and upper tail-coverts bright yellowish green; tail bluish green, edged with yellowish green, the latter beneath dingy yellowish green; before the eye, slightly above, and broadly beneath, ferruginous, intermixed with greenish yellow; upper mandible red, tip and under one also pale."

In the adult the crown of the head is bright grass-green; face, sides of the head, and sides and back of the neck fine deep red, suffused with a vinous bloom; moustache jet-black; back yellowish green, suffused with delicate blue; lower part of the back verditer-blue; upper tail-coverts green; wings green, washed with orange on the centre; primaries deep bluish green, with a tinge of yellow on the margin and blackish brown inner webs; under surface pale greenish yellow; sides of the body and under wing-coverts wax-yellow; centre tail-feathers blue; lateral tail-feathers grass-green; bill red, paler at the tip.

The bird is represented on the accompanying Plate of the size of life.



Gould, John. 1858. "Malacca Parrakeet, *Palaeornis malaccensis* [Pl. 10]." *The Birds of Asia* 6(X), -. <https://doi.org/10.5962/p.323420>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/122488>

**DOI:** <https://doi.org/10.5962/p.323420>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/323420>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Smithsonian Institution Libraries

**Copyright & Reuse**

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.