PTILOCOENAS CRUENTA.

J. Gould and H. C. Richter del. & lith.

Walter. imp.

PHLOGÆNAS CRUENTA.

Red-breasted Pigeon.

- La Tourterelle grise ensanglantée*, Sonn. Voy. a la Nouv. Guin., p. 52, pl. 21.
——— *blanche ensanglantée*, Sonn. Id., p. 51, pl. 20.
Red-breasted Turtle, Lath. Gen. Syn., vol. iv. p. 657.
Sanguine Turtle, Lath. Id., p. 657, and Gen. Hist., vol. viii. p. 91.
Columba cruenta, Gmel. Edit. Linn. Syst. Nat., tom. i. p. 785.—Lath. Ind. Orn., vol. ii. p. 611.
——— *sanguinea*, Gmel. Id., p. 785.—Lath. Ind. Orn., vol. ii. p. 611.
Columbi-galline poignardé, Knip et Temm. Les Pig. part ii (Les Colombi-gallines), p. 16, pl. 8 et pl. 9, var.
Columba Luzonica, Scop. Del. Flor. et Faun. Insub.
——— *nivea*, Scop., var.
Red-breasted Pigeon, Lath. Gen. Hist., vol. viii. p. 90.
Calenas Luzonica, Gray, List of Spec. of Birds in Coll. Brit. Mus., part iii. p. 18.
Calenas ? luzonica, Gray, Gen. of Birds, vol. ii. p. 478, *Calenas*, sp. 3.
——— *cruenta*, Cab.
Phlegœnas luzonica, Reich. Syst. Av., t. ccxxv. fig. 1265, var., et tab. ccxxvii. f. 2479.
Phlegœnas cruenta, Bonap. Consp. Gen. Av., tom. ii. p. 88, *Phlegœnas*, sp. 1.
Phlogœnas cruenta, Sclat. in Proc. of Zool. Soc., 1863, p. 377.
Phlegœnas luzonica, Wall. Ibis, 1865, p. 392.

AMONG the many advantages afforded to the ornithologist by the unrivalled collection of birds contained in the menagerie of the Zoological Society of London, are the opportunities given him from time to time of becoming acquainted with living examples of species of which probably he had before only seen but indifferent skins; he is thereby enabled not only to observe their individual peculiarities, but to ascertain many details as to their colouring, particularly of their soft parts, which, from their fading immediately after death, could not otherwise be ascertained. The circumstance of several living examples of the old *Columba cruenta* of Gmelin, the Red-breasted Pigeon of Latham, being at this time (March 1866) in the Gardens of the Society, enables me to give a correct delineation of a bird which has been made the type of the genus *Phlogœnas* by Reichenbach—a form of which four species are all that are known. They are all insular birds, being natives of Manilla, Java, Sumatra, the Celebes, and New Guinea. The one here represented is said to be from the Philippines, but from which of them is still uncertain, although we have been aware of the existence of the bird for nearly eighty years; Latham merely says:—"Perouse met with these, which he called 'Stabbed Doves,' at Morvula, one of the Philippine Islands." Judging from the living examples in the Zoological Society's Gardens there appears to be little or no outward difference in the sexes. Like many other members of the *Columbidæ* or family of Pigeons, they readily become accustomed to captivity; and if any foreign species could be domesticated and acclimatized in this country, the Red-breasted Pigeon seems to be the one with which the experiment would be most likely to succeed, as their long legs would indicate them to spend much of their time on the ground.

Forehead and crown delicate grey; occiput and hinder part of the neck deep violaceous grey with purple reflexions; back, scapularies, lesser wing-coverts, and sides of the breast slaty grey with purple and red reflexions, each feather with a lunate mark of metallic green at the tip; throat and breast white below, while all the under surface is of a clear pale cinnamon hue, at the junction of the two colours a large patch of blood-red, giving the bird a wounded appearance, whence the specific name; middle and greater wing-coverts reddish purple for three-fourths of their length, their tips being grey and forming three bands across the wing; the primaries and secondaries deep greyish brown, narrowly margined with reddish; two middle tail-feathers greyish brown, the remainder grey at the base, crossed in the middle by a band of black and tipped with ash-grey; irides dark brown; bill blackish brown; nostrils grey; legs and feet purplish red.

The figures are of the size of life. The plant is the *Sonerila margaritacea*.

Gould, John. 1866. "Red-breasted Pigeon, *Phlogoenas cruenta* [Pl. 51]." *The Birds of Asia* 6(XVIII), -. <https://doi.org/10.5962/p.323435>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122488>

DOI: <https://doi.org/10.5962/p.323435>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323435>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.