UROCISSA FLAVIROSTRIS.

J. Gould and H. Richter, del. et lith.

E. H. Mandel & Walton, Imp.

UROCISSA FLAVIROSTRIS.

Yellow-billed Blue Pie.

Psilorhinus flavirostris, Blyth, Journ. Asiat. Soc. Beng., vol. xx. p. 28.—Ib. Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 93.

Calocitta flavirostris, Bonap. Consp. Gen. Av., p. 381, *Calocitta*, sp. 3.

Urocissa flavirostris, Cab. Mus. Hein., p. 87.—Horsf. and Moore, Cat. of Birds in Mus. East Ind. Comp., vol. ii. p. 578.—Gould in Proc. of Zool. Soc., part xxvii. p. 200.

HERE at least we have a bird which the most sceptical of ornithologists must admit to be specifically distinct from the other Blue Pies of India and China, *U. occipitalis* and *U. sinensis*,—its much duller colouring, combined with its bright yellow bill, its lengthened and slenderly-formed tarsi, and the absence of all spotting on the crown, being characters which are surely sufficient to indicate its individuality; Mr. Blyth's indications, however, of the points of difference, which I give below word for word, will be the best evidence on this point. Of course, it would be useless to compare it with either of the red-billed species; nor can it, in my opinion, be confounded with the bird to which I have given the name of *U. cucullata*. It is true that both the *flavirostris* and *cucullata* have yellow bills, but there, as will be seen on reference to the figures and descriptions of the two species, their similarity ends.

In pointing out the specific differences of this bird, Mr. Blyth says, "General plumage of a much duller colour than in the preceding (*occipitalis*); the bill of the recent specimen bright yellow, and the white of the occiput reduced to a narrow transverse band, with a broad collar of black below it, surrounding the hind-neck, and never any white tips to the feathers immediately above it; legs and toes small and slender."

The native countries of the *Urocissa flavirostris* are Sikkim and Bhotan.

The vast primæval forests of the southern slopes of the Himalayas, extending over an area embracing many degrees of longitude, must still contain an abundance of undiscovered stores for the student of zoology; for if we find such conspicuous objects as the members of the genus *Urocissa* differing in countries only a few degrees apart, there also will be found numerous species of more diminutive birds, insects, and mammals which at present are entirely unknown to us.

The sexes are alike in size and colouring.

Head, neck, throat and breast black; within the black at the back of the neck a crescentic mark of bluish white; all the upper surface dull bluish brown; wings dull brownish blue; the inner webs of the primaries and secondaries brownish black; all the secondaries crescented with white at the tip; primaries margined with dull brownish white and with a small oblique mark of white at the tip; upper tail-coverts dull blue tipped with black, behind which is a narrow transverse line of bluish white; two centre tail-feathers blue tipped with white; the remainder blue tipped with white, the two colours separated by a broad band of black, immediately behind which is a mark of white, which gradually expands into a wide band crossing both webs as the feathers recede from the centre; under surface dull bluish white; bill and legs yellow.

The Plate represents the bird about three-fourths of the size of life.

Gould, John. 1861. "Yellow-billed Blue Pie, *Urocissa flavirostris* [Pl. 50]." *The Birds of Asia* 5(XIII), -. <https://doi.org/10.5962/p.323496>.

View This Item Online: <https://www.biodiversitylibrary.org/item/121124>

DOI: <https://doi.org/10.5962/p.323496>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323496>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.