TROCISSA OCCIPITALIS.

J. Gould and W. C. Richter, del. & lith.

H. W. Henslow & Walton, imp.

UROCISSA OCCIPITALIS.

Nepaulese Blue Pie.

- Pica erythrorhyncha*, Vig. in Proc. of Comm. of Sci. and Corr. of Zool. Soc., part i. p. 173.—Gould, Century of Birds, pl. 41.
- Cyanurus erythrorhynchus*, Hodgs. in Gray's Zool. Misc. 1844, p. 84.
- Psilorhinus sinensis*, Gray, Cat. of Spec. and Draw. of Mamm. and Birds pres. to Brit. Mus. by B. H. Hodgson, Esq., p. 101.
- *albicapillus*, Blyth, Journ. Asiat. Soc. Beng., vol. xv. p. 28, young.
- Urocissa albicapillus*, Cab. Mus. Hein., p. 87, note.
- Psilorhinus occipitalis*, Blyth, Journ. Asiat. Soc. Beng., vol. xv. p. 26.—Hutton, Journ. Asiat. Soc. Beng., vol. xvii. part ii. p. 681.—Adams in Proc. of Zool. Soc., part xxvi. p. 479.
- Urocissa occipitalis*, Cab. Mus. Hein., p. 87, note.—Adams in Proc. of Zool. Soc., part xxvii. p. 172.—Gould in Proc. of Zool. Soc., part xxvii. p. 200.

If this be not the oldest known species of this elegant and truly Indian group of Pies, it is the one most generally seen in the museums of Europe. To say that it is more beautiful than its near allies would scarcely be correct, as the species of this genus are all equally elegant in their form and equally lovely in tints: these charming members of the family CORVIDÆ must indeed form conspicuous objects among the foliage of the forest trees of Upper India, and the landscape be greatly enlivened by their presence.

We have much yet to learn respecting the true habitats of the various species of the genus *Urocissa*, each of which evidently enjoys a range over a distinct part of the country without encroaching on the others' territory. Of the three species inhabiting the temperate regions of the southern slopes of the Himalayas, the present would appear to frequent the middle or Nepaulese country, the *U. cucullata* the more western districts to the north-west of Simla, and the *U. flavirostris* Sikkim and Bhotan.

The *U. occipitalis* may be at once distinguished from the *U. flavirostris* and the *U. cucullata* by the spotted character of the feathers of the forehead and the rich red colouring of the bill. There are still two other species bearing even a greater resemblance to this bird, namely, the *U. Sinensis* and the *U. magnirostris*; from the former of these it differs in the lighter colouring of the nape and in the greater extent of the white tipping of the tail-feathers, and from the latter by the much smaller size of its bill.

Captain Hutton states that "This species occurs at Masuri throughout the year, collecting into small parties of four to six during winter. It breeds at an elevation of 5000 feet, in May and June, making a loose nest of twigs externally, lined with roots. The nest is built on trees, sometimes high up, at others about eight or ten feet from the ground. The eggs are from three to five in number, of a dull greenish ash-grey, blotched and speckled with brown dashes, confluent at the larger end. Diameter $1\frac{1}{8} \times \frac{1}{8}$ inch; the ends nearly equal in size. The bird is very terrene in its habits, feeding almost entirely on the ground."

Dr. Adams informs us that this species "is not uncommon in the jungles and woods of the lesser ranges round Simla and Kussowlee," that "the habits of the bird resemble those of the Magpie," and that "the presence of a cheetah, or a bird of prey, is often discovered by the chattering of these beautiful creatures."

The Hon. F. J. Shore says, "A specimen of this bird which I kept in captivity, though it refused other food, ferociously pounced upon living birds, presented to it by way of experiment, and eagerly devoured them. Seen among the foliage of the trees, it is an elegant and conspicuous object, flitting from bough to bough, with long and flowing tail, and full of vivacity in every movement."

The sexes are so much alike in size and colouring that it is impossible to distinguish them without the aid of dissection.

Bill and legs orange-red; forehead and crown black, with a spot of white at the tip of each of the feathers of the latter; cheeks, sides of the neck, throat and breast black; occiput and back of the neck white; all the upper surface dull purplish blue; shoulders and the outer webs of the primaries and secondaries fine blue, their inner webs brownish black; all the secondaries crescented with white at the tip; primaries margined along the middle portion of their outer web with bluish white, and with a small linear mark of white near the tip; upper tail-coverts dull blue at the base, fading into bluish white towards their apices and largely tipped with black; middle tail-feathers blue, largely tipped with white; the next on each side blue, not so largely tipped with white as the middle ones, the blue and the white separated by an imperfect band of black blending posteriorly with the blue; the four lateral feathers are blue at the base and white at the tip, the two colours being separated by a broad and distinct band of deep black; immediately behind this black band, a triangular white mark occurs on the inner web, which gradually increases in extent until on the outer feather it forms an oblique band across both webs; all the under surface creamy white with a very faint wash of blue.

The Plate represents the bird about three-fourths of the size of life.

Gould, John. 1861. "Nepaulese Blue Pie, *Urocissa occipitalis* [Pl. 47]." *The Birds of Asia* 5(XIII), -. <https://doi.org/10.5962/p.323498>.

View This Item Online: <https://www.biodiversitylibrary.org/item/121124>

DOI: <https://doi.org/10.5962/p.323498>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323498>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.