VIVIA INNOMINATA.

VIVIA INNOMINATA.

Speckled Piculet.

- Picumnus innominatus*, Burton, in Proc. of Zool. Soc., part iii, 1835, p. 154.—Blyth, Journ. Asiat. Soc. Beng., vol. xii. p. 1005.—Id. Cat. of Birds in Mus. Asiat. Soc. Calc., p. 65.—Gray & Mitch. Gen. of Birds, vol. ii. p. 432, *Vivia*, sp. 11.—Gray, Cat. of Spec. and Draw. of Mamm. and Birds pr. to Brit. Mus. by B. H. Hodgson, Esq., p. 114.—Malh. Mon. des Piciés, vol. ii. p. 278, vol. iv. pl. 116. figs. 5 & 6.
- Vivia innominata*, Bonap. Consp. Gen. Av., p. 140.—Horsf. & Moore, Cat. of Birds in Mus. East-Ind. Comp., vol. ii. p. 677.—Jerd. Birds of India, vol. i. p. 300.—Beav. Ibis, 1865, p. 411; 1869, p. 415.
- *nepalensis*, Hodgs. Journ. Asiat. Soc. Beng., vol. vi. p. 107.
- Piculus nipalensis*, Hodgs. Journ. Asiat. Soc. Beng., vol. x. p. 29.—Id. Gray's Zool. Misc., 1844, p. 85.
- *rufifrons*, Hodgs. Gray's Zool. Misc., 1844, p. 85.
- Wee-wee* of the Nepaulese (Hodgson).

This little frequenter of branches, to which the generic name of *Vivia* has been applied, is an inhabitant of the Himalayas, and is the only species of the genus yet discovered. It is clearly a representative, in the Old World, of the many members of the genus *Picumnus* in the New. They all bear a very general resemblance, and in their respective localities doubtless perform the same offices, whatever they may be; I say whatever they may be, because we really know little or nothing of the habits and economy of these diminutive creatures. Insects extracted from the bark of the larger trees probably constitute their principal food; but whether they are truly scansorial, like the Woodpeckers, and merely ascend the trees in search of them, or if they have the power of running down as well as up their boles, like the Nuthatches, we know not. Of their cry, if they have any, of their mode of nidification, whether their eggs are white like those of the Woodpeckers, Barbets, and Wrynecks, or speckled like those of the Nuthatches, we are alike ignorant; and any information on these points would prove acceptable and, doubtless, interesting.

Mr. Jerdon and Capt. Beavan have each written a few lines on this species, which I shall take the liberty of repeating here; for, of my own knowledge, I have nothing to say respecting it.

"This interesting little bird," says Mr. Jerdon, "is found throughout the Himalayas, and in no other locality that I am aware of. Its range extends, so far as known, from three to six thousand feet, or so. It is found in tangled brushwood, and among dead and fallen trees in damp spots, hunting about among the decaying bark for various insects. It is said to breed in holes of trees.

"Its nearest ally is the *Picumnus minutus* of Temminck, from South America, which is very similar in colouring, but has a stronger bill, with the head red on the forehead, black with white spots posteriorly, like so many Woodpeckers, to which, in fact, these little birds are nearly related; the tongue of *V. innominata* is strictly *Picine*."

"Dr. Jerdon says," remarks Capt. Beavan, "that this species is found throughout the Himalayas, and 'in no other locality that I am aware of.' But an observation from Mr. Blyth in the Journal of the Asiatic Society of Bengal for 1859, p. 416, shows that it also occurs in Burmah. I procured a specimen at Simla on the 3rd of July, 1866, which is now, I believe, in Col. Tytler's collection. The dimensions of the bird, in the flesh, were as follows:—Length 4·125, wing 2·25, tail 1·25, tarsus ·4375, spread of foot 1·0625, bill from front 1·4375, from gape ·5, extent 7 inches. Legs light blue; bill bluish lead-colour; head apparently brown. The tongue of this species, as Dr. Jerdon remarks, is like that of a Woodpecker, having several barbs along its extremity, resembling, in this respect, many of the arrows manufactured by savage nations, who, doubtless, conceived their first idea of such a weapon by being struck with the form of a Woodpecker's tongue; at least, such is my opinion."

The sexes differ but little in outward appearance, and that only in the colouring of the crown.

The male has the forehead chestnut-red; all the upper surface and wings yellowish green, tinged with ashy on the head, nape, and ear-coverts, and somewhat brighter on the edges of the secondaries; two white lines on each side of the head, one above and the other beneath the eye; primaries dark brown; two middle tail-feathers black on the outer web and white on the inner; the next four on each side entirely black; the remainder black, with whitish outer webs and tips; under surface white, tinged with yellow, and with a spot of black at the tip of each feather, which, approaching more nearly on the lower part of the abdomen, assume the appearance of interrupted bars; irides brown; bill and legs lead-colour.

In the female the forehead is green, like the other parts of the upper surface.

The Plate represents the two sexes, of the natural size.

Gould, John. 1870. "Speckled Piculet, *Vivia innominata* [Pl. 39]." *The Birds of Asia* 6(XXII), -. <https://doi.org/10.5962/p.323500>.

View This Item Online: <https://www.biodiversitylibrary.org/item/122488>

DOI: <https://doi.org/10.5962/p.323500>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323500>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: Not in copyright

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.