TROCTES MASSENÆ.

TROCTES MASSENÆ.

Massena's Trogon.

SPECIFIC CHARACTER

Mas. *Staturâ magnâ ; vertice, corpore suprâ pectoreque nitidè viridibus ; loris, plumis auricularibus et gutture nigris ; alâ medid cinerâ nigro delicatè striatâ, primariis extus albo marginatis ; rectricibus caudæ duabus intermediis purpureo-viridibus, ad apicem nigris ; reliquis nigris ; pectore corporeque subtus coccineis.*

Fœm. *Vertice, corpore suprâ, gutture, pectore abdomineque superiore intensè cæruleo-cinereis ; abdomine imo, lateribus crissoque coccineis.*

Male.—Crown of the head, all the upper surface, and chest fine green ; lores, ear-coverts, and throat black, gradually passing into the green ; centre of the wing finely pencilled with black on a grey ground ; primaries black, margined on their external webs with white ; two centre tail-feathers purplish green, tipped with black, the remainder wholly black ; breast and under surface rich scarlet ; bill yellow ; feet greyish olive.

Total length 14 inches, bill $1\frac{1}{2}$, wing $7\frac{1}{2}$, tail 7.

Female.—Crown of the head, the upper surface, throat, chest, and upper half of the belly dark bluish grey ; lower part of the belly, flanks, and under tail-coverts scarlet ; wings and tail black, the secondaries and lesser wing-coverts powdered with extremely fine spots of grey ; upper mandible black ; lower mandible yellow ; feet dark brown.

Trogon Massena, Gould, Mon. of Trog., pl. 16.—Gray and Mitch. Gen. of Birds, vol. i. p. 70, *Trogon*, sp. 19.—Gray, List of Spec. of Birds in Coll. Brit. Mus., part ii. sect. i. *Fissirostres*, p. 40.—Bonap. Consp. Gen. Av., tom. i. p. 149, *Trogon*, sp. 18.—Sclat. Proc. of Zool. Soc., 1858, p. 96, & 1859, p. 387.—Sclat. and Salv. Ibis, 1859, p. 132.—Salv. Ibis, 1861, p. 146.—Sclat. Cat. of Coll. of Amer. Birds, p. 277.—Cab. Journ. für Orn., 1862, p. 174.—Sclat. and Salv. Proc. of Zool. Soc., 1864, p. 364.—Salv. in Proc. of Zool. Soc., 1867, p. 151.

Curucujus Massena, Bonap. Consp. Vol. Zygod., p. 14. gen. 2 b. 16.

Troctes Massenæ, Cab. et Heine, Mus. Hein., Theil iv. p. 204.

ORNITHOLOGISTS will doubtless always be divided in opinion with regard to the value of the minute subdivisions which are found to exist in all large families of birds ; on observing that the members of each of these subdivisions are characterized by some peculiar feature, some will be inclined to raise them to the rank of genera, and to assign to them a distinctive designation, while others will be content with merely indicating their existence. With few exceptions, I have kept the American Trogons under two generic appellations—*Pharomacrus* and *Trogon* : one of the exceptions is *Troctes*, of which the present bird may be considered a typical example, as it certainly is the largest species known, its size being nearly equal to that of the greatest of the Train-bearers, *Pharomacri*. It is a very robust bird, and is remarkable for the total

absence of the white pectoral crescent, a character common to so many members of its family; I may remark, too, that were its habits and economy better known, we should doubtless find that its very powerful bill is fitted for some special purpose.

The *Troctes Massenæ* is very common in Guatemala, Costa Rica, and probably in all the other countries southward to the Isthmus of Panama. Specimens from the latter country are smaller than those obtained in more northern localities.

The following particulars respecting this species have been kindly furnished me by Mr. Osbert Salvin, a gentleman who is so intimately acquainted with the ornithology of Central America that all that flows from his pen must be considered of value :—" *Trogon Massenæ*, though abundant in many parts of the vast forests which clothe the warmer portions of the eastern side of the great Central-American Isthmus from Mexico to Panama, does not appear to occur in similar tracts on the western or Pacific side of those countries; nor is it, that I am aware of, to be found in the more broken and thinner forests of the central portion of the Isthmus. Though originally described as occurring in Mexico, it does not seem to be very abundant there, as the only recent records of its capture in that country are given by Mr. Sclater, who states that M. Boucard obtained specimens at Playa Vicente, and that others passed through the hands of the MM. Verreaux (*cf.* P. Z. S. 1858, p. 96, & 1859, p. 387). In Guatemala, Massena's Trogon is very abundant in the province of Vera Paz; and during my excursions to the lowland forests in 1860, and again in 1862, when Mr. Godman and I were together, I had frequent opportunities of observing it. As I said before, it is a bird which inhabits the densest forest, never emerging into the sunlight or the brushwood of an old clearing. It usually flies amongst the lower branches of the forest trees, but still far from the ground. Its habits are quick and spasmodic while in motion, and the very reverse when perched and at rest. A bird is observed to fly past overhead; it alights on a bough, and in a moment assumes an attitude that would lead one to suppose it had not stirred for an hour. When thus perched, the glittering green of the upper plumage and breast is inconspicuous, but the brilliant red of the under parts is an object of mark against the dark foliage of the trees. Its cries are various, but harsh and discordant, none of them being so soft as the call-note of the Quetzal. Its food consists principally of ripe fruits, which are plucked whilst the bird is on the wing. Occasionally a caterpillar is added to its repast. The first specimen I saw was in the mountains of Rashché, at an elevation of about 2500 feet above the sea; but at Choctum, which is only half that elevation, it occurred in numbers; not only did we obtain examples ourselves, but every collection made by natives was well stocked with them. Specimens were also procured for us near Teleman, a village situated just where the river Polochic becomes navigable for canoes. Mr. Leyland found *Trogon Massenæ* at Omoa and San Pedro, and in the notes referring to his collection (P. Z. S. 1859, p. 53) he states that 'these birds are met with in pairs; they are very active, making a chattering noise, varied occasionally by a great outcry.' In Costa Rica it appears to be equally abundant; M. Carmiol has recently forwarded us specimens from a place called Valza; and the collection of the Smithsonian Institution, at Washington, contains a plentiful supply from the same sources. A little further southwards, Enrique Arcé found it in Veragua, and has sent us skins from the Cordillera of Tolé and Santiago de Veraguas, whilst from the Panama railway line we have had forwarded to us numerous examples, by that indefatigable naturalist Mr. James McLeannan, the 'track-master' of Lion Hill Station; the forest surrounding his house seems to be full of these birds. The most southern recorded locality for the occurrence of *Trogon Massenæ* is the river Truando and the delta of the Atrato, where examples were obtained by the Messrs. Wood, who accompanied the surveying expedition under Lieut. Michler, as recorded by Mr. Cassin in the Proceedings of the Academy of Natural Sciences of Philadelphia, 1860, p. 135."

The Plate represents the two sexes, of the size of life.

Gould, John. 1869. "Massena's Trogon, *Troctes massenæ*." *A monograph of the Trogonidæ or trogons* 2, -. <https://doi.org/10.5962/p.323631>.

View This Item Online: <https://www.biodiversitylibrary.org/item/264437>

DOI: <https://doi.org/10.5962/p.323631>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323631>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.