HARPACTES HODGSONI, *Gould*.

J. Gould and H. Richter, del. & lith.

I. Walter, Imp.

HARPACTES HODGSONI, Gould.

Hodgson's Trogon.

SPECIFIC CHARACTER.

Mas. *Capite, collo, pectore corporeque subtus intense sanguineis.*

Fœm. *Capite, collo, pectore corporeque superiore cinnamomeo-brunneis.*

Male.—Head, neck, and breast deep blood-red, separated from the rich scarlet of the under surface by a crescentic mark of white; back and upper tail-coverts cinnamon-brown, brightest on the rump; wing-coverts and tertiaries striated with fine wavy lines of black and white; primaries black, margined on their outer webs with white; two centre tail-feathers rich chestnut-brown, tipped with black; the next on each side black, with two thirds, from the base, of the outer web, and a fine line down the shaft on the inner web, rich chestnut-brown; the next on each side black, with the basal half of the outer web rich chestnut-brown; the three lateral feathers on each side black, largely tipped with white, which extends some length down the outer web; “bill deep smalt-blue, becoming black along the culmen and at the tip; irides chestnut-brown; orbital skin deep lavender-blue; legs and feet pale lavender” (Jerdon).

Total length $13\frac{1}{2}$ inches, *wing* $5\frac{1}{2}$, *bill* $\frac{3}{4}$, *tail* 8.

Female.—Head, neck, breast, and upper surface cinnamon-brown, palest on the rump and upper tail-coverts; under surface pale crimson, separated from the cinnamon of the throat, as in the male, by a narrow crescent of white; undulations on the wing black and brown instead of black and white.

Trogon Hodgsonii, Gould, Mon. of Trog., pl. 34.

Harpactes Hodgsonii, Gould, Mon. of Trog., Syn. Spec. *Harpactes*, sp. 6.—McClell. in Proc. Zool. Soc., part vii. p. 166.—Gray and Mitch. Gen. of Birds, vol. i. p. 71, *Harpactes*, sp. 3.—Gray, Cat. of Spec., and Draw. of Mamm. and Birds presented to Brit. Mus. by B. H. Hodgson, Esq., p. 56.—Blyth, Cat. of Birds in Mus. Asiat. Soc. Calcutta, p. 80.—Bonap. Consp. Gen. Av. tom. i. p. 151, *Harpactes*, sp. 3.—Horsf. and Moore, Cat. of Birds in Mus. East Ind. Comp., vol. ii. p. 713.

——— *Hodgsoni*, Jerd. Birds of India, vol. i. p. 202.—Beav. in Ibis, 1865, p. 407.—Blyth, in Ibis, 1866, p. 342.

Trogon (Harpactes) Hodgsonii, Gould, Mon. of Trog., List of Plates, sp. 34.

Harpactes erythrocephalus, Gray, List of Spec. of Birds in Brit. Mus., part ii. sect. i. *Fissirostres*, p. 45.—Blyth, Cat. of Birds in Mus. Asiat. Soc. Calcutta, app. p. 322.

Pyrotrogon Hodgsoni, Bonap. Consp. Vol. Zygod., p. 14. gen. 5e. 39.—Cab. et Heine, Mus. Hein., Theil iv. p. 160.

“*Trogon dilectus*, F. B. Hamilton's MSS., vol. i. p. 63,” Horsf. and Moore, Cat. of Birds in Mus. East Ind. Comp., vol. ii. p. 713.

Suda sohaghin, Bengalese (male), Dr. Hamilton.

Hummesha Peeara ("always thirsty"), Hindoos (male), Hamilton.

Cuchcuchea, Bengalese (female), Hamilton.

Sakvor pho, Lepchas, Jerdon.

"THIS handsome Trogon," says Mr. Jerdon, "is found in the Himalayas, from Nepaul eastward, in Assam, Sylhet, Arrakan, and Tenasserim. It prefers hilly places at from 2000 to 4500 feet. At Darjeeling I found it chiefly at about 4000 feet, frequenting darkly shaded valleys and flying from tree to tree at no great elevation, or a few of them together, keeping near the same spot, making sallies every now and then, and seizing insects on the wing. It feeds on Coleoptera chiefly. Tickell, who lately observed it on the Tenasserim hills, about 3000 feet and upwards, says that it flies in small troops, is active and vociferous in the morning, solitary and quiet during the heat of the day. I had the eggs of the Trogon brought me at Darjeeling: they were said to have been taken from a hole in a tree; and were two in number, white and somewhat round. There was no nest, it was stated—only some soft scrapings of decayed wood." (Birds of India, vol. i. p. 203.) Capt. Beavan found it among heavy timber trees in the valley of Little Runjeet river, Darjeeling.

In naming the present species after B. H. Hodgson, Esq., I was actuated by a desire to pay a just compliment to a gentleman who, besides diligently fulfilling his official duties as British President in Nepaul, laboured with the greatest assiduity and with the richest result, not only in the field of natural but in many other departments of literature and science.

The Plate represents the two sexes, of the natural size. The plant is the *Benthamia fragifera*.

Gould, John. 1869. "Hodgson's Trogon, *Harpactes hodgsoni*, Gould." A *monograph of the Trogonidæ or trogons* 2, -. <https://doi.org/10.5962/p.323632>.

View This Item Online: <https://www.biodiversitylibrary.org/item/264437>

DOI: <https://doi.org/10.5962/p.323632>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323632>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.