TROGON ATRICOLLIS, Vieill.

TROGON ATRICOLLIS, Vieillot.

Black-throated Trogon.

SPECIFIC CHARACTER.

Trog. vertice, dorso pectoreque summo viridibus; tectricibus alarum et secundariis extus nigris cinereo vermiculatis; primariis extus albido limbatis; regione parotidæ guttureque nigris; torque pectorali albo indistincto; rectricibus duabus intermediis cuprescenti-viridibus, duabus proximis nigris externè viridibus, reliquis nigris albo latè terminatis albo et nigro transfasciatis.

Fcem. Capite, dorso, gutture et rectricibus sex intermediis brunneis, his nigro terminatis; rectricibus reliquis obsoletiùs fasciatis; ventre aurantiaco sordidiore.

Rostrum flavum, vix olivascens.

Adult male.—Beak yellow, clouded with olive; throat and ear-coverts black; chest, top of head, and entire upper surface green; two centre tail-feathers green, with slight bronzy reflections, the two next on each side black, with their extreme outer edges of the same green as the two middle ones, the tips of all being black, the three outer ones on each side black, tipped with white, and regularly barred on the outer web with black and white as well as towards the tip of the inner one; wing-coverts and secondaries black, finely and delicately freckled with grey; primaries black, with a narrow external edging of white; under surface fine orange, separated from the throat by an indistinct pectoral band of white; feet brown.

Female.—Head, throat, upper surface, and six middle tail-feathers brown, the latter slightly tipped with black, the remaining tail-feathers as in the male, but not so regularly barred; the wing-coverts freckled with brown on a black ground; the primaries black, margined on the outer edges with white; under surface orange, but not so pure as in the male.

Total length about 9 or 10 inches; wing $4\frac{1}{2}$; tail 6.

Yellow-bellied Green Cuckoo. Edwards, Gleanings Nat. Hist. pl. 331.

Couroucou à queue rousse de Cayenne. Buff. Pl. Enl. 736.

——— *orangé.* Levaill. H. Nat. Couroucous, p. 14, pl. 7, 8, 9.

Trogon rufus. Gm. S. N. i. p. 404.

——— *atricollis.* Vieill. N. Dict. d'Hist. Nat. viii. p. 318.—Id. Gal. des Ois. i. pl. 31.—Gould, Monogr. Trogon. pl. 8.—Gray, Hand-l. B. i. p. 83.—Scl. & Salv. Nomencl. Av. Neotr. p. 104.

——— *viridis.* Spix, Av. Bras. i. p. 50 (*nec* Linn.).

Trogonurus lepturus. Swains. An. in Menag. p. 331.

Trogon chrysochlorus. Natt. & Pelz. Sitz. Akad. Wien, xx. p. 49.—Pelz. Orn. Bras. p. 20.

——— *scalaris.* Licht. in Mus. Berol. (*fide* Cabanis).

Pothinus atricollis. Cab. & Heine, Mus. Hein. Th. iv. p. 180.

“THERE can be little doubt,” says Mr. Salvin, “that the figures of the old authors, Edwards, Buffon, and Levaillant, quoted above, all apply to this bird, and also that a strict application of the law of priority points to the exceedingly inappropriate title of *Trogon rufus*, bestowed upon it by Gmelin, as the name this bird ought to bear.

"Setting aside the name of *T. rufus* as inappropriate, and *T. lepturus*, proposed by Swainson, which is a pure synonym of *T. atricollis*, as well as the MS. title of Lichtenstein's quoted by Dr. Cabanis, the other names given to this bird all refer to races or supposed races occurring in different districts of its range. Thus the South-Brazilian bird has been separated by Herr von Pelzeln as *T. chrysochlorus*, and the Central-American as *T. tenellus* by Dr. Cabanis, Vieillot's name, *T. atricollis* being applicable to the race found in Guiana and the valley of the Amazons.

"Having on several occasions, both by myself and with Mr. Sclater, closely examined numerous specimens of this bird from all parts of its range with a view to test the right of these various races to rank as species, I have always come to the conclusion that such a rank could not be granted them without rendering it exceedingly difficult, if not impossible, to assign to every specimen its right name. To show how varied is the state of opinion on this point, I may add that Dr. Cabanis, Mr. Gould, and Mr. Lawrence, uphold the distinctness of *T. tenellus*, the two former rejecting *T. chrysochlorus* as distinct from *T. atricollis*. Dr. Finsch agrees with Herr von Pelzeln in maintaining the specific rank of *T. chrysochlorus*, suggesting (as it appears to me rightly) that the birds described in the 'Museum Heineanum' really were *T. chrysochlorus* and not *T. atricollis*. I think myself that the differences observable are so slight that they may be regarded rather as tendencies to vary in a particular direction in a particular locality than as fixed, definite, specific characters. At the same time I must say that, if their separation be maintained, the South-Brazilian has equal, if not stronger, claims to be considered distinct from the Guiana race than the Central-American has. The Guiana bird is in fact almost precisely intermediate between the races found in Brazil and Central America. The white bands of the tail are widest and fewest in *T. tenellus*, narrowest and most numerous in *T. chrysochlorus*. The central tail-feathers have a bluish shade in *T. tenellus*, and are most bronzy in *T. chrysochlorus*. The wing-coverts are coarsely mottled in the former, finely in the latter. In all these points *T. atricollis* is intermediate."

Leaving the northern form, *T. tenellus*, to be noticed in a separate article. I will now trace the range of *T. atricollis*, including with it *T. chrysochlorus*.

Buffon alludes to its occurrence in Cayenne. It also occurs in Demerara, where Schomburgk met with it; but westwards of this district I have not yet come across any recent allusion to its occurrence in Venezuela or Trinidad, though Levaillant mentions its presence in the latter island. Throughout the valley of the Amazons it is found apparently in abundance. Natterer obtained nine examples during his voyage at Nas Lages, Borba, Marabitanas, Cocuy, and Barra do Rio Negro; while on the Upper Amazons, or Solimoes, Spix met with it, and subsequently Mr. E. Bartlett within the frontiers of Peru at Chamicurros. In Southern Brazil Prince Maximilian found it at Bahia, whence, and also from Rio de Janeiro, numerous specimens are constantly sent to Europe. In the province of Sao Paulo Mr. J. F. Hamilton found it pretty generally distributed throughout the forest, but not abundant in any one locality. He adds that it is of rather solitary habits, and that he never saw two together. In the same district, at Ypanema, Natterer secured fifteen examples, and upon these he bestowed the name *chrysochlorus*, afterwards described by Herr von Pelzeln. *Trogon atricollis* is also found still further south, in the province of Rio Grande do Sul, whence Mr. Rodgers has sent us specimens.

The figures in the Plate are of the natural size.

Gould, John. 1875. "Black-throated Trogon, *Trogon atricollis*, Vieillot." A *monograph of the Trogonidæ or trogons* 4, -. <https://doi.org/10.5962/p.323651>.

View This Item Online: <https://www.biodiversitylibrary.org/item/267606>

DOI: <https://doi.org/10.5962/p.323651>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323651>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.