MILVUS MIGRANS, Bodd.

MILVUS MIGRANS.

Black Kite.

Falco migrans, Bodd. Tab. de Pl. Enl., p. 28, no. 472.

Accipiter milvus, Pall. Zoog. Rosso-Asiat., tom. i. p. 356.

Falco ater et *F. austriacus*, Gmel. edit. Linn. Syst. Nat., tom. i. p. 262.

— *fuscus-ater*, Meyer, Taschenb. Deutschl. Vög., tom. i. p. 27, et tom. viii. p. 11.

— *ater* et *F. fuscus*, Brehm, Vög. Deutschl., p. 53.

Milvus niger, Briss. Orn., tom. i. p. 413.

— *austriacus* et *F. ater*, Daud. Orn., tom. ii. p. 149.

Hydroictinia atra, Kaup, Class. der Säug. u. Vög., p. 115.

Milvus (Hydroictinia) migrans, Gray, Hand-list of Birds, p. 26.

IN 1867 Mr. Hancock made known the circumstance of an individual of this species having been killed in Northumberland; and almost simultaneously a specimen was transmitted to me from Northern Australia,—facts which will give the reader an idea of how widely this species ranges over the globe. That there might be no mistake in the matter, I submitted the Australian bird to the inspection of J. H. Gurney, Esq., who immediately said it was identical with the European *Milvus migrans*; and Mr. Hancock's testimony will, I am sure, be deemed sufficient as to the identity of the British-killed or Northumberland example. Either as a bird of passage or as a migrant this species is said to inhabit most of the central portions of Europe, Asia Minor, and almost the whole of Africa; we also find it in the lists of the birds of many other countries.

"The geographical distribution of the Black Kite" says Professor Newton, in his edition of Yarrell's 'British Birds,' is extensive. Though not found in Norway, Sweden, or Finland, in Russia it reaches as far to the north as Archangel, and thence across Siberia, becoming rarer to the eastward, and hardly observed beyond the Lena. It is said by Pallas to winter in Persia, where De Philippi also found it. It is very common in the Caucasus; and Messrs. Dickson and Ross obtained it at Erzeroum. In Palestine, according to Canon Tristram, it arrives about the beginning of March in immense numbers, and scatters itself over the whole country. There is much discrepancy in the accounts of recent travellers as to its occurrence in Egypt, some stating that it is very abundant there, and others that they never met with it, and that a near ally (*Milvus ægyptius*) must have been mistaken for it. The explanation of the difficulty probably lies in the fact that, while *M. ægyptius* is a resident in Egypt, *M. migrans* is a bird of passage only and may not always stop for the convenience of other travellers on its way down or up the Nile valley. Drs. Von Heuglin and E. A. Brehm include it as a bird of Eastern Kordofan and Abyssinia and Mr. Blanford; found it to be extremely common both in the highlands and lowlands of the latter country. Mr. Chapman sent specimens procured on the Zambesi to Mr. Layard; and Mr. Edward Newton shot a bird, pronounced by Mr. Gurney to be of this species, in Madagascar. Mr. Layard also records an example killed at Colesberg, in the Cape Colony; and Andersson met with it in Damara Land, where it arrives in autumn in large numbers, and remains throughout the breeding-season. In Western Africa it has been obtained at Bissao and on the Niger. It occurs in Morocco, and is very common in Algeria, breeding in the Atlas, but not occurring to the south of that range of mountains. In Europe it is said to be met with occasionally in Portugal and in Spain, where it breeds, as it also does in several parts of France. It does not seem to have occurred in Belgium; but the Leyden Museum contains a specimen killed in Holland."

With respect to its solitary occurrence in England Mr. Hancock says (in 'The Ibis' for 1867, p. 253):—

"A fine mature male example of the Black Kite, *Milvus migrans*, came into my possession in a fresh state on the 11th of May, 1866. It was taken in a trap by Mr. F. Fulger, the Duke of Northumberland's gamekeeper, a few days before, in the Red-deer park at Alnwick. This is, I believe, the first time that this fine rapacious bird has occurred in Britain. The plumage was in very good condition, except on the lower part of the body (where it had sustained some injury from the trap), and agrees with that of mature specimens, in my collection, which I received from the Continent some years ago. It was proved by dissection to be a male."

Throughout the whole of the countries embraced in its wide range the Black Kite is migratory, proceeding northward in spring and returning southward in autumn—thereby fully meriting the earliest appellation, that of *migrans*, bestowed upon it by Boddaert.

Mr. Salvin, writing of the bird as seen by him in the Eastern Atlas, says, in 'The Ibis' for 1859, p. 184:—

"During the breeding-season it is much more abundant in the Souk-Harras district than *M. regalis*.

Indeed, with the single exception of the Kestrel (*Tinnunculus alaudarius*), it is the commonest rapacious bird in the Eastern Atlas. Like *M. regalis* it selects for the position of its nest the roots of a shrub growing out of a rock, and builds a structure composed principally of sticks, with a lining of rags, wool, &c., while on the surrounding branches are fantastically hung old pieces of burnouses of various colours. The Black Kite plays the part of scavenger in the districts where it abounds; and over every French settlement and Arab village several may be seen flying boldly around, on the look-out for any fragment of carrion that may be lying about. Its fearless and familiar manner and beautiful flight render this bird decidedly one of the most interesting in the country. The eggs are laid from the middle to the end of April, and are more distinctly and deeply marked than those of the other species. A series of the eggs of the two from the same district present a marked contrast. This bird is known to the Arabs by the name 'Hadayia soda'—the 'Black Hadayia'.

Of the Black Kite in Palestine, Tristram remarks (in 'The Ibis' for 1865, p. 256):—"No sooner has the Red Kite begun to retire northwards than the Black Kite, never once seen in winter, returns in immense numbers from the south, and about the beginning of March scatters itself over the whole country, preferring especially the neighbourhood of villages, where it is a welcome and unmolested guest, and certainly does not appear to attack the poultry, among which it may often be seen feeding on garbage. It is not strictly gregarious, though very sociable; and the slaughter of a sheep near the tents will soon attract a large party of Kites, which swoop down, regardless of man and guns, and enjoy a noisy scramble for the refuse, chasing each other in a laughable fashion, and sometimes enabling the wily Raven to steal off with the coveted morsel during their contention. It is the butt of all the smaller scavengers; and it is evidently most unpopular with the Crows and Daws, and even with the Rollers, who enjoy the amusement of teasing it in their tumbling flight, which is a manœuvre most perplexing to the Kite. The nest, generally in a tree, often in a glen, is a grotesque, untidy structure, decorated with all sorts of rags and rubbish, apparently to attract observation. The eggs are invariably two, and, as a rule, are more richly coloured than those of the *Milvus regalis*."

"This species, which does not remain [in Southern Spain] throughout the winter," says Mr. Howard Saunders (in 'The Ibis' for 1871), "breeds a full fortnight later than *M. regalis*; and the first week in May a patch of wood of an acre or two in extent will certainly yield upwards of a score of eggs, it being a sociable bird. Even when nesting somewhat apart, it has always a colony of Spanish Sparrows to keep it company. The usual number of eggs is two."

This association of the Spanish Sparrows with the Black Kite is also mentioned by Lord Lilford, who says (in 'The Ibis' for 1866):—"The Black Kite's nest contained three eggs; and in its foundation were three nests of the Spanish Sparrow. In almost every nest of the two species of Kite we find in Spain there were one or more nests of this Sparrow, besides, in most instances, a large colony in the immediate vicinity."

Messrs. Elwes and Buckley, in their 'List of the Birds of Turkey,' published in 'The Ibis' for 1870, state that the Black Kite is "extremely common on some parts of the Danube, where it breeds on the islands, which are covered with a dense thicket of willows and a few poplars. In these trees it begins to build in April, and lays about the first week in May—in a very small nest, which, at first sight, would not seem large enough for the eggs of a Crow. It is also found in the interior of the country, and in the large towns, where it acts as a scavenger."

Respecting the *Milvus migrans* in South Africa, Dr. Exton wrote to Mr. Layard:—"This Kite is said by the Matabili to be 'the king's bird,' and is in consequence much respected by them. A chief's son examining my specimen said, 'We never kill that bird.'

"It is remarkably bold and fearless, dashing down at your very feet for a stray scrap of flesh, or attempting to carry off meat hung up to dry in the native fashion. It does not seem very choice in its food. The stomachs of those I examined contained locusts and lizards; and I have seen family parties dining, after the manner of Vultures, off the putrid carcass of an ox. They breed about the time the locust-larvæ become developed, the young birds taking wing when the 'hoppers' are becoming strong on the ground. They then congregate in flocks; and I have counted between eighty and ninety hovering over an army of infant locusts—and have seen them in still greater numbers, some on the ground busily devouring the 'hoppers,' others perched on the neighbouring trees gorged with a full repast."—*Ibis*, 1869, p. 362.

The figure is about three fourths of the natural size.

Gould, John. 1873. "Black Kite, *Milvus migrans* [Pl. 23]." *The birds of Great Britain* 1, -. <https://doi.org/10.5962/p.323803>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221495>

DOI: <https://doi.org/10.5962/p.323803>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323803>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.