MUSCICAPA COLLARIS, Bechst.

MUSCICAPA COLLARIS, *Bechst.*

White-collared Flycatcher.

Muscicapa collaris, Bechst. Orn. Taschenb., tom. i. p. 158.

——— *albicollis*, Temm. Man. d'Orn., 2nd edit., tom. i. p. 153, et tom. iii. p. 84.

——— *streptophora*, Vieill. Faun. Franç., p. 145, pl. 151. figs. 2, 3.

——— *atricapilla*, var. β , Gmel. edit. Syst. Nat., tom. i. p. 835.

——— (*Hedymela*) *collaris*, G. R. Gray, Hand-l. of Birds, part i. p. 322.

——— *albifrons* et *albicollis*, Brehm, Vög. Deutschl., p. 223.

I CAN furnish no evidence of this bird having been killed in England beyond what is stated in my 'Birds of Europe;' but I recollect perfectly seeing a specimen in the flesh at the house of the late Mr. Leadbeater in Brewer Street, which, as far as my memory serves, was destined for the collection of the late Mr. Lombe, of Norfolk, who was very particular that it should contain as many British-killed specimens as possible; and that the specimen I saw at Mr. Leadbeater's went to Mr. Lombe is almost certain from the following note lately received from Mr. J. H. Gurney, jun.:—"In your 'Birds of Europe' you state that you saw the White-collared Flycatcher in a collection of British birds, on which authority Jenyns and Doubleday introduced it doubtfully. I have just detected an adult male in the late Mr. Lombe's collection. It is one of the few birds marked in the catalogue as having been received in a fresh state. Mr. Lombe got his best birds of Leadbeater; and at his death they were moved from Great Melton to Wymondham, where his daughter (Mrs. Clark) keeps them."

The circumstance of a White-collared Flycatcher having been killed in England need not excite surprise, inasmuch as it is as plentiful on the Continent as the Pied Flycatcher (*Muscicapa atricapilla*). Both are migratory, spending the summer in the northern and central parts of Europe, and retiring southward to warmer climates in winter. When the *Muscicapa collaris* becomes better known to our living ornithologists and field-naturalists it is not improbable that we shall hear of its occasionally occurring on our island. Although the two birds are found on comparison to be very different, each possessing characters sufficiently distinct to warrant their specific separation, those characters are not so easily seen in the living birds among the foliage of trees; and hence the rarer species may have escaped notice.

As already mentioned, the White-collared Flycatcher is abundant in the central provinces of continental Europe, it is also somewhat plentiful in various parts of France, but less so in the northern portion of that country. Degland says he has found it at Lisle in May, and that it breeds in considerable numbers in Lorraine. Temminck says that it is never found in Holland, and but seldom in Central Italy. Count von der Mühle observed it in Greece for a few days during its spring passage in April, in breeding-plumage and in considerable numbers, whence he thinks it probable that it breeds in Roumelia. Captain Loche includes it in his list of the birds of Algeria. The best account of the bird and its habits that I find recorded is comprised in the following somewhat free translation of some passages in Bailly's 'Ornithologie de la Savoie':—

"The White-collared Flycatcher is much less common in Savoy than the *Muscicapa atricapilla*, even at the time of its two migrations, which take place from the 8th to the 15th of April, and at the end of August or the commencement of September. It is at the latter season that it is most abundant, particularly if warm rains are prevalent; it then remains a little longer than usual with us, no doubt in consequence of the abundance of insects that occur under those circumstances, and are so necessary for its subsistence. In spring it arrives in pairs, rather than singly as at the end of summer. Some pairs breed in the northern part of our territory, particularly in those districts rendered humid by the thick forests of Chamounix, Maurienne, Tarentaise, and the environs of Albertville.

"Both sexes unite in the construction of the nest towards the middle of May. It is formed with moss, lichens, blades of grass, and straw and flexible roots, lined with hair, fine grass, feathers, and down. It is negligently constructed, and is placed in a perforated tree, in a cavity abandoned by a Woodpecker or Tit, for which these birds often dispute the ownership with the Nuthatch and Wryneck. The eggs, which are four or five in number, rarely six, are sometimes nearly round, but ordinarily are pointed, and of a uniform bluish tint, but are occasionally washed with green or a light greenish brown. The young are fed by both parents with small winged insects, for which they diligently search. They attend to their young

for some time after they have left the nest, and do not quit them until they are perfectly able to take care of themselves.

"The Collared Flycatcher does not resort so frequently to the tops of trees as the *Muscicapa grisola*, even during fine weather, but evinces a preference for the centre and lower portions of trees and shrubs, and bushes on the borders of ditches and rivers, where insects are particularly numerous during the heat of the day. In rainy weather it remains on the lower branches, whence it readily darts upon any insect that may be passing or fluttering near the ground. When obliged to resort to the ground for food, it alights on a stone, or any other slight elevation, and waits for its prey. Its actions are very similar to those of *M. grisola*; and it exhibits a similar air of stupidity and melancholy. Like that species it admits of a close approach. Its voice is very different, stronger, sharper, and seems to resemble the syllables *pitt, pitt*. During the breeding-season and the education of the young the male gives utterance to a warble as brief as that of *M. atricapilla*, mainly composed of sharp notes, but not entirely devoid of sweetness. In autumn, when it is fat, its flesh is not to be despised, especially as at that season the bird feeds upon fruits, such as the mulberry and fig.

"This species departs from Savoy before the commencement of the month of October; but a few of the young remain to the 8th or 12th of that month."

Besides possessing a broad white collar at the back of the neck, this species differs from *M. atricapilla* by having a much greater amount of white on the forehead and a more extensive band of white on the primaries. Temminck states that the tail is wholly black; and such is the case with the specimen now before me; but I suspect that in some specimens the outer tail-feathers are margined with white. The female, as will be seen on reference to the Plate, differs very considerably from the male. It is said that the young males are very similar to the female, or, if any difference occurs, they are rather darker. "On the approach of spring the plumage of the young male blackens wherever it is ash-coloured in the female. One or both lateral tail-feathers are black, with white edges, but the latter disappear entirely in males upwards of two years old. In winter there is no difference between males and females" (BREE).

The Plate represents an adult male and either a female or a young male, of the size of life. The plant is the *Ulmus major*, with its seed-vessels.

Gould, John. 1873. "White-collared Flycatcher, *Muscicapa collaris* [Pl. 18]." *The birds of Great Britain* 2, -. <https://doi.org/10.5962/p.323833>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221554>

DOI: <https://doi.org/10.5962/p.323833>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323833>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.