BUDYTES CINEREOCAPILLA.

BUDYTES CINEREOCAPILLA.

Grey-capped Wagtail.

- Motacilla cinereo-capilla*, Savi, Nuov. Giorn. Lett., no. 57, p. 190.—Idem, Orn. Tosc., tom. iii. p. 216.
——— *borealis*, var. β , Blas. List of Birds of Eur., Eng. edit., p. 12.
——— *flava cinereocapilla*, Schleg. Rev. Crit. des Ois. d'Eur., p. xxxviii.
——— *Feldeggi*, Mich.
——— *dalmatica*, Bruch.
——— *melanocephala*, Savi (nec Licht.), summer dress.
Budytes cinereocapilla, Bonap. Consp. Gen. Av., tom. i. p. 249, *Budytes*, sp. 2.
——— *nigricapilla*, Bonap. ibid., *Budytes*, sp. 4?
-

A VERY beautiful specimen of this Wagtail having been killed in Cornwall, by Mr. Vingoe, of Penzance, it becomes necessary to give a figure and an account of the species in the 'Birds of Great Britain,' which I am happy to do, inasmuch as it affords me an opportunity of giving some further information respecting the puzzling members of the genus *Budytes*. At present their synonymy is in the utmost confusion; and the opinions of ornithologists as to which is and which is not entitled to rank as a species are equally conflicting, some believing that several of the so-called species are identical, while others are disposed to separate them still further than has yet been done; and it is not until numerous specimens of both sexes from all the different countries frequented by these birds are brought together, placed side by side, and carefully compared, that the vexed question as to the number of species which really exist can be satisfactorily determined. It is even a question with me whether the bird I have figured under the name of *Budytes flava* be really the true *Motacilla flava* of Linnæus, or if the one here represented be not the bird so named by the illustrious Swede. If this should ultimately prove to be the case, then my specific name of *neglecta* must be restored, and the above list of names sink into synonyms. The confusion in the synonymy and the conflicting opinions of ornithologists have in a great measure arisen from the circumstance of all the species being nearly of the same size, and from the females and young being very similar in their colouring. To define each species unerringly it becomes necessary to compare the fully adult males of all with each other; and if this be carefully done, I venture to assert that it will be an easy matter to distinguish the various members of this elegant group of birds, and to prove that they comprise even more than have yet been characterized. To affirm for one moment that the yellow-headed *Budytes Rayi* of our island and the black-headed *B. melanocephala* (Licht.) of the Lower Nile, the Crimea, and Asia Minor are identical, is to my mind simply absurd. After a careful study of these Wagtails during the last forty years, I believe I may confidently state that four of them pass the summer in Europe—and that these are the *Budytes Rayi* (which at that season frequents the British Islands), the *B. flava* (vel *B. neglecta*) of Western Europe and Sweden (?), the *B. cinereocapilla* of Southern and Central Europe (which migrates to Norway in spring for the purpose of breeding), and, lastly, the *B. melanocephala*, Licht. (distinguished by its jet-black head and deep yellow breast and throat), of the eastern portions of Europe, particularly the Crimea, Asia Minor, and the Nile valley.

It would, perhaps, be out of place to enumerate here the species of the genus *Budytes* which are extra-European; but I may mention that birds of this form inhabit most parts of the Old World. Unless we except the assertion that a *Budytes* has been discovered in Alaska, none are found in America; nor is there any bird therein which might be regarded as representing these lively little birds. In India and China there are more species than in Europe, all of which, I believe, are different from the European.

Professor Schlegel appears to consider that the *Budytes cinereocapilla* is confined to Italy; but I can assure him that I both saw and shot numerous examples on the Dovre-fjeld, in Norway. Some of these were much darker than others, and, moreover, had a dark patch on the centre of the breast, which appeared to be due to the abrasion of the yellow portion of the feathers having rendered their black bases visible.

Respecting the specimen killed in Cornwall Mr. Rodd informs me that "it was observed for some days about the place where it was at last captured, viz. amongst some loose cabbage-plants on the ridge of bank-sand which separates the eastern marsh from the sea-beach, between Penzance and Marazion, very near the Marazion Railway Station. It was in the month of May. It uttered a very neat and melodious song, which, considering the time of the year, may be regarded as a love-song; but no mate was seen at the time."

Of all modern authors, it appears to me that Bonaparte is the one who has taken the most correct view

of these grey and black-headed Wagtails, for which I must refer my readers to his valuable 'Iconografia della Fauna Italica.'

On reference to my MSS. I find the following note respecting this bird as observed by me in Norway, written while the matter was fresh in my memory. "From a careful examination of several specimens obtained on the Dovre-fjeld and in its neighbourhood, I am inclined to regard *Budytes cinereocapilla* and *B. nigricapilla*, Bonap. (*M. melanocephala*, Savi., nec Lichtenstein) as one and the same species. Males killed on approaching the Dovre-fjeld had dark grey heads, while those obtained near Jerkin had jet-black crowns. The dark-grey-headed birds had scarcely a trace of the stripe through the eye; the females, on the contrary, had the head olive-grey, a conspicuous stripe through the eye, and an indistinct mark of black on the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-fjeld, where the bird may be seen in all its glory from the middle of June to the middle of July on all the hill-sides to about or even above the line of vegetation. The crowns of those killed were quite black; they had no stripe through the eye; upper part of the throat beneath the bill white; the remainder of the under surface rich deep yellow, with a conspicuous patch of black on the chest; their backs were fine green, their bills black, their feet brownish black, and their eyes black. They were fine sprightly birds, and uttered notes quite different from those of the Yellow Wagtail of England."

As it is almost impossible to determine to which species the numerous notices that have appeared under the heading of *Budytes cinereo-capilla* have reference, I hesitate to quote any of them, with the exception of the following, from the pen of Lord Lilford, which, doubtless, pertains to the present bird. He states (in 'The Ibis' for 1860) that it "arrives in great numbers in Corfu about the middle of April, at which season it is to be found in small flocks in all the low meadows and maize-fields of the island. I never could find a nest of this species, though I have observed a few pairs during the whole summer."

In conclusion, I may remark that the colouring and markings of the head of the male occasionally present slight differences, some specimens having the crown, lores, and ear-coverts black, others uniform dark sooty grey; and in others, again, the grey is broken by a faint white stripe above the ear-coverts, as seen in the Penzance example.

I must not omit to acknowledge the persevering kindness of Dr. Leith Adams, than whom no one of my friends rendered me more valuable assistance towards the illustration of these Wagtails. During his sojourn in Malta he collected for me a very large series, obtained during their passage from Africa to Europe in the spring, and on their return in autumn, at which latter season the young either accompanied the adults or followed them after a short interval. No examples of either the western *Budytes Rayi* or the eastern *B. melanocephala* were found among the birds I had the pleasure of receiving from Dr. Adams, the only species being *B. cinereocapilla* and *B. flava*.

The upper figure in the accompanying Plate represents the Penzance specimen; the two lower ones are from male and female examples shot by myself on the Dovre-fjeld, in Norway, on the 4th of July, 1856.

Gould, John. 1873. "Grey-capped Wagtail, *Budytes cinereocapilla* [Pl. 5]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323897>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323897>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323897>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.