ANTHUS ARBOREUS. *Bechst.*

ANTHUS ARBOREUS, *Bechst.*

Tree-Pipit.

? *Alauda trivialis*, Linn. Syst. Nat., tom. i. p. 288.

—— *minor*, Mont. Orn. Dict.

Anthus arboreus, Bechst. Naturg. Deutschl., tom. iii. p. 706.

Motacilla spipola, Pall. Zoog. Ross.-Asiat., tom. i. p. 512.

Alauda turdinae, Scop.

Anthus trivialis, Flem. Brit. Anim., p. 75.

Pipastes arboreus, Kaup, Natiirl. Syst., p. 33.

Anthus foliorum, Brehm, Vög. Deutschl., p. 326.

—— *juncorum*, Brehm, ib., p. 326.

—— *herbarum*, Brehm, ib., p. 327.

I WISH I could imbue the minds of all my readers with the same degree of love for our native birds that reigns in my own, or could set forth in clear and truthful language the many charms of this comparatively plain-plumaged little bird; but as I am unable to do this to the extent or in the manner I desire, I must content myself with penning what I may to this end, and leave the accompanying illustration to show what the bird is like. First, then, I may state that the Tree-Pipit, although abundantly dispersed over every part of England and the whole of Scotland, except in the extreme north, is only a summer visitant to these islands; for as truly as the Swallow leaves us in the autumn for warmer climes, so does the Tree-Pipit. It will not even await the chilly blasts of autumn before it wings its way across the seas to the coasts of France, Spain, and Portugal, whence it proceeds by slow degrees towards the shores of the Mediterranean, which it also crosses for the more genial latitudes of the south. Here it awaits the turning of the northern portion of our planet towards the sun, and at the proper time profits by the warmth of that luminary and returns to us in April or the beginning of May, when the leaves are expanding in our woodlands and the elms in the hedgerows, when the wheat and the rye are carpeting the fields with the maiden green of their first sheathings, when the cowslip and the buttercup bespangle the mead; for it is then that the country is fitted for this bird. How great a difference is observable in the habits and economies of the Tree- and the Meadow-Pipit, species so closely assimilating in general appearance that even the ornithologist has sometimes to look twice before he can say with certainty which is which. There are, however, characters in their plumage, particularly the strongly marked edgings of the wing-coverts of the Tree-Pipit, which serve to set the matter at rest; or if we wish for a more convincing proof, an examination of the claw of the hind toe will furnish an infallible one, that of the Meadow-Pipit being long, straight, and slender, while the Tree-Pipit's is short, curved, and comparatively thick.

The *Anthus arboreus*, as its name implies, spends much of its time on trees, but only on those which skirt open plains and border fields of corn or the grassy glades of the forest. Here the male usually sits and pours forth his pretty simple song; but when the female is incubating, he mounts in the air, and warbles over his strain on quivering wings as he descends to the branch whence he started, or to the ground—a movement which is repeated again and again, as if in the exuberance of joy. The Meadow-Pipit, which, on the other hand, is a stationary species, is one of the hardiest of our little birds; for let the frost be ever so nipping, and the cold so great as to turn the sturdiest of us away from the windy hillsides, the Meadow-Pipit is there, a stone wall or turf dyke being its highest resting-place.

I have purposely digressed from the more immediate subject of the present paper, in order to show that two birds closely assimilating in size, colour, and structure, are very different in their habits and economy. These differences have always been noticed by ornithologists, and have induced some of them to place them in distinct genera; in the present work, however, I keep all our Pipits in the genus *Anthus*; were I not to do so every species would have a different generic name.

Baillly, in his work on the 'Ornithology of Savoy,' has given a long and somewhat graphic history of the Tree-Pipit; but as much of his account is very similar to what I have already written, it will not be desirable to repeat it here. In that country, as with us, the bird is a migrant, arriving in May and retiring southwards again in August, September, and October. A few remain to breed, both in the lowlands and in the mountains; but they do not ascend to any great height, a low temperature appearing uncongenial to them. According to Baillly the Tree-Pipit is highly prized by epicures and "wine-connoisseurs" as an article of food, on account of the delicate flavour of its flesh and its extreme fatness, which is probably due to the abundance of food it finds among the vines and figs, to which it is doubtless attracted by the insects

abounding there, rather than by the fruit, which, I believe, it never eats. Being constantly seen in such situations, it is known by the names of *Vinette*, *Becfigue*, and *Becfigue de Vignes*. In Savoy it plays a very important part as an article of diet, and it is there captured or killed in various ways, principally by a kind of clap-net; but the gun is sometimes employed, and fifteen or twenty may be shot at one discharge while resting during their autumnal migration; with us, however, it is never seen in such abundance, and I question if the cost of powder and shot required to procure a dish of Tree-Pipits would not be too great to render it remunerative, or if the bird-catcher would be compensated for his time, patience, and trouble. In the way of eating all kinds of little birds, we and our continental brethren are very different people; and I can easily imagine, from what I have heard of strings of Robins and Wrens being exposed for sale in the markets of Rome and other places, that the present bird with its rolls of fat would form for them a *bonne bouche* not to be despised.

Hitherto I have only spoken of this bird as frequenting England, Scotland, and a portion of Europe; its range, however, is far more extensive, since it has been observed all over the central portions of the continent, Madeira, North Africa, Palestine, Persia, and some parts of India: further west than this it would not be safe to assign it a locality, since the species of this form inhabiting China and Japan is, I believe, the *Pipastes agilis*. As is the case with the other members of this form, the sexes are alike in colouring, and the young at a month old differ but little from the first autumn moult of their parents.

As a songster the Tree-Pipit is in some respects unsurpassed by any British bird. Though its notes may perhaps be deficient in the softer modulations which characterize those of the Blackcap and the Redbreast, in power they exceed those of the Nightingale itself. Its joyous outburst rings through the surrounding groves in a manner almost astonishing even to those most accustomed to it, while they make a stranger suspect that he is listening to the tones of some truant Canary. The bird seems to delight in singing during the sunny intervals of showery weather. At such times it may be observed springing into the air and, while on the wing, with quaint gestures, pouring forth its loud triumphant carol, which suddenly ceases as it alights on some neighbouring tree.

Notwithstanding the curved form of its short hind claw, it runs over the ground with great agility; at the same time it is less shy and recluse than its congeners, is more tame in disposition, and displays the elegance of its form by exposing itself readily to the view of the spectator while lightly perched on some bare branch of a tree, the end of a vine-pole, or the crown of a hillock, a stone, or any conspicuous elevation.

The nest is a slight structure, placed with little artifice, in a depression on the side of a bank or among the grasses in the open field or shrubbery, frequently under the shelter of a bush or tuft of herbage. With regard to its eggs, Mr. Hewitson remarks that there are none "which present so many, or such distinct varieties as those of the Tree-Pipit. No one would at first believe them to be eggs of the same species; and it was not until I had captured the bird upon each of the varieties, and also received them from Mr. H. Doubleday similarly attested, that I felt satisfactorily convinced upon the subject. All are easily procured." The most common kind are somewhat smaller than the others, and of a deep reddish hue sprinkled with fine dots of a livelier tint; the others are of as many shades of pale lilac, purplish, or buffy red, more or less minutely spotted, and blotched with darker tints of the same hue.

"The nest of this species," adds Mr. Hewitson, "is composed chiefly of dry grass mixed with moss, lined with finer grasses, and sometimes with a few hairs. It is placed on the ground, rarely far distant from trees or brushwood, and is frequently found in woods and plantations by the side of a drive or footpath. The favourite resort of the Tree-Pipit is a grassy bank on the margin of a wood, especially if ornamented by a few single trees, on the branches of which it delights to perch. In such a situation you are sure to find its nest and eggs towards the end of May or during the month of June. The eggs are four or five in number."

Feathers of the head and all the upper surface brownish black, broadly margined with reddish grey; wing-coverts brownish black, tipped with greyish white; primaries deep brown; outer tail-feathers whitish, with a broad stripe of dark brown occupying the basal three-fourths of the margin of the inner web, the next on each side dark brown, with a deeply indented mark of whitish at the tip of the inner web, the remainder wholly dark brown; above the eye a streak of buff; ear-coverts brown; under surface fawn-colour, becoming paler on the chin and centre of the abdomen, and having a series of dark-brown streaks down the sides of the neck, across the breast, and on the flanks; bill fleshy brown, lighter on the under mandible; legs and feet fleshy white.

The Plate represents the two sexes, of the natural size, on the white poplar (*Populus alba*, Linn.).

Gould, John. 1873. "Tree-Pipit, *Anthus arboreus* [Pl. 14]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323905>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323905>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323905>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.