EMBERIZA CITRINELLA, Linn.

EMBERIZA CITRINELLA, *Linn.*

Yellowhammer or Yellow Bunting.

Emberiza citrinella, Linn. Faun. Suec., p. 84.

——— *sylvestris*, Brehm, Vög. Deutschl., p. 294.

——— *septentrionalis*, Brehm, ib., p. 294.

Citrinella septentrionalis, Kaup, Natürl. Syst. p. 142.

THE present beautiful bird is known throughout the greater portion of the British Islands by the name of Yellowhammer, a term which Yarrell considered to be a corruption of Yellow Ammer, the word "ammer" being a common and well-known German name for many of the Buntings; he has therefore called this species the Yellow Bunting or Yellow Ammer. The Scotch biographer of our native birds, Macgillivray, also described it as the Yellow Bunting, and adds the following list of provincial names as applied to it:—Yellowhammer, Yellow Yeldring or Yoldring, Yellow Yowley, Yellow Yite, Yoldring, or Yeldrock, Skite, Devil's bird, Buidhein, Buidheag Bhuachair; while Thompson, of Ireland, mentions only the following:—Yellow Bunting, Yellow Ammer, and Yellow Yorlin. Of all these terms that of Yellow Bunting is undoubtedly the most correct; but Yellowhammer is the one by which it is generally known to the school-, the herd-, and the plough-boy. Like those sturdy sons of the soil, it is strictly a native of, and a permanent resident in, this country (for it never leaves us either in summer or winter), and is alike common in every district, from north to south and from east to west, from the low fluviatile county of Lincolnshire to the high peaks of Derbyshire, from the Lothians to the hills of Rosshire.

In the early mornings the Yellowhammer may be seen on the dew-bespangled sprays of the field-side, and there, while perched on some prominent twig, emits his singular ditty long before the vernal migrants have arrived. As summer advances, no bird is more showy, nor is there one whose appearance is more striking among the hedges which skirt the green lanes; the dense coppice and the thick wood he shuns; it is among the wide shaws in low valleys, and the bushes which grow on commons and wastes, that the Yellowhammer finds a home congenial to his tastes. It has now paired, and the couples only wait for the thorn-bush to be covered with leaves, and the ditch-side overgrown with grasses and herbage, before they commence their nests.

"Just by the wooden bridge a bird flew up
Seen by the cow-boy as he scrambled down
To reach the misty dewberry. Let us stoop
And seek its nest. The brook we need not dread,
'Tis scarcely deep enough a bee to drown,
As it sings harmless o'er its pebbly bed—
Aye, here it is! Stuck close beside the bank,
Beneath the bunch of grass that spindles rank
Its husk-seeds tall and high: 'tis rudely planned
Of bleached stubbles and the withered fare
That last year's harvest left upon the land,
Lined thinly with the horse's sable hair.
Five eggs, pen-scribbled o'er with ink their shells,
Resembling writing scrawls, which Fancy reads
As Nature's poesy and pastoral spells:
They are the Yellowhammer's; and she dwells
Most poet-like 'mid brooks and flowery weeds."

During spring and summer the Yellowhammer is associated with the *Rubus fruticosus* from the period of its flowering to that of its fully ripened fruit, the well-known Blackberry; but when the spring is past and "summer is over and gone," it betakes itself to the open fields and seeks its food on the ground, where it finds a plentiful supply of seeds, small-shelled mollusks, &c. As winter approaches, it assembles in flocks, and mingles with Finches and Sparrows around the outstanding ricks, and even ventures within the precincts of the farmyard in quest of grain or other kinds of food which such places afford. Soon, however, spring again appears, and with it comes a change of diet; for insects and their larvæ are then eagerly devoured—a kind of food with which the Buntings also feed their offspring.

No one of our native birds varies so much in colouring as the Yellowhammer; the differences in this respect are, however, too trivial to be regarded as specific, the variations being confined to the intensity or richness of its hues, and to the presence or absence of markings on the head. Some males have this part of a beautiful clear yellow, while others have a well-defined light-chestnut moustache bounding the lower part of the face; others, again, have the head and cheeks suffused with dark brown, without a trace of the

moustache. My collection contains many such varieties, of which those from chalk districts are the most beautiful. On the continent of Europe other varieties occur, and examples from Norway far exceed in size those from more southern countries, especially those of the British Islands. Having mentioned these facts, I must now state in what countries the Yellow Bunting is found besides our own. As regards Europe, it will not be necessary to say more than that it occurs in all parts of the continent, from the shores of the Mediterranean to Norway, Sweden, and Lapland; it also inhabits the whole of the northern regions of Africa, from Tunis to the Nile; further eastward it becomes scarce, but I possess the skin of a female from Erzeroom. I have never seen a specimen from India, neither does Schrenck include it in his List of the Birds of Amoorland.

M. Bailly, in his work on the ornithology of Savoy, states that "during the mornings of the months of October and November flocks of this Bunting pass through Savoy. They come from the north, traverse the valleys, and generally direct their course towards the southern countries of Europe. Some flocks, mingled with the Chaffinches and Cirl Buntings, emigrate from Savoy in order to pass the winter in more genial regions, and return again towards the end of February and beginning of March; but considerable numbers constantly remain there even during the most severe frosts. The Yellowhammer readily submits to confinement, where it will subsist on millet and hemp-seeds or buckwheat, but has no song to solace those who may keep it. When destined for the aviary, care must be taken that it be not frightened, lest it be destroyed by dashing itself violently against the bars of its prison. Its flesh is delicate and much sought after." Many modes of capturing it for the purposes of the table are resorted to. In the environs of Yenne, and particularly at St. Paul, it is hunted unmercifully. The villagers keep the entrances of their barns and granaries wide open during the day (in order that the birds may enter in quest of subsistence), conceal themselves behind the buildings, and, as soon as they have seen a number enter, rush with shouts to the doors and suddenly close them. Most of the frightened birds attempt to escape by the windows, and there become entangled in nets placed for the purpose, while those that continue to fly about are pitilessly killed with blows from rods or switches.

M. de Selys-Longchamps informs us, in his 'Faune Belge,' that "the Belgian villagers kill many of these birds when the ground is covered with snow, by a kind of sporting called *ramaille*. On a moonless night a man goes along the sides of old hedges, holding in his hand a torch of burning straw, towards which the Yellowhammers, quitting the hedges, direct themselves, and are immediately knocked down by two or three other men with large branches of thorn."

Mr. Hewitson states that the nest "may be found upon almost every briery hedge-dyke; it is most commonly placed upon the ground, but not unfrequently occupies the centre of a thick bush. It is formed outwardly of straws, bits of moss, sometimes sticks and coarse grass, finer towards the interior, which is finished with roots and a few hairs. The female sits closely, and is not easily driven from the nest. The singularly marked eggs, which are from three to five in number, always form a large portion of the bird-nesting spoils which fill the string of the schoolboy. Although subject to great variation, they are usually so characteristic of the species that it would not be easy to confound them with those of any other bird, except the rarer eggs of the Cirl Bunting. In form they are occasionally very long and oval, and sometimes as round as a marble. I have seen a single specimen of this egg so much suffused with colour that it might have been mistaken for that of the Cuckoo."

The adult male in summer has the head, cheeks, ear-coverts, and throat, chest, and centre of the abdomen lemon-yellow, the feathers surrounding the crown streaked with dusky; the lower edge of the ear-coverts is also similarly streaked; nape yellowish, passing into a greenish-olive tint on the lower part of the neck; feathers at the back and wing-coverts black in the centre and largely margined with sandy red; lower part of the back dark rust-red; primaries dusky black, narrowly edged with pale yellow; secondaries and tertiaries also dusky black, more broadly margined with rufous; tail dusky black, the two centre feathers margined with rufous, and the two outer ones on each side with a broad patch of white down the apical portion of the inner web; breast and flanks yellowish red, each feather darker in the centre; bill bluish horn-colour; irides, legs, toes, and claws light brown.

The Plate represents the two sexes, of the size of life, on a flowering branch of the Bramble, *Rubus fruticosus*; the beautiful little plant accompanying it is the *Pyrola minor*, obligingly sent to me for the embellishment of this work by the Rev. H. Harpur-Crewe.

Gould, John. 1873. "Yellowhammer or Yellow Bunting, *Emberiza citrinella* [Pl. 22]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323913>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323913>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323913>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.