GLYCOSPINA HORTULANA .

GLYCYSPINA HORTULANA.

Ortolan Bunting.

- Emberiza hortulana*, Linn. Faun. Suec., p. 84.
——— *Badensis*, Gmel. edit. Linn. Syst. Nat., tom. i. p. 873.
——— *chlorocephala*, Gmel. ib., p. 887.
——— *Tunstallii*, Lath. Ind. Orn., vol. i. p. 418.
——— *Malbeyensis*, Sparrm. Mus. Carls., tom. i. tab. 21.
——— *citrinella*, var., Flem. Hist. Brit. Anim., p. 77.
——— *pinguescens*, Brehm, Vög. Deutschl., tom. i. p. 295.
Citrinella hortulana, Kaup, Natürl. Syst. p. 142.
Hortulanus chlorocephalus, Bonap. in Parz. Cat. Ois. d'Eur., p. 4. sp. 131.
Glycyspina hortulana, Cab. Mus. Hein., Theil i. p. 128.
-

BIRDS not only contribute to the pleasure of man by their songs, their thousand actions, interesting ways, and varied habits and economies, but they also contribute their fair share towards his sustenance. Besides the amount afforded by the great group of Gallinaceæ, the marsh-, and water-birds, many of the smaller species form no inconsiderable item in supplying his wants and desires. In the acquirement of delicacies for the pampering of his appetite many devices are resorted to, some of which are accompanied by a degree of cruelty much to be deprecated: I allude to the means adopted to obtain the celebrated "pâté de foie gras," to the repeated takings of the nest of the esculent Swallow, and the continued robbery of the eggs of the Plover—acts which cause the Goose to die a miserable death, the Swallow to exhaust itself in the reconstruction of its glutinous cradle, and the Plover to weaken its system by a forced reproduction of its eggs. In Europe three little insessorial birds are especially regarded as delicacies for the table—the Wheatear, the Beccafico, and the Ortolan. The last of these (the bird represented on the accompanying Plate), which, although not indigenous to our island, is sometimes found here, is sent from the continent to London by thousands, but the great bulk of the community, the middle class and the poor, never see or partake of them; it is the noble and wealthy epicures alone who can afford to gratify their appetites with this costly "bonne bouche." The marchioness and the lady of the alderman consult their poulterers as to the more choice viands for the summer season, that nothing may be wanting for their next *recherché* dinner. A dish of Ortolans being the "chose," the fattening-cage is resorted to, and a dozen or more are taken out and killed if they have not already been trampled to death by their companions, or died during the night from apoplexy or sheer obesity, the little birds, which in a state of nature scarcely exceed the size of a Tit, having become a mass of fat the appearance of which is very unlike that of an ordinary plucked Bunting. This essence of millet and canary-seed is certainly not to be despised; and the clubman of our great metropolis frequently doubles the amount of his dinner-bill by ordering a couple of Ortolans. Alas for the fate of the little birds! which have been netted in Italy, Savoy, and France, and sent hither to be fattened and stifled in low cages, where they have not sufficient room to flutter their wings, for the special gratification of those whose gustatory enjoyments dominate over their other senses. The procuring of a string of Larks is, comparatively speaking, unattended with cruelty; a trapped Wheatear is killed in a moment, a Beccafico is destroyed with shot, or taken from the meshes of a net and immediately dispatched; but the Ortolan, captured during his migration from the southern to the northern parts of Europe, where he would otherwise have paired, loved, and bred, terminates the last summer of his existence in the cruel manner above described.

It is somewhat surprising that a bird so common on the continent, and which is a migrant withal, should not have been more frequently seen in this country; few, however, as are the instances of its occurrence, they are sufficient to entitle it to a place in our avifauna. The first specimen recorded as having been taken in England was captured by a London birdcatcher in Marylebone Fields, and was figured in Brown's 'Illustrations of Zoology,' a work published in 1776: this is the example described by Latham in the third volume of his 'Synopsis of Birds,' and moreover was the source, according to Yarrell, from which Gmelin, Lewin, Montagu, and others copied their descriptions. Bewick's figure was taken from a specimen caught at sea off the Yorkshire coast by the master of a trading vessel, and which subsequently passed into the possession of G. T. Fox, Esq. Another was killed near Manchester in 1827. A fourth, taken near London, in a birdcatcher's net, in 1837, along with Yellow Buntings, was deposited in the aviary of the Zoological Society of London in the Regent's Park. The bird is said to have been seen in Norfolk. Mr. Rodd has recorded that an example was killed at Trescoe, one of the Scilly Islands, in 1851; the next year one was

shot near Worthing, in Sussex, and in 1859 another at Lowestoft, in Suffolk. On the continent it is most widely distributed—from the hot countries of Spain and Italy to Holland, Sweden, Finland, and Norway. During my visit to Norway in 1856, few birds came more frequently under my notice; sometimes it was seen by the roadside, at others it was perched on stone walls and among small shrubby trees, but mostly on the fences of gardens and fields: my attention was attracted to it by the conspicuous colouring of its head and the tameness of its disposition. In a note sent to me by H. E. Dresser, Esq., that gentleman says, “The Ortolan breeds all over Finland; yet its nest is not so often found as might be expected. It arrives at Uleaborg in the latter end of May. I saw a good number on the 3rd of June, while walking along the banks of the Ija river; but it was too early to procure any eggs.” Speaking of the bird as seen in Savoy, M. Bailly states that it arrives in that country between the 15th and 20th of April, in small companies of four or six individuals, or in pairs, but that it is principally between the 25th and 30th of that month that it appears in great numbers. “As soon as they arrive they pair, and spread over the country, some resorting to the vineyards, oat-, and wheat-fields, others to places surrounded by hedges and copses, and others again to sandy or stony situations. These migratory flights are performed during serene and moonlight nights; and it is then that we frequently hear their cry of *thû, thê, thiû, thû, thê*, the last syllable of the three cries being ordinarily a little higher in tone than the others. They depart again at the end of August or September. The old birds are the first to leave, and are generally accompanied by the young ones that have already moulted: the others migrate as soon as they have finished changing their feathers; and these are often accompanied by one or two old birds, which usually fly at their head, as if to guide them. In the southern provinces, where they are very common, they are considered a great delicacy, and are fattened for the table in great numbers. For this purpose they are shut up in dark places, sometimes only lighted by the rays from a lantern, or in a cage covered over with silk, except the seed-drawer, to which light is admitted that they may see to feed. Millet seed, steeped for a moment in boiling water, is alone given to them. In these prisons the Ortolans, deprived of all exercise, and provided with an abundance of food, soon become covered with an amount of fat which would not fail to suffocate them, were they not killed in time to prevent it.”

When compared with the birds forming the genus *Emberiza* as now restricted, the bill of the Ortolan Bunting will be found to have a more prolonged and otherwise different form—a circumstance which has induced M. Cabanis to separate it into a distinct genus, *Glycyspina*, which, concurring with him in opinion, I have adopted.

The nest, which seems to be invariably placed in a slight hollow on the ground in fields of corn, is very similar to that of the Skylark, but is somewhat more compact; it is formed of dry grass and small roots, lined with finer grasses and hairs. The eggs are from four to six in number, of a bluish white, speckled and spotted with black.

The food consists of grain and various seeds when ripened, to which insects are added in the early part of the season.

As is the case with the other members of the family, much difference occurs in the colouring of the sexes, the male alone having the green head, which has obtained for it the trivial name of Greenheaded Bunting.

In summer the adult male has the head and cheeks of a greyish green; the back reddish brown, each feather with a darker centre; wings brownish black; the primaries narrowly edged, and the tertiaries and wing-coverts broadly margined, with reddish brown; tail brownish black, the two outer feathers on each side with a broad patch of white on their inner web, near the tip; chin, throat, and upper part of the breast yellowish green; under surface reddish buff, lighter on the abdomen and under tail-coverts; irides brown; bill reddish brown; legs, toes, and claws pale reddish-brown.

The female has the head of a greyer tint and streaked with dark brown, the breast streaked with dark brown, and her general colour not so rich as in the male.

The Plate represents the two sexes, of the size of life, on a branch of the Barberry, *Berberis vulgaris*, Linn.

Gould, John. 1873. "Ortolan Bunting, *Glycospina hortulana* [Pl. 27]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323918>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323918>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323918>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.