PLECTROPHANES NIVALIS.

PLECTROPHANES NIVALIS.

Snow-Bunting, or Snowflake.

Emberiza nivalis, Linn. Faun. Suec., p. 82.

——— *montana et mustelina*, Gmel. edit. Linn. Syst. Nat., tom. i. p. 867.

——— *glacialis*, Lath. Ind. Orn., vol. i. p. 398.

Hortulanus glacialis et montanus, Leach, Syst. Cat. of Indig. Mamm. and Birds in Brit. Mus., pp. 15, 16.

Plectrophanes nivalis, Meyer, 3. Theil des Taschenb. deutsch. Vög., p. 58.

——— *borealis, montanus et mustelinus*, Brehm, Vög. Deutschl., pp. 305, 306.

If we compare the avifaunas of the arctic and antarctic circles, we find that they are of very opposite characters—that probably no insessorial bird inhabits the southern land, and that Penguins, Gulls, and Petrels are almost the only birds that live in that inhospitable region. How different is it in the great polar basin of the north! how varied are the forms of birds which dwell therein! and what myriads of individuals there occur! If not congenial to the human race, it may be considered the elysium of a certain portion of the feathered creation: for where can they live more peacefully, or obtain their food with less trouble?—the great warmth of the brief summer and perpetual daylight being conducive to the life of so many of the lower animals upon which they live. If the shores of these ice-bound seas be not favourable to the growth of shrubs and trees, sufficient vegetation and insect life exist to nurture several little insessorial birds, of which, perhaps, the most conspicuous is the Snow-Bunting, a species which, from its abundance and extreme tameness, must be to the Esquimaux what the Sparrow and the Wagtail are to us. No northern country yet visited by Europeans, either in the Old or the New World, has been too near the pole for its existence; for, however high the latitude that has been attained, there the Snow-Bunting has been found breeding; and I strongly suspect that if the pole itself were reached, and islands or a mainland were to be found there, this interesting bird would be seen enlivening the waste; at least we may naturally conclude that such would be the case from the circumstance of its having been met with in every part of the arctic circle yet visited—Spitzbergen, Nova Zembla, Lapland, Russia, and Siberia in the Old World, while Greenland, from its position, unites it with the New. The remains of the departed and excellent Franklin and his heroic companions lie in the midst of the breeding-grounds of this species; and well does its mourning attire accord with the dreary resting-place of those brave men, many of whom were my personal friends. Heaven only knows whether the skeletons of our countrymen may not have afforded shelter for the nest of the beautiful Snowflake; for Captain Lyons found one placed in the bosom of a departed Esquimaux child—“a situation,” remarks the Rev. F. O. Morris, “suggestive of affecting thoughts, but the history connected with which must remain unknown until that day when both land and sea shall give up their dead.” One would imagine that birds are among the happiest of created beings; without effort, and with but little labour, they transport themselves from place to place, and from country to country, their wing-powers enabling them almost to annihilate time and space. They are certainly more independent than quadrupeds; for no impediments obstruct their passage from one region to another. They inhale freely the ambient air, through which they pass on their wonderfully constructed wings. To-day the Snowflake may be in the British Isles, to-morrow in Iceland, the day following in their breeding-grounds in Greenland, whence, the task of incubation accomplished, and the feathering of the young perfected, they return southward to avoid the rigours of winter, which would be fatal to so frail a bird. The individuals which leave us for the temperate regions of the northern hemisphere, singly or in pairs, in spring, return again in the autumn in flocks, and frequently hundreds together.

“This neat and elegant bird,” says Dr. Richardson, “breeds in the northernmost of the American islands, and on all the shores of the continent, from Chesterfield Inlet, to Behring’s Straits. The most southerly of its breeding-stations in the New World, that has been recorded, is Southampton Island, in the sixty-second parallel. Its nest is composed of dry grass, neatly lined with deer’s hair and a few feathers, and is generally fixed in the crevice of a rock, or in a loose pile of timber or stones. The eggs are greenish white, with a circle of irregular umber-brown spots round the thick end, and numerous blotches of subdued lavender-purple. On the 22nd July, 1826, in removing some drift timber lying on the beach of Cape Parry, we discovered a nest on the ground containing four young Snow-birds. Care was taken not to injure them; and while we were seated at breakfast, at the distance of only two or three feet, the parent birds made frequent visits to their offspring, at first timidly, but at length with the greatest confidence, and every time bringing grubs in their bills. The Snow-Bunting does not hasten to the south on the approach of winter with the same speed as other summer birds, but lingers about the forts and open places, picking up grass-seeds, until the snow becomes deep; and it is only during

the months of December and January that it retires to the southward of the Saskatchewan. It usually reaches that river again about the middle of February; two months afterwards it attains the sixty-fifth parallel of latitude; and in the beginning of May it is found on the coast of the Polar Sea. At this period it feeds upon the buds of the *Saxifraga oppositifolia*, one of the most early of the arctic plants; during the winter its crop is filled with grass-seeds. In the month of October, Wilson found a large flock running over a bed of water-plants, and feeding, not only on their seeds, but on the shelly mollusca which adhered to the leaves; and he observes that the long hind claws of these birds afford them much support when so engaged. The young are fed with insects."—*Faun. Bor.-Am.* vol. ii. p. 246.

In his 'Notes on the Ornithology of Lapland,' the late Mr. Wheelwright says:—"Although the Snow-Bunting did not appear to remain during the winter, we observed small flocks of it during our whole journey up north of Hernosand; and very soon after we arrived at Quickiock I shot specimens in nearly pure winter dress; they seem to leave the lowlands for the fells early in May. We never found a nest, although the bird breeds abundantly among the fells, and we shot old birds in their summer dress as well as young fliers in the end of July, and one of the latter as early as the 6th of that month. I am not surprised that we did not find the nest; for the wildest and most desolate spots on the fells appeared to be their summer home. On these fells there are thousands of acres, we might say many miles, covered with nothing but loose shingly slate and ironstone, and boulders of erratic rock, which are most difficult to traverse; and here we always saw the Snow-Bunting during the breeding-season; but when the young could fly they appeared to descend lower down on the fells."

During the months of September and October the British Islands are visited by numbers of these birds, the eastern coast particularly, the great promontory of Norfolk being a favourite place of resort; but it is less numerous in the southern and western parts of our islands. That it proceeds still further south is evident, Mr. Frederick Du Cane Godman having seen it in the Azores; it is said that it also visits the Canaries. I believe we have no direct evidence that the bird has ever bred with us; but we may reasonably assume that a few now and then remain for that purpose; for Macgillivray states, in his 'Natural History of the Deeside and Braemar,' that he "met with this species early in August in the corry and on the summit of Lochnagar, on the Glas-mheal, in the western corry of Cairn Toul, on the summit of Ben-na-muic-dhui, and in several other localities. Mr. Cumming and Mr. Brown inform me that it resides there all summer and breeds. In winter it frequents the valleys from Castletown to Ballater in small flocks. According to Mr. Stewart it breeds on Ben-Aun."

"Seen against a dark hill-side or lowering sky," says Mr. Saxby, in the 'Zoologist,' "a flock of these birds presents an exceedingly beautiful appearance, and it may then be seen how aptly the term 'Snowflake' has been applied to the species. I am acquainted with no more pleasing combination of sight and sound than that afforded when a number of these birds, backed by a dark grey sky, drop as it were in a shower to the ground, to the music of their own sweet tinkling voices."

So much diversity occurs in the colouring of the Snow-Bunting during its progress from youth to maturity that examples have been described as pertaining to different species, besides which a seasonal change takes place which transforms the bird amazingly; jet-black and snow-white is the characteristic of the breeding-dress of the male, while that of the female is browner, with streaks of a darker hue on the head and back, and she has much less white on the shoulder; the under surface, however, is white as in the male; the bills of both sexes at this season are jet-black.

The young, as may be seen in the accompanying Plate, are very unlike the adult, being olive-brown above, streaked with black, while the belly is tawny and the white shoulder-marks slightly tinged with the same hue.

The Plate represents a male, a female, and a nest of young, of the natural size, in summer plumage. The reduced tawny-coloured figure represents the dress in which the birds are frequently seen in autumn and winter. The red-flowering Lichen is the *Cladonia cornucopioides*.

Gould, John. 1873. "Snow-Bunting or Snowflake, *Plectrophanes nivalis* [Pl. 31]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323922>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323922>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323922>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.