AGIOTHUS LINARIA.

ÆGIOTHUS LINARIA.

Mealy Redpole.

- Fringilla linaria*, Linn. Faun. Suec., p. 87.
Linaria borealis, Vieill. Nouv. Dict. d'Hist. Nat., tom. xxxi. p. 341.
—— *Holbælli*, Brehm, Vög. Deutschl., p. 280?
Passer linaria, Pall. Zoog. Ross.-Asiat., tom. ii. p. 25?
Linaria minor, Leach, Syst. Cat. of Spec. of Mamm. and Birds in Brit. Mus., p. 75
Fringilla linaria borealis, Jenyns, Man. Brit. Vert. Anim., p. 139.
—— *borealis*, Temm. Man. d'Orn., 2nd edit. tom. iii. p. 264.
Linaria canescens, Gould, Birds of Eur., vol. iii. pl. 193.
Linota canescens, Yarr. Hist. of Brit. Birds, edit. of 1838, vol. i. p. 508.
Fringilla (Acanthis) borealis, Keys. & Blas. Wirb. Eur., p. xli.
—— *canescens* (partim), Schleg. Rev. Crit. des Ois. d'Eur., p. lxiii.
Acanthis linaria, Bonap. et Schleg. Monog. des Loxiens, pl. 52.
—— *holboelli*, Bonap. et Schleg. ib., pl. 53?
Ægiothus linarius, Cab. Mus. Hein., Theil i. p. 161.
Fringilla (Acanthis) linaria, Middend. Sib. Reise, tom. ii. pl. 11, p. 150.
Ægiothus linaria, Baird, Pacific Railr. Rep., Birds, p. 428.
-

For many years after ornithology had become a science, two British birds of this form were confounded under one name; and I regret to say that a mass of confusion exists with regard to their true synonymy, which I fear it will scarcely be possible to unravel; indeed I know not how it can be satisfactorily effected without an examination of the type specimens to which the various appellations were applied—a measure now hardly possible. We may, however, arrive at one certain conclusion—namely, that the species to which Linnæus applied the name *Fringilla linaria* was one commonly found in Sweden; and as the only Redpole which fulfils this condition is the present, we may safely infer that it is the bird he described by that appellation.

Although the Mealy Redpole is occasionally very numerous in our islands, it must be regarded as but an accidental winter visitor. The bird-catchers, who are generally excellent observers after their own way, will tell you that this season the Mealy or Stone Redpole is more numerous than the common or lesser species, and that their takes of this bird have consequently been great—a fact which may be confirmed or refuted by a visit to that great emporium of living birds, the Dials; or they may tell you that no Stone Redpoles have been seen either in this or the last year, or that for several seasons they have not seen a bird.

“The Mealy Redpole,” says Mr. Stevenson in his ‘Birds of Norfolk,’ “can scarcely be called an annual winter visitant, although flocks of more or less extent may be met with in several consecutive seasons; but now and then, from some cause not easily explainable, their total absence is remarked upon by our bird-catchers, and, as I have frequently experienced when most wanting a specimen to supply some loss in my aviary, not a bird has been netted the whole winter through. Their numbers and appearance also cannot always be accounted for by the severity of the weather, in this country at least, either at the time of, or subsequent to, their arrival on our coasts. In 1847 and 1855, the latter a very sharp winter, they were extremely plentiful; and in 1861, from the middle of October to the close of the year, probably the largest flocks ever noticed in this district, were distributed throughout the country. Hundreds of them were netted by the bird-catchers, being far more plentiful than the lesser species, and many still retained the rich flame-coloured tints of the breeding-season. Yet the weather throughout this period was not unusually severe; and in the previous winter of 1860–61, hardly a bird was taken, though remarkable for its intense frosts; and again in 1863 and 1864 they were equally scarce, with an almost equal degree of cold.

“I am not aware that the nest of this species has ever been found in Norfolk; but Mr. Alfred Newton has recorded the occurrence of a male specimen, in full breeding-plumage, at Riddlesworth, in July 1848, which he had ‘no doubt had bred there;’ I was also assured by one of our Norwich bird-catchers, that in the spring of 1862, after the large influx of the previous autumn, he observed a flock of twenty or thirty as late as the middle of April. Both the Mealy and Lesser Redpoles, from their tameness and engaging actions, are most desirable additions to the cage or aviary, but from their happy contented natures are liable to grow too fat, and, like Ortolans, when overfed, drop off the perch in a fit of apoplexy. Mr. Charles Barnard, of this city, had a brood of young Mealy Redpoles hatched off in his aviary at Stoke, in July 1860—a very uncommon circumstance with this species.”

The irregularity in the visits of this bird cannot, of course, be accounted for; all we know is that the bird

is ever plentiful in the north of Europe, and that it breeds over the greater part of Norway and Sweden. In the former country I observed it breeding on the Dovrefjeld, while in Lapland the late Mr. Wolley obtained numerous richly coloured specimens, nests, and eggs, and, in some notes communicated to Mr. Hewitson, says, "The Mealy Redpole is seen in most seasons throughout the winter in Lapland, though the greater number go southwards. Even in the breeding-time it seems to be a gregarious bird; for a considerable number of nests are to be found in a small space of the birch-forest; and the region of birch trees seems to be its proper habitat. In 1854, I principally met with it in a small district, at a great elevation, towards the Norwegian frontier. In 1855, when all soft-billed birds were so scarce after the preceding severe winter in the south, the Mealy Redpoles were abundant everywhere, from the very strands of the Arctic Ocean, over the mountains, where the nests were often close to the ground, as indeed they sometimes are in other situations, to the extensive forests of Munioniska."

"The nest of the Mealy Redpole," says Mr. Wheelwright in his 'Spring and Summer in Lapland,' "is one of the most beautiful I ever saw—perfectly cup-shaped, built of fine sticks, then a layer of fine grass, and next an interior lining of the white down of the willow and white feathers of the Willow-Grouse."

The eggs are said to be five or six in number, and, as figured by Mr. Hewitson, are of a very pale blue or bluish white, in some cases sparingly, and in others more profusely, speckled with pale rufous round the thicker end.

Every observer of nature must have noticed that a red colouring largely pervades the nuptial and summer dress of the *Loxiidae*. These bright colourings offer a striking contrast to the green foliage of the trees among which they respectively breed, give light as it were to the sprays of the pine, and life to the snow-covered branches of the spruce, their rosy breasts showing like living flowers among the birches and stunted willows during the inclement season of early spring.

Mr. Newton informs me that he considers only three species of this form to have been clearly defined. In this case the present bird must be subject to a greater variation in the size and form of its bill than any other small bird that has come under my notice. In some specimens the bill is as large, long, and pointed as that of a Goldfinch; in others it is short and triangular (the normal form of the genus); while I possess two examples, killed by myself on the Dovrefjeld, in which the bills differ from both slightly in form and altogether in colour, being nearly black. Mr. Stevenson and many other observers have noticed these differences; but no one, I believe, except Brehm, has been bold enough to characterize them as distinct species. They are indeed sadly puzzling to the ornithologist; but it is possible, if not probable, that they are dependent mainly on season. In the Brambling and Chaffinch the bill entirely changes its colour with the time of year; and it may well be, though from a different cause, that the same is the case with its form. Excluding from consideration birds that have been kept in cages, it will be found on examination that Redpoles having the longest bills are those which have been obtained towards the end of summer. This fact leads one to suspect that the peculiarity may be owing to the birds' having at that season lived almost exclusively on soft food (insects, buds, and the like), which would occasion no wearing away of the mandibles as is the case at other times of the year, when hard seeds form their principal if not their only diet. Again, in specimens obtained in winter, especially in high northern latitudes, the bill is so thickly clothed with feathers at its base that its apparent length is very greatly diminished, though if these be taken into account it will be found not so very much differing, either in size or shape, from what it is easily seen to be in summer. Future observations, no doubt, will set this point at rest.

The Mealy Redpole is subject to precisely the same changes of plumage as the Lesser Redpole: in the summer the rosy tints of its breast are most beautiful, and the further you proceed north the finer and brighter do they appear to be. In winter the male has the lores and throat brownish black; feathers of the head dusky, with a patch of deep red on the crown; back of the neck and upper part of the back pale yellowish brown, lower part of the back and rump greyish white; wings and tail dark brown, margined with brownish white; wing-coverts the same, the whitish tips forming two bands across the wing when closed; breast and fore part of the flanks suffused with rose-red; all the feathers of the upper surface and flanks with a small streak of brown down the centre; abdomen white; bill dull yellowish; irides, legs, and feet brown.

The female at this season is very like the male, but is paler in her general hue, and has no trace of the rose-red on the breast and flanks.

In summer the upper surface of the males becomes of a more uniform and darker brown, from the absence of much of the yellow bordering of the feathers; the head becomes of a deep blood-red; the breast, throat, and upper part of the flanks of a very rich rose-red, a trace of which colour appears on the rump and upper tail-coverts; and the flanks are conspicuously striated with blackish brown.

Specimens from North America agree exactly with our bird; but in that country there is certainly another species also, the true *Ægiothus canescens*, a much larger species.

The Plate represents two adult males and a female, life-size, and a branch of the alder (*Alnus glutinosa*).

Gould, John. 1873. "Mealy Redpole, *Ægiothus linaria* [Pl. 51]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323941>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323941>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323941>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.