DRYOCOPUS MARTIUS.

DRYOCOPUS MARTIUS.

Great Black Woodpecker.

- Picus martius*, Linn. Faun. Suec., p. 34.
— *niger*, Briss. Orn., tom. vi. p. 21.
Dryocopus martius, Boie, Isis, 1826, p. 977.
Dendrocopus pinetorum, Brehm, Vög. Deutschl., p. 185, tab. 13. fig. 3.
— *martius*, Brehm, ibid., p. 185.
Carbonarius martius, Kaup, Natürl. Syst., p. 131.
Dryotomus martius, Swains. Faun. Bor.-Amer., p. 301.
Picus (*Dryocopus*) *martius*, Keys. et Blas. Wirbelth. Eur., p. 34.
Dryopicos martius, Malh. Mém. Acad. Nation. Metz, 1849, p. 320.
Dryopicus martius, Malh. Mon. Pict., tom. i. p. 31, tab. 10. figs. 5, 6, 7.
-

THE Great Black Woodpecker was first described as a British bird by Latham, in 1785, on the authority of his friend Mr. Tunstall, who informed him that it had sometimes been seen in Devonshire. During the interval between that date and 1869, no less than thirty notices of its occurrence in various English counties have been recorded; and every ornithologist of repute, from Montagu to Yarrell, have included it as a conspicuous species in our avifauna; besides which many other persons—collectors, sportsmen, and country gentlemen—have stated to me, and, doubtless, to many others, that, in such and such a wood, on some named day, they had seen a Great Black Woodpecker, one or more affirming that they had followed the bird from tree to tree without being able, from its extreme shyness, to get within shooting distance. In the ‘Field,’ of January 29, 1870, Mr. J. E. Harting gave a detailed account of the first twenty-five recorded instances of the occurrence of the bird in our island; and in a note with which he has recently favoured me, he has enumerated a number of others, making a total of thirty reported instances of this bird in Great Britain. “It is not likely,” remarks Mr. Harting, “that they can *all* be mistakes.”

I regret, however, that I am unable to endorse this opinion. I had hoped that so fine a Woodpecker would have headed our indigenous *Picidæ*; but, of the many persons who have asserted the occurrence of the bird, not one, I believe, has been able to verify his statement by pointing out where an authenticated British-killed specimen is to be seen; nor, I think, can any public museum or private collection produce one. To omit, however, from the present work a species the appearance of which in our island has been asserted by so many respectable authorities, would scarcely be just. I therefore give a figure of it, for the purpose of showing those who are unacquainted with the bird what it is like, and how it differs from the Greater Spotted Woodpecker, the Jackdaw, and the Nutcracker, one or other of which has probably been mistaken for it.

Apart from England, and even just over the North Sea, the Great Black Woodpecker is tolerably common. In the pine-forests of Norway, Sweden, and Finmark it finds a congenial home, breeds, and rears its young. It also inhabits Switzerland, Sicily, Turkey, Greece, the Ionian Islands, Germany, Russia, Siberia, and Japan; and, if I mistake not, Mr. Swinhoe showed me specimens from Northern China. It is not found in the Atlas range, in North Africa; at least it is not included in Loche’s ‘List of the Birds of Algeria’; consequently the European shores of the Mediterranean is the limit of its range in a southern direction.

From one of the most interesting accounts of the bird as observed in Sweden I take the liberty of making a lengthy extract, nearly in the author’s words:—“Towards the latter end of May 1856,” says Mr. Simpson, “I happened to be staying with a Dane, the overlooker of a large forest belonging to Count L—. When he heard that I had come all the way from England to find the ‘Bo’ of the Spilkråka (*Picus martius*), he sent for his chief woodman to inquire what chance there was of getting one. The woodman said he had frequently seen birds throughout the spring, and had in former years noticed their ‘Bo,’ but that it was generally so high that nobody could get at it; that this year a pair of birds were known to frequent the edge of a clearing about four miles distant, and that if we would accompany him early the next morning we might possibly discover the object of our search. This was cheering intelligence, and caused us to make an early start. Our way lay chiefly through a monotonous wood of spruce-firs, very uninteresting in appearance and apparently destitute of any species of bird. But on crossing the clearing (a square of about 1000 yards), a Spilkråka was seen to slip quietly away from the upper part of a tall spruce and to fly towards the far corner of the square, where he uttered a single warning cry and disappeared. It took us a very short time to cross the remaining space in the direction he had gone, and it speedily became manifest that the object

of our journey was attained. Not far from the edge of a marsh stood the remains of an ancient birch some 30 feet in height; near the top was a fresh hole, out of which peered the unmistakable head and bill of a *P. martius*, gazing at the party without evincing much alarm. A tap from the woodman's axe caused the bird to retreat, when we commenced operations, pending which the old ones remained in the vicinity. Sometimes they came within gunshot and then swung round to the other side of the tree, poking so much of the head forward as to enable them to see what was going on. In this position they would beat a rapid and angry tattoo upon the trees, and occasionally utter, in addition to their ordinary cry, one of the most peculiar notes I ever heard from a European bird, and which more resembled the sharp and momentary ring of a shrill-toned bell than any other sound with which I am acquainted. We afterwards heard this note whilst operating upon another nest; but it appears not to be uttered except under great excitement, and then, perhaps, only by the female. The entrance-hole was within two feet of the top of the tree. On examining it I found the edges much rougher than those of the holes made by *P. viridis*, the sides sloping slightly upwards towards the interior. In shape it was elliptical, the horizontal being perhaps a fourth larger than the perpendicular diameter. The roof of the cavity was quite honeycombed by the strokes of the bird's bill; and this peculiarity was noticed in the two nests subsequently discovered. The depth of the cavity below the entrance-hole was rather less than two feet, quite sufficient to prevent my ascertaining what lay at the bottom. An alarm that the old stump was giving way with my weight brought me very speedily to the ground. The woodman meanwhile cut down a young spruce, and had it reared up against the ancient birch to prevent the anticipated catastrophe. He then went up himself and laid open the nest with his hands alone so far as to introduce his arm into the cavity. To my intense delight he proclaimed that there were eggs at the bottom. I immediately ascended, in order to see them *in situ*; but that was impossible without further enlarging the hole, for which my fingers were not strong enough. In the interval the party below had rigged up a long pole with a cap attached to the end of it, into which I deposited the eggs, and then slipped down in time to receive them once more into my hands. They were five in number, one much fresher than the others, which were partially incubated. In their smooth ivory texture these eggs very much resemble the well-known eggs of *P. viridis*. The fresher one was glossy white and sufficiently transparent to show the colour of the yolk; the others were slightly wood-stained, and presented a duller appearance, owing to their having been incubated. When blown this difference was removed, but the wood-stains of course remained and somewhat detracted from their beauty. In shape they resemble the eggs of *P. viridis*, and are not so much larger as I expected would be the case. The hole was of this year's making; the bird, however, is not in the habit of always making a fresh hole. Of the three inhabited nests discovered, two were new and one old."

The Great Black Woodpecker, like its congeners, is very shy and wary, and its power of flight appears to be very limited; its food consists of insects, which it searches for in the crevices of the bark of decaying trees. Latham states that "it is so very destructive to bees that the Baschirians in the vicinity of the River Ufa, as well as the inhabitants of other parts, who form holes in the trees 25 or 30 feet from the ground wherein the bees may deposit their store, take every precaution to hinder the access of the bird, and, in particular, to guard the hive with sharp thorns; notwithstanding which the Woodpecker finds means to prove a most formidable foe, and is most numerous where the bees are in the greatest numbers."

Specimens of the Great Black Woodpecker are often brought in the flesh from the Continent to the London markets, and I suspect that it is from this source that some of the reputed British examples have been derived; at least I know of one instance in which a specimen was purchased there, taken into the country, and palmed off as such.

The following notes of the colouring &c. of this bird were taken from an example in the flesh purchased in Leadenhall Market and kindly submitted to me by Mr. Gatcombe. The eyes were bright king's yellow; the naked orbits reddish black; basal portion of the sides of both mandibles wax-white, the remainder black; legs and feet olivaceous black; general plumage dull black, slightly tinged with green on the neck and chest; crown of the head scarlet.

The female has the back part of the head only scarlet.

The figures represent a male and female, of the size of life.

Gould, John. 1873. "Great Black Woodpecker, *Dryocopus martius* [Pl. 73]." *The birds of Great Britain* 3, -. <https://doi.org/10.5962/p.323959>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221726>

DOI: <https://doi.org/10.5962/p.323959>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323959>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.