PHASIANUS COLCHICUS Linn.

PHASIANUS COLCHICUS, *Linn.*

Common Pheasant.

Phasianus colchicus, Linn. Syst. Nat., tom. i. p. 271.

IN accordance with the expressed wishes of a great number of the subscribers to the present work, I here give a figure and description of the Common Pheasant. For my own part I consider it as much out of place in the 'Birds of Great Britain' as it would be to include therein similar delineations of the Peacock, the Turkey, and several other birds which have at various periods been introduced, and, favoured with fostering care, have lived and thriven among us; my opinion, however, has been met with the argument that, from the great number of years which have elapsed since its introduction, and the fact that it has become partially wild in our woods, it should form an exception. The true habitat of the Common Pheasant is doubtless Asia Minor and the western portion of Asia, whence in all probability the first living examples were brought to our islands; but even this is doubtful, since the bird has for many ages been known to be an inhabitant of Turkey, Bulgaria, and other parts of eastern Europe, and it is just as probable that our birds came thence.

Daniell, in his 'Rural Sports,' says Pheasants were brought into Europe by the Argonauts 1250 years before the Christian era, and are at present found in a state of nature in nearly the whole of the Old Continent.

"It may surprise the sportsman to read that this bird, which he finds wild in forests which can scarcely be said to have an owner, was brought from the banks of the Phasis, a river in Colchis, in Asia Minor, and artificially propagated with us and in other parts of the globe. History assigns to Jason the honour of having brought this bird, on his celebrated expedition, from the banks of the Phasis; and hence the modifications of the word, viz. Phasianus in Latin, Pheasant in our own language, Faisan in French, and Fagiano in Italian. The ancient Colchis, from which the specific name is derived, is the Mingrelia of the present day; and there, it is said, this splendid bird is still to be found wild and unequalled in beauty. The price Pheasants bore, according to Echard's 'History of England,' A. D. 1299 (being the 27th of the reign of Edward the First), was fourpence. At the same period the value of a Mallard was three halfpence, a Plover one penny, and a couple of Woodcocks three halfpence."—YARRELL'S *British Birds*, vol. ii. p. 278.

In those good old times the Pheasants which roamed about the woods and coppices of the British Islands were pure in blood, and adorned with all the pristine colours and markings of a true species; now, however, owing to the introduction of other kinds, and the crossings that have taken place, our country is tenanted by a set of mongrels, each individual, or at least by far the greatest portion of our stock, exhibiting an indefinite kind of coloration, so that scientifically there really is no interest in this bird as regards Britain; and in a utilitarian point of view, much harm has, in my opinion, accrued by the introduction of foreign blood into the veins of our own old stock,—not that I for a moment deprecate the infusion of new blood when it can be obtained from a distance and from individuals of the same species; for every breeder and physiologist is aware that the result would be a beneficial one. The introduction, however, of the Chinese *Phasianus torquatus* and the Japanese *P. versicolor* has plainly shown this in a certain way only. The first hybrids from either of these two birds with our own true *P. colchicus* are often wonderfully fine birds, generally, if not always, much larger in size, of far greater weight, and adorned with a plumage the colouring of which is often more beautiful than that of either of their parents. Nature, however, does not favour such liberties; for, interesting as they may seem in the eyes of an ordinary observer, the inutility of such unnatural proceedings is at once rendered manifest by the infertility, or partial infertility, of these larger and variously coloured individuals. No other result could in fact be expected, since every thinking person must at once perceive they are nothing more or less than true mules—or if not mules, that they would seldom or never breed *inter se*; and they are in fact shy of breeding with either of the parent species to which they are most nearly allied. Now these results having been certified by hundreds of experiments, it must be evident to our landed proprietors, sportsmen, and keepers that no beneficial effect has been brought about by mixing two or three species of Pheasant in the same covert, or even the crosses from any two of them. In making this somewhat sweeping statement I must, however, add that from what I have myself personally seen, and the information that has reached me on the subject, I do not positively affirm that hybrids are in all instances non-prolific, but that much uncertainty prevails on the matter. Some clutches of eggs may turn out pretty well, others be half addled; and the young that burst the shell often grow up a rickety and weakly stock.

In my work on the birds of Asia will be found a full account of the two birds mentioned as having been recently introduced into the British Islands, namely *P. versicolor* from Japan, and *P. torquatus* from Southern China, together with several other species of this beautiful group of birds, respecting which it is only necessary to mention here that they nearly all inhabit the northern part of that great land-section of the globe called Asia, and that none of them are found in India.

Of all the true Pheasants the *P. colchicus* in a pure state is the darkest in colour, and may be always recognized by the deep chestnut hue of its rump, a dark green stripe over each eye, and the uniform redness of its flank-feathers; the Chinese *P. torquatus* is conspicuous for the light silvery green mark over each eye, the glaucous green of its rump, and the light buff colouring of its flank-feathers; while the Japanese bird, with its splendid green breast and sides, differs materially from both. The birds usually shot in our woods exhibit an intermixture of all these tints and markings, no two being precisely alike.

As an evidence that the same colours in cross-bred birds cannot be perpetuated I append two notes which I find among my MSS. bearing upon this point.

“Burdett, the clever keeper of the Earl of Craven, informs me that a Pheasant which had a narrow ring round its neck the first year had a very broad one during the second, and that in the third it had totally disappeared. ‘I am positive of this,’ he says, ‘as it was never taken out of the pen in which it was kept.’”

“Mr. J. H. Gurney bred some extremely beautiful first-cross birds between a Green Pheasant, obtained at the Earl of Derby’s sale, and the species common in his woods at North Repps, in Norfolk; and although they appeared to be extremely healthy, and some of them exceeding four pounds in weight, the race could not be perpetuated, Mr. Gurney assuring me that after an interval of a few years there was no strain of the green bird left.”

On this head too, Mr. Stevenson, in his ‘Birds of Norfolk,’ has the following passage:—“From personal observation and inquiry, however, during the last two or three years, it appears that evidences of this cross, even in the coverts where these hybrids were most plentiful, are now scarcely perceptible, the strong characteristics of the Chinese bird apparently absorbing all the less-marked, though darker, tints of the Japanese. One of these birds, killed in 1853, weighed upwards of four and a half pounds; and many examples which were stuffed for the beauty of their plumage will be found in the collections of our county gentlemen.”

The accompanying illustration represents an old and true *Phasianus colchicus*, which has met with a fate to which hundreds of its brethren are annually subjected.

Gould, John. 1873. "Common Pheasant, *Phasianus colchicus* [Pl. 12]." *The birds of Great Britain* 4, -. <https://doi.org/10.5962/p.323971>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221609>

DOI: <https://doi.org/10.5962/p.323971>

Permalink: <https://www.biodiversitylibrary.org/partpdf/323971>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.