LIMOSA RUFA.

Bar-tailed Godwit.

- Scolopax Lapponica*, Linn. Faun. Suec., p. 62.
Limosa rufa, Temm. Man. d'Orn., 1815, p. 432.
Fedoa rufa, Steph. Cont. of Shaw's Gen. Zool., vol. xii. p. 77.
Scolopax leucophæa, Lath. Ind. Orn., vol. ii. p. 719.
Limicola Lapponica, Vieill. Nouv. Dict. d'Hist. Nat., tom. iii. p. 250.
Totanus leucophæus, Bechst. Naturg. Deutschl., tom. iv. p. 237.
—— *gregarius*, Bechst. ibid. p. 258.
Limosa ferruginea, Pall. Zoogr. Ross.-Asiat., tom. ii. p. 180.
—— *Meyeri*, Leisl. Nacht. zu Bechst. Naturg. Deutschl., tom. ii. p. 172.
—— *Noveboracensis*, Leach, Syst. Cat. of Indig. Mamm. and Birds. in Brit. Mus., p. 32.
Fedoa Meyeri, Steph. Cont. of Shaw's Gen. Zool., vol. xii. p. 75.
—— *pectoralis*, Steph. ibid. p. 79.
Actitis limosa, Ill. Prod. Mamm. et Av., p. 262.
Limosa lapponica, G. R. Gray, Gen. of Birds, vol. iii. p. 570, *Limosa*, sp. 3.
-

THE above long list of synonyms will show how much attention this well-known species has received from ornithologists; I trust, however, that it will not be encumbered with an additional name, and that the appropriate one of *rufa* will in future be retained as the specific appellation of the bird; *Lapponica* is certainly inappropriate, whatever may be its claims to priority; for the bird is not confined to Lapland, but is perhaps more generally dispersed than any other species of the genus.

If we take a glance at the northern regions of the Old World, as displayed on a map or an artificial globe, we see a few degrees to the eastward of the North Cape, between 70° and 75° N. lat., the island of Nova Zembla, standing out boldly in the Icy Sea; looking further east, we find the mainland of Siberia extending to a still higher latitude. These, and other parts of Siberia, with the interminable marshes and moorlands which skirt the rivers of that great country running into the Icy Ocean, from the sea of Kara, in the west, to Behring's Straits, in the east, northern Lapland and Finmark are, probably, the great nurseries of the Bar-tailed Godwit; and it is to those countries that the individuals which winter in Europe, China, and Japan most likely retire for the purpose of reproduction; I speak of Europe and China conjointly, because I think that the migratory movements of the birds may take place simultaneously in both those parts of the globe, and I may almost venture to assert that the individuals which return to Europe in the autumn have been breeding in Lapland, Finmark, and Nova Zembla. Some of these western Godwits winter with us in the British Islands, others on the estuaries of the rivers and sea-shores of the Continent; others, again, cross the Mediterranean, and distribute themselves in all favourable situations throughout Africa. The same influence, I have no doubt, affects the Chinese and Japanese examples which have bred in eastern Siberia; for they do not all stop in that country, but, like the western birds, impulsively seek a warmer climate, some individuals, I believe, extending their range, through the Philippine and all the intervening islands, to the continent of Australia, and even to Tasmania, on the shores of which island, in all suitable localities, I found numerous examples in their grey attire, but none in the red or breeding-dress. In asserting that it visits Australia I may be accused of discrepancy, since in my folio work on the birds of that country, and in my Handbook on the same subject, I treated the southern and northern Bar-tailed Godwits as distinct species; but I now believe them to be identical, and that the greater and less amount of spotting on the rump is a character not to be wholly depended upon. If this latter view be correct, how widely does this bird range! and how distant does it wander from its breeding-place! Young birds generally wander further from their birth-place than adults; and hence it probably is that we never see an Australian example in the red dress, which may not be acquired until the bird is on its return to the place in which it was bred.

Over Britain the dispersion of the Bar-tailed Godwit is very general; and wherever a tidal estuary or a low flat beach occurs, there it is sure to be found during the months of autumn. In the spring great accessions to those which have wintered with us take place, particularly on our eastern coasts, the additions mainly consisting of birds *en route* from more southern countries. They are now in their finest dress, having exchanged the grey plumage of winter for the rich chestnut livery of the approaching nuptial season; when thus attired in all the freshness of the spring moult, the bird, for a Sandpiper, is truly beautiful. May is the month in which they pass over Suffolk, Norfolk, and Lincolnshire, resting there for

a short time on the broads and inland waters. Unlike the Black-tailed Godwit, which makes, or rather did make, breeding-places of those localities, the present bird passes them by and proceeds to the higher latitudes of Lapland, Siberia, and Russia, in which a few solitary nests have been found; these localities, however, are, I am sure, merely the outskirts of its summer quarters. To justify this assertion, and to show that the outskirts only of their breeding-grounds have been entered on, I may state that I have at this moment before me the sale-list of the duplicate eggs from Mr. Wolley's collection, in which it is recorded that four pounds were given for one egg of this bird, and five pounds for another,—prices which must surely be considered enormous, and which clearly indicate that the most persevering oologist has not yet visited the bird's true breeding-quarters.

The Bar-tailed Godwit has two distinct lines of migration—a western one through Europe to Africa, and an eastern from China, through the Philippines, and across the line to Australia,—the great peninsula of India, and probably Burmah, forming a kind of neutral ground wherein it is seldom seen—so rarely, indeed, that Mr. Jerdon does not include it in his 'Birds of India,' although Mr. Blyth says that, "as a species occurring rarely in the Himalaya, it must be added to the Indian list." The immense range of the Himalaya Mountains, as we all know, are backed by almost interminable deserts of high land, forming a barrier over which either the Godwit or any other southern migrant would seldom attempt to pass.

"Eggs said to be those of the Bar-tailed Godwit," says Mr. Hewitson, "were brought from the North by the Rev. J. B. Tristram and also by Mr. Wolley, and with evidence sufficient to justify me in figuring them. The Bar-tailed Godwit was seen close to the eggs; the Black-tailed Godwit was never seen at all." Mr. Tristram, referring to the Bar-tailed Godwit, says:—"I found the bird in Finnmark, and shot several specimens in breeding-plumage, without any trace whatever of the black tail; I got the nest, and shot the bird, a female, the same morning, close to the spot, but I did not flush her off the nest. Mr. Wolley's eggs corresponded so exactly with mine that we could not distinguish them; but they were at once to be recognized among twenty eggs of the common Godwit. I have no moral doubt whatever on the subject, though my evidence does not amount to absolute demonstration." Mr. Wolley says this species breeds in marshes, chiefly in the neighbourhood of mountains, and that it gets up so warily from its nest that it is difficult to find it. The eggs figured are from Rowa, near Kittila, in Finland; I cannot, however, with Mr. Tristram, see anything about them to distinguish them from eggs of the other species; their close resemblance was to be expected.

"Dr. Middendorff, who met with this species breeding in numbers on the river Taimyr, in Lapland, says that it was not easy to find the eggs, as the birds would leave their nests with loud cries when he was at a great distance from them. The eggs are found at the end of June, two or three in a nest."—*Col. III. Eggs of Brit. Birds*, vol. ii.

"That the Bar-tailed Godwit breeds in the neighbourhood of Quickiock, in Lapland," remarks Mr. Wheelwright, "is pretty certain, although I never found the nest; but I fancy that although both the Godwits may breed here and there, in Lapland they principally do so more to the east. Both species are summer visitants around Vardol, in Finland. Sommerfelt however, has, never known the egg of the Bar-tailed to be taken there, although it breeds near Enara."

Speaking of the flight of the Bar-tailed Godwits, MacGillivray says it is "moderately rapid, light, somewhat buoyant, and greatly resembles that of the Curlew, and especially the Whimbrel. They emit a loud shrill whistle, and on rising utter a lower modulated sound somewhat like that of the bird just mentioned. Walking for the most part sedately, but sometimes running, they tap and probe the mud and sands in search of worms and other small animals, frequently mingling with Sandpipers and Ring-Plovers, along with which they often repose on the sands and neighbouring pastures at high water."

The number of Godwits that appear in different years varies greatly—but few in some, and many in others; generally speaking, too, the numbers that are seen in spring are much less than in autumn.

The sexes, like those of the allied species, differ very considerably in size, the female being by far the largest of the two.

The Plate represents the male in his spring dress, and the female in winter, of the size of life.

Gould, John. 1873. "Bar-tailed Godwit, *Limosa rufa* [Pl. 51]." *The birds of Great Britain* 4, -. <https://doi.org/10.5962/p.324008>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221609>

DOI: <https://doi.org/10.5962/p.324008>

Permalink: <https://www.biodiversitylibrary.org/partpdf/324008>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.