GALLINULA CHLOROPUS.

GALLINULA CHLOROPUS.

Moorhen.

Fulica chloropus, Linn. Syst. Nat., tom. i. p. 258.

Gallinula, Briss. Orn., tom. vi. p. 3. pl. 1.

Gallinula chloropus, Lath. Ind. Orn., vol. ii. p. 770.

——— *parvifrons*, Blyth.

——— *akool*, Jerdon.

THE information gained by the ornithologist who makes the Birds of the world his study should not, in my opinion, be withheld from those who take an interest in our native birds only. It will, I am sure, not be considered out of place, then, if, in some instances, I preface my remarks by briefly noticing a few of the other known species of the same immediate form, as I shall thereby be enabled to show that if similar physical conditions exist in other countries, however distant they may be, there will frequently be found birds of the same kind, and very often, but not always, of the same species. To render my remarks more clear, let me state that the distant country of Australia is tenanted, like England, by a Swift and a Swallow; it has also a Coot and a Moorhen. North America, too, has a Moorhen and a Coot. These representative species (for they are not the same) are most interesting to the ornithologist, and tend greatly to enhance the importance of his favourite science. The emigrant, who quits his native home, bidding adieu to all that has ever been dear to him, also hails with great pleasure these representative birds of the land of his adoption, and, of course, considers them the same; but were examples from the two countries placed side by side, he would soon perceive their differences, slight though they may be. Enough has been said to convey the meaning intended by these remarks, which are not made for the scientific ornithologist, but for the information of those readers of 'The Birds of Great Britain' whose occupations, and perhaps tastes, prevent them from entering deeply into this branch of natural science.

In Great Britain the Moorhen is a constant resident; and although it may be less numerous in some counties than in others, its distribution may be said to be general; for England, Ireland, Scotland, and even the Islands of Orkney and Shetland alike enumerate it as among the commonest of their birds. A much more extended range than that of the British Islands is, however, enjoyed by this bird, even more extensive than the entire continent of Europe; for I believe it is resident in all suitable situations throughout Africa, from the shores of the Mediterranean to the Cape of Good Hope; India, too, is inhabited by Moorhens, which offer no specific differences from our own bird; and I have a specimen now before me from Shanghai in China, presented to me by John R. Reeves, Esq. My readers will therefore readily perceive how extensively this bird is spread over the Old World. That the southern part of the peninsula of India is the extent of its range towards the equator in that country is, I think, very probable.

The situations affected by the Moorhen in the British Islands are the sedgy banks of lakes and rivers, reedy ponds, moats, beds of osiers, wet ditches, &c. It is naturally shy and recluse, avoiding the presence of man by immediately retreating from the open to the thick covert, or shrouding itself from observation by diving, and remaining half submerged until the apprehended danger has disappeared; this natural shyness, however, is frequently overcome, and the bird is rendered semi-domesticated, when its singular but graceful actions contribute to our pleasure and amusement. On the water the Moorhen is as buoyant as a cork, and its powers of swimming are as perfect as those of most water-birds; and it is by no means ungraceful on the land. Its neatly trimmed body, richly coloured bill, and red-gartered green legs contrast strongly with the sombre tint of its plumage. It frequently leaves the sedgy coverts, and quietly walks over the grassy meads, lawn, or pleasure-ground or garden, with all the ease of a land-bird, flirting its tail from side to side, and conspicuously displaying the white under-coverts, especially during the season of courtship. At this particular period the Moorhen becomes animated and noisy. The short, harsh cry of the males is now heard from the midst of the thick reeds; and if a close inspection were admissible, many battles and skirmishes between them would be witnessed for the choice of a mate. The Moorhen's energies are also aroused in an extraordinary manner, when a rat swims across the river, and perchance lands in the neighbourhood of its nest. I have seen a Moorhen furious at such an intrusion; and from the manner in which the rat has been assailed, I suspect he would avoid that locality for the future. Boldness and pugnacity appear to be part of the Moorhen's nature, and its quarrelsome disposition renders it an unpleasant neighbour to any peaceful bird that may live in close contiguity. This leads me to a trait in its character which will not redound to its credit; still it ought to be known. The Moorhen comes walking over the lawn, turning its head first to the right, then to the left, jerking its short, uplifted tail, apparently all peace and amiability;

but should the chick of a Fowl or Pheasant or a duckling cross his path, a single stroke of his pointed bill lays the little innocent dead at his feet, almost without a kick or struggle; and many losses to the keeper and the housewife have occurred which are not charged to the Moorhen. I was first made aware of this habit of the bird by one of the keepers of Sir S. Morton Peto, Bart., at Somerleyton; shortly afterwards I read a similar statement in the 'Zoologist;' and in order to determine the truth of these assertions, I inquired of James Hunt, one of the keepers of the Zoological Society, if such a habit in the bird had come under his notice; and I soon found all I had previously heard was true to the very letter. The following is his reply:—

"In answer to your inquiries respecting the Moorhens in the ponds in these Gardens, I beg to say they have always been very annoying and often destructive to young ducks. I have known them to kill several of a hatch, and partly to pull them to pieces. If the old duck attempts to assist them, the Moorhen immediately attacks, and sometimes mounts on her back, and continues tormenting her until she shakes it off by main strength, or dives under water, and so gets clear. The Moorhen often takes possession of the nest-boxes fixed in the ponds, and builds her nest therein, when no bird, not even a Goose, dare approach within some yards of it; in fact, the greater part of the time of the Moorhen during the breeding-season is taken up in annoying and tormenting the other birds in the pond."

The following interesting extract from the 'Zoologist' must close this part of the Moorhen's history. "At the beginning of July," says H. J. Partridge, Esq., of Hockham Hall, near Thetford, in Norfolk, "the keeper having lost several Pheasants about three weeks old, from a copse, and having set traps in vain for winged and four-footed vermin, determined to keep watch for the aggressor, when, after some time, a Moorhen was seen walking about near the copse; the keeper supposing that it only came to eat the young Pheasants' food, did not shoot it until he saw the Moorhen strike a Pheasant, which it killed immediately, and devoured all the young bird, except the leg- and wing-bones. The remains agreed exactly with those of eight found before. Perfect confidence may be placed in the correctness of this statement."

After this, let me say something more pleasing respecting the Moorhen, but before so doing direct the reader's attention to the accompanying Plate, whereon is depicted a newly hatched brood, one of three or four which are annually produced. That they are beautiful and interesting, no one, I think, will gainsay; yet how seldom are they seen, and how little is the colouring of the bird during the first few days of its existence known! This may be attributed to two causes—one the situations in which the nest is placed, the other the very short period (at most four or five days) during which this peculiar colouring is retained. Immediately on emerging from the shell, these infant birds take to the water, and follow their parents through labyrinths of thick and tangled herbage, at one part of the day sunning themselves on the prostrate rushes, at another threading the floating leaves of the water-lilies, both yellow and white; as night approaches, their sensitiveness to cold prompts them to seek shelter under the wings of the careful mother: the clucking male is now assiduously attentive, and protects both her and her progeny from danger, flirts his white tail, and exhibits evident signs of pleasure. These newly-hatched chicks, which a few hours previously were breathlessly imprisoned within the hard shell, have sprung into life with all the active energies of their parents, and, uttering a cheeping note, follow them about, swimming over the glassy pool, scrambling over the floating reeds in pursuit of insects, with the quickness of thought, and avoiding danger by diving beneath the surface with remarkable adroitness. From this period a great change takes place; for as the bird increases in size, feathers take the place of the downy covering, the characteristic colouring being olive-brown above and hoary white beneath, particularly on the throat and under surface; the bill now becomes of a uniform olive-green, a colour which it retains for the first year; after this, or in the second year, the adult livery is assumed, the feathers become more glossy, the bill assumes its brilliant hues of red and yellow, and the gaily coloured garter offers a conspicuous contrast to the yellowish green of the legs; the eye is now more in unison with the red of the bill and the frontal plate, and the Moorhens are in their finest attire—both sexes alike in their colouring, to a shade, even to the red and yellow bill. The average weight of several individuals was from 13 to 15 ounces, the lighter birds always being females. What more can I say respecting this well-known denizen of our marshes, the companion of the Will-o'-the-wisp? perhaps more would be tedious to the reader; with a description of its flight, its food, and its nest and eggs, I shall therefore close the Moorhen's history. Its wing-powers are not great, still they are sufficient to transport the bird from one part of the river to another, and for its nightly sallies, which are made known to all who live on the sides of rivers by the peculiar cry it emits while flying overhead, when darkness shrouds the object that gives utterance to it from our view: this is the voice of the Moorhen. His food consists of aquatic insects and their larvæ, mollusks of various kinds, every species of grain, and the shoots of young wheat or other cereals. The nest is sometimes placed on the flat branch of a tree, at others on stumps near the water's edge, among reeds, or on large floating masses of weeds. It is usually made of rushes, and is somewhat carelessly constructed. The eggs, which are from six to eight or nine in number, are of a reddish white, thinly spotted and speckled with dull orange-brown; they are one inch and eight lines long by one inch three lines broad.

The Plate represents male, female, and brood, about life-size. The plant is the *Nymphæa alba*.

Gould, John. 1873. "Moorhen, Gallinula chloropus [Pl. 85]." *The birds of Great Britain* 4, -. <https://doi.org/10.5962/p.324039>.

View This Item Online: <https://www.biodiversitylibrary.org/item/221609>

DOI: <https://doi.org/10.5962/p.324039>

Permalink: <https://www.biodiversitylibrary.org/partpdf/324039>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.