DAFILA ACUTA.

DAFILA ACUTA.

Pintailed Duck.

Anas acuta, Linn. Faun. Suec., p. 44.

— *caudacuta*, Leach, Syst. Cat. of Indig. Mamm. and Birds in Coll. Brit. Mus., p. 38.

Dafila caudacuta, Steph. Cont. of Shaw's Gen. Zool., vol. xii. p. 127, pl. 49.

— *acuta*, Eyton, Hist. of Rarer Brit. Birds, p. 60.

Anas longicauda (Briss.), Brehm, Vög. Deutschl., p. 868.

— *caudata*, Brehm, *ibid.*, p. 869.

Trachelonetta acuta, Kaup, Natürl. Syst., p. 115.

Querquedula acuta, Selb. Ill. Brit. Orn., vol. ii. p. 311.

— *caudacuta*, Macgill. Man. of Nat. Hist. Orn., vol. ii. p. 170.

Phasianurus acutus, Wagl. Isis, 1832, p. 1235.

AMONG the British members of the *Anatidæ*, or family of Ducks, there are three species which have especial claims to our notice: these are the Mallard, the Teal, and the Pintail. From the first have sprung all our domestic varieties, the excellence of whose flesh need not be dwelt upon; neither is it necessary to speak in praise of the delicate flavour of that of the little Teal; equal, if not superior, to both, is the flesh of the Pintail, a bird which is characterized by a greater elegance of form, and a more pleasing contour, than either of its congeners—one, moreover, which sits on the water with especial grace, and swims with unusual speed. All three species assimilate in the readiness with which they become partially domesticated, imparting life to our lakes and ornamental waters, in their wide and almost general distribution over the globe, and in the total change of costume which takes place at opposite seasons of the year. The plumage of the Pintail, though not so gay and contrasted as that of the Mallard or the Teal, is very pleasing, from its chaste and harmonious colouring; in form the bird is slender and elegant, its neck being considerably lengthened, and its two central tail-feathers prolonged to such an extent as to have obtained for it the name of Sea-Pheasant. How great is the contrast between the long central tail-feathers of this species, and the short curled ones of the Mallard!

In the British Islands the Pintail is very local; and although our eastern decoys supply the markets of London with a goodly number about a month before and after Christmas-time, few are either taken or shot at other seasons. Montagu states that it is most abundant in the north of England and Scotland, and especially in the Orkney Islands. "This assertion, however," says Mr. Selby, "I must in part contradict, as the result of long observation tells me it is of *rare occurrence* in the *northern counties* of England; and the same may be said of the southern districts of Scotland." Montagu's assertion, however, with regard to the Orkneys, is confirmed by the late Robert Dunn, who states that it is tolerably plentiful there, particularly in the island of Sanda, where it frequents the inland lakes more than the sea-shore, and leaves early in spring; he never met with it in the neighbouring Shetlands. In Cornwall, Mr. Rodd tells us that it is common at the Land's End in severe weather.

"A winter visitant to this country," says Yarrell, "it remains here till the spring, and is obtained by wild-fowl shooters on the coast as well as by fenmen on the rivers and lakes of the interior. It is one of the first species taken when the decoys begin to be worked in October." In Ireland it is a regular winter visitant in small numbers, and, as in England, is locally distributed. With regard to its distribution over other parts of the world, I may mention that it is tolerably common in North Africa, in the Grecian Islands, and thence throughout Europe, from the Mediterranean to the extreme north of Lapland, in Palestine, Asia Minor, throughout India, the Amoorland, China, Formosa, and Japan; in America it is found from the fur-countries to Honduras, and, doubtless, in all those parts of Mexico that are suitable to its habits. In confirmation of this vast extent of range, I may quote the following authorities.

Mr. E. C. Taylor, Dr. Leith Adams, and Mr. S. S. Allen, all enumerate it among the birds of Egypt; Captain Loche states that it inhabits the three provinces of Algeria; the Rev. Mr. Tristram mentions that specimens were shot near the brook Kedron, in Palestine; Mr. Jerdon says it is one of the most numerous winter visitants to India, frequenting large tanks and jheels, often in immense flocks; Mr. Swinhoe includes it in the lists of birds seen by him at Amoy, and between Takoo and Peking, in China, and in the island of Formosa; Captain Blakiston and Mr. H. Whitely in the birds of Japan; Von Schrenck in those of Amoorland; Dr. Walker obtained specimens, during the voyage of the 'Fox,' at Godhaven, in Greenland; Captain Blakiston in British North America; Mr. Brown includes it in his 'Synopsis of the Birds of Vancouver's Island;' Dr. Baird states that it is found over the whole of North America; and Mr. Salvin saw it at Belize, and observed it to be common on the Lake of Dueñas during winter.

That the Pintail does not habitually breed in this country is certain; but Mr. John Hancock informed Mr. More that he has known it to breed spontaneously in a swamp in Northumberland, which is now drained, and he believes the bird still breeds occasionally on the Northumbrian moors.

The evidence adduced by Mr. Henry Milner and Mr. Wolley in the following extract from the last edition of Mr. Hewitson's 'Illustrations of the Eggs of British Birds,' tends to prove that the eggs of the Pintail are unusually small for the size of the bird, and, moreover, are very like those of the Long-tailed Duck (*Harelda glacialis*); hence it will be read with interest.

"A few years ago," says Mr. Wolley, "I was very much surprised at the appearance of an egg given me by a gentleman (Mr. Henry Milner), who had brought it from Iceland in 1846, and who assured me it was out of a nest from which he himself had shot a female Pintail as it rose. It seemed so small for the bird, was so different from eggs previously supposed to be genuine, and looked like what I had been accustomed to consider Long-tailed Duck's. This single egg from Iceland, I accordingly valued very highly, and looked upon it as a veritable Pintail's, though this discovery of Mr. Milner's, like all others founded upon single nests, perhaps still wanted confirmation.

"In common with some other ornithologists, I had long been almost in a state of despair about most of those Ducks which do not occasionally, at least, breed in Great Britain. It was this which, more than anything else, determined me to take a journey to the far north; and, for many reasons, the fenny regions beyond the gulf of Bothnia seemed the most promising. On the 7th of June, 1853, I was some hundred miles up the river which forms the boundary between the territories of the King of Sweden and the Czar. Stopping at a house by the water-side, I could get nothing to eat but a few eggs, among which were nine of some kind of Duck, the appearance of which was exactly like the one I have mentioned as being brought from Iceland; but, having no means of identifying them, I dropped them into the kettle without the least remorse. On the 14th of June, some hundred miles further north, in fact within half an English mile of where I am now writing (Muonioniska), after a long and fruitless search for eggs, a Duck fluttered up a few yards off. There was a rush to the spot, greatly to the peril of the nest, sunk as it was in the moss. It was lined with down, and contained four eggs. The place was marshy, a few yards from the forest, on the rise of the hill. At midnight I went again to try and obtain the bird; it was just taking a circle over the nest, and it bent its long neck down to see that all was safe. I had a good look at it, as the sun was still shining. Twelve hours afterwards I had a shot at it as it rose rather wildly; but it did not seem to be hurt, and, as I had to continue my journey, I now reluctantly took the eggs; but I hoped that the down would serve to identify them, for amongst it were several breast-feathers. In the meantime, if I could trust my eyes, the bird was a hen Pintail; the eggs were, perhaps, a week sat upon, and just like some others I had attributed to the same bird. On the 18th of June, I and my line of beaters put up the old ones from three nests at different times in the course of twenty-four hours in a large marsh. I saw two very well, one of which I examined with my glass as it stood with its neck up in an open place some sixty or seventy yards off. It was a Pintail. All the eggs were nearly hatching, and the young, of which I preserved one or two, were all of the same species. I also kept the down and scattered feathers from each nest; and now I considered I had genuine Pintail's eggs of my own taking. But the most permanent proof was still wanting—the skin of a bird I myself should obtain from the nest. It was not till last season that I got this proof. On the 20th of May, 1854, I visited the same marsh; and in a little wooded island of a few yards in circuit, a Duck rose almost under my feet, and I shot it, feeling sure it was a Pintail, as it proved to be. There were six eggs, a day or two sat upon. The nest was made of a few twigs, mixed and lined with down from the mother's breast. It is usually made of long bleached grass, or anything that comes to hand. This bird breeds generally in marshes, and not very near large pieces of water. The eggs seem to be usually six or seven in number. The Pintail is one of the earliest breeders among the Ducks. They appear as soon as the water begins to open, and may be seen standing in pairs at the edge of the ice. As soon as the Ducks are hard sitting, the Drakes go about in flocks, having apparently deserted their mates." The eggs are of a clay-colour, slightly tinged with olive, and measure about two inches in length by one inch and a half in breadth.

In Lapland Mr. Wheelright always found the nest of the Pintail in the small willow plantations that skirt the foot of the fells. He obtained his first nest on the 4th of June.

Different as is the garb of the two sexes in winter, as shown on the accompanying Plate, soon after the female has incubated her eggs, the male throws off his finery, and assumes a dress so like that of his mate, that, except in size, the two sexes are very similar in appearance; the summer dress, however, is carried but a short time; for early in autumn the fine winter costume of the male is again assumed. Selby believed that these changes were produced by a change in the colour of the feathers, rather than by a renewal of them; but I think that this is not the case, and that the feathers are shed upon each occasion.

The Plate represents an old male, of the size of life, with a reduced figure of a female in the distance.

Gould, John. 1873. "Pin-tailed Duck, *Dafila acuta* [Pl. 18]." *The birds of Great Britain* 5, -. <https://doi.org/10.5962/p.324062>.

View This Item Online: <https://www.biodiversitylibrary.org/item/222497>

DOI: <https://doi.org/10.5962/p.324062>

Permalink: <https://www.biodiversitylibrary.org/partpdf/324062>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.