PODICEPS RUBRICOLLIS.

PODICEPS RUBRICOLLIS.

Red-necked Grebe.

Podiceps rubricollis, Lath. Ind. Orn., tom. ii. p. 783.

Colymbus rubricollis, Gmel. edit. Linn. Syst. Nat., tom. i. p. 392.

——— *subcristatus*, Gmel. ib., p. 390.

——— *Parotis*, Gmel. ib.

——— *griseigena*, Bodd.

Pedeithya subcristata, Kaup.

Colymbus cucullatus, Pall. (Bonap.).

——— *nævius*, Pall. (Bonap.).

To speak of this bird as a rarity in our country would be incorrect; still it is so in a certain sense. Britain is not its true home; and the individuals that have been seen are migrants from the Continent, and particularly from the north.

It is only during the spring and breeding-season that the curious tippets and high-coloured necks and egrets adorn the varied members of the genus *Podiceps*, and it is seldom in this state of finery that the Red-necked Grebe is killed in our island; still it has occurred in this dress, and thus I have represented it. My Plate will have the greater interest when I state that the feathers of the crown and neck are drawn in their natural position, as seen in an individual which lived for some time with Mr. Bartlett, now Superintendent of the Zoological Gardens in the Regent's Park. Mr. Bartlett informed me that the bird became quite tame and familiar, and in a domesticated state changed from its winter to its full summer dress, in which state it died. It is in autumn and winter, then, when the head and neck are without ornament, or like the rest of the body, that the Red-necked Grebe visits the Norfolk and Suffolk broads and other extensive meres and inland waters, and the larger rivers of our island: instances of these occurrences are on record too numerous to mention; for in every county, from Cornwall to Orkney, it occasionally appears. In his 'List of Cornish Birds,' Mr. Rodd speaks of it as "quite as often occurring as the last species (*P. cristatus*), frequenting the same localities. Sometimes killed towards the spring, when some of the red feathers appear, characteristic of its nuptial livery." Macgillivray says, "I have procured this species, with all the other Grebes, in the Frith of Forth." Thompson, in his work on the 'Birds of Ireland,' states that he had opportunities of examining five specimens; in the stomach of one of these were found the remains of several shrimps (*Crangon vulgaris*) and fishes, with ear-bones of small Gadidae, a pipe-fish (*Syngnathus acus*) ten inches in length, and a number of feathers of the bird's own body: none of these five individuals were adult.

The following note on the occurrence of this bird in Norfolk has been transmitted to me by Mr. Stevenson of Norwich:—

"A regular though not very numerous visitant late in autumn and early spring, appearing on our broads and inland waters between the beginning of November and the middle of March. Most of the specimens obtained are in immature plumage; but I have seen adult birds in their winter dress, and some also with traces of the red throat. According to Messrs. Gurney and Fisher (writing in 1846), a pair of this species occasionally remain to breed in this country; but these instances are, I imagine, extremely rare. A very beautiful specimen in full summer plumage, in Mr. Gurney's collection, was shot at Yarmouth about the 2nd of April, 1848, and another at Scotland, on the 22nd; but since that date their latest appearance here in spring to my knowledge, has been the 18th of March. The late Mr. Hunt, of Norwich, in his 'List of Norfolk Birds,' states that a pair of these birds were once killed near the Foundry Bridge in this city."

Temminck states that the Red-necked Grebe is nowhere more plentiful than in Holstein; and Mr. Dann informed Mr. Yarrell that it "is common, during the breeding-season, on many of the shallow reedy lakes at the head of the Bothnian Gulf, particularly between Pitea and Lulea. They seem to be confined to the vicinity of the coast of the Baltic. I have never met with them anywhere in the interior of the country, except in Scona and in the southern provinces of Sweden, although the whole of Northern Scandinavia abounds with lakes. The character of these lakes, where alone I have seen and procured specimens of the Red-necked Grebe so far north as latitude 66°, is precisely similar to that of the broads in Norfolk and the meres of Holland, where some of the Grebes are so numerous. Swedish ornithologists have confined the locality of this Grebe to the southern parts of Sweden; but having procured the old and young birds in August, and seen them in considerable numbers, two years in succession, in the same localities, no doubt can exist that they are regular visitants. The eggs I did not see; but the peasants, on finding a nest, are in the habit of leaving one egg, and the female will continue to lay, as long as one is left, until nature is

exhausted. These Grebes are by no means shy, and when undisturbed, amongst the reeds and grass, keep up an incessant croaking. They swiftly glide through the water, and dart through thick, entangled masses of weeds and grass with the ease and rapidity of a fish." The late Mr. John Wolley, acting on information supplied him by Mr. Dann, found that this bird was well known on the Kalix Noir, in the north of Sweden, and subsequently succeeded in getting specimens of its eggs from the locality where that gentleman met with it many years before.

Independently of the countries mentioned above, I may state that it inhabits every other part of Europe; it is included in the 'List of North African Birds' by Captain Loche; and specimens have been transmitted from Trebizond in Persia. It does not extend its range to the peninsula of India; but I find it included by Schrenck in his account of the Birds of the Amur-land, and Temminck says it is found in Japan. In Greenland there is a bird of this form, which is so similar to the *Podiceps rubricollis* that they have been considered identical by some, while others have regarded it as distinct, and have assigned to it the specific designation of *P. Holbælli*. Among my MSS. I find a note to the following effect:—"American specimens agree with European, except in being somewhat larger." Dr. Baird, who calls it *Podiceps griseigena*, evidently considers the bird identical with ours. It is likely, however, the American and Greenland birds may be the same, and distinct from the true *P. rubricollis*, in which case the name of *P. Holbælli*, assigned to it by Reichenbach, should be retained.

In no respect do the sexes differ in colour: the same law which affects the male is also carried out in the female; both assume the ornamental head-dress in summer, which gives place to a more sombre hue in winter.

No difference occurs in the nidification of this Grebe from that of the other members of the genus; the nest is placed on the surface of the water, among aquatic herbage and reeds, of which materials it is also built. The eggs are four or five in number, of a pale greenish white, and are somewhat smaller in size than those of *P. cristatus*. They are often stained by the materials with which the nest is built, till they acquire a rich orange-red hue; and it seems commonly the case for eggs of the Grebe to be more brilliantly dyed than those of any other species.

Crown of the head and back of the neck dark olive-brown; upper surface of the body brownish black; cheeks and throat brownish grey, bordered with greyish white; primaries brownish black; secondaries white; front of the neck, chest, and upper portion of the flanks rich rusty red; breast and abdomen silvery white; bill brownish horn-colour, except at the base, which, with the gape, is orange-yellow; irides red; tarsi clouded with pea-green; upper side of the toe bluish white, particularly in the centre of the lobes.

The young bird of the year has neither the red neck nor the elongated head-feathers; the throat is brownish, and the abdomen less silvery; the part of the neck which is red in summer is brown in the youthful state; irides brown; base of the bill paler orange.

The Plate represents the two sexes of the size of life, and a reduced bird in the distance. The plant is the Buckbean (*Menyanthes trifoliata*).

Gould, John. 1873. "Red-necked Grebe, *Podiceps rubricollis* [Pl. 39]." *The birds of Great Britain* 5, -. <https://doi.org/10.5962/p.324083>.

View This Item Online: <https://www.biodiversitylibrary.org/item/222497>

DOI: <https://doi.org/10.5962/p.324083>

Permalink: <https://www.biodiversitylibrary.org/partpdf/324083>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.