HYDROCHELIDON NIGRA.

HYDROCHELIDON NIGRA.

Black Tern.

- Sterna fessipes*, et *nævia*, Linn. Syst. Nat. (1766), tom. i. p. 228.
—— *nigra*, Briss. Orn. (1760), tom. vi. p. 211.
—— *plumbea*, Wils. Am. Orn., vol. vii. 1831, pl. lx., young.
Hydrochelidon nigra, Boie, Isis, 1822, p. 563.
—— *nigricans*, Brehm, Vög. Deutschl., p. 794.
—— *obscura*, Brehm, ibid., p. 795.
—— *fessipes*, G. R. Gray, Gen. of Birds, vol. iii. p. 660, *Hydrochelidon*, sp. 5.
—— *plumbea*, Lawr. Gen. Rep. 1858, p. 864.
Virelva nigra, Steph. Cont. of Shaw's Gen. Zool., vol. xiii. p. 167.
-

THE generic term *Hydrochelidon* was instituted by Boie for a small section of the *Sternidæ* possessing certain peculiarities of structure accompanied by an equally peculiar style of colouring, and whose habits and economy are unlike those of the other members of their family, from which they also differ in the kind of situations they frequent. The ordinary Terns or Sea-Swallows, genus *Hirundo*, are true sea-birds, which either retire to shingly beaches or ascend the larger rivers with banks of a similar character for the purpose of breeding, incubate their two eggs on the bare ground, fly with a laboured flapping motion of their wings, and descend upon their piscine food with a perpendicular stoop like that of the Kingfisher. The members of the genus *Hydrochelidon*, on the other hand, frequent the inland fresh waters (lakes and rivers) rather than the open sea, feed principally upon large winged insects, which they take in the air, and deposit three or four eggs in a nest of weeds placed in a tuft of grass in the midst of reed-beds. The true sea-Terns have short webbed feet, while the feet of the marsh-Terns, as the birds of the present form are called, are more lengthened and have the interdigital membrane but little developed. The Black Tern, which is a migrant, comes to us in the spring from a warmer climate, and, if any suitable locality be left in which it may remain unmolested, will reproduce its kind during its stay. On this point Mr. Stevenson, of Norwich, writes to me :—"Although this species formerly nested in considerable numbers, both on our eastern broads and in the fens of the western part of Norfolk, it has almost ceased to breed in the county. I know of but one or two solitary instances of a pair remaining to breed with us during the last few years ; Mr. Newton informs me that in the spring of 1852, owing to the extent of land then under water from the immense floods of the previous winter, two or three pairs bred in Feltwell fen, where they had not been known to remain for some years. Draining, and the abominable system of indiscriminate egging, are the principal causes why this bird, the Black-tailed Godwit, and other species now proceed further north. The old birds regularly appear on the coast every spring during the months of April and May, and again with their young in autumn (August, September, and October). The Black Tern is said to have formerly bred at Winterton, near Yarmouth ; and Lubbock, in a communication to Yarrell, says 'the great breeding-place, in a wet alder-carr at Upton, where, twenty years back, hundreds upon hundreds of nests might be found at the end of May, has been broken up some years.'"

During the numerous visits I have made to the middle portion of our beautiful Thames in the month of May, for the last forty years, I have seldom missed seeing the Black Tern hawking over the reaches in the neighbourhood of Maidenhead, Cookham, and Marlow. At that season they are apparently passing over our island from the Bristol Channel to some eastward localities, and they merely stay for a few hours, in their course down the river, one day at Henley, the next at Maidenhead or Windsor, thence proceeding to the Nore and other parts of the eastern coast. From the 10th to the 19th of May, 1866, several solitary individuals passed my boat ; and a similar occurrence took place in the same locality the succeeding year.

Occasionally I have seen the Common Tern, the Arctic Tern, and the present species in the same reach at one time ; an example of each of the three species, all of which fell to my own gun in 1866, may be seen at Taplow Court, Mr. C. Pascoe Grenfell having kindly accepted them as a memento of these birds occasionally occurring in his neighbourhood. In autumn the young of the Black Tern sometimes occurs ; and the flying figure in my Plate was taken from an individual killed in the same locality.

Nothing can be more easy than to distinguish the Black Tern, on the wing, from the ordinary sea-Swallows ; so different are they in appearance that, if seen flying together at the end of a reach half a mile distant, the fisherman will remark, "There is one of the black Terns."

As before stated, the flight of the typical or sea-Terns appears to be of a laboured character ; the marsh-

Terns, on the other hand, pass through the air with a more bounding motion, turning to the right or to the left and dipping perpendicularly towards a Dragonfly or any other insect that may have attracted their attention. Their aerial actions have been compared to the hawking of the Nightjar; and I may add that they are not unlike those of the Great Bat (*Vespertilio noctula*).

"The evolutions of the Black Tern are so rapid, and its turns so short," says Montagu, "that by this means it sometimes escapes the talons of predacious birds, as we had once an opportunity of witnessing. In a very hard gale of wind many Terns were sporting over the water, when a Peregrine Falcon passed like a shot, singled out his bird, and presently coming up with the chace, made a pounce; but the great dexterity of the Tern avoided the deadly stroke, and took a new direction. The Falcon by his superior velocity, soon regained sufficient elevation to successively repeat his pounces, but at last relinquished the pursuit."

Scotch writers state that the Black Tern is not common in that northern country; and those who have written on the Birds of Ireland, that it is an occasional visitor only, chiefly in autumn, and in an immature state.

Mr. Rodd when speaking of its occurrence in the extreme west of England, informs us that it is "generally observed in the autumnal months, and nearly every year, in more or less numbers, both on the sea-side and inland. In its full black plumage it is rarely met with in Cornwall."

Temminck, in his 'Manuel d'Ornithologie,' says:—

"*Habite*: les rivières et les bords des lacs d'eaux douces, mais particulièrement des marais; très-accidentellement sur les côtes maritimes; assez abondant dans le nord, jusques au cercle arctique; très-nombreux en Hollande et dans les grands marais de la Hongrie. Vit en grandes troupes dans les marais de Tombole et d'Ostia, mais n'y vient pas avant les premiers jours d'avril.

"*Nourriture*: insectes ailés et vers aquatiques.

"*Propagation*: niche en grandes bandes dans les marais, parmi les roseaux clair-semés et sur les grandes feuilles de nénuphar qui flottent sur les eaux; pond depuis deux jusqu'à quatre œufs, d'un olivâtre clair marqué de nombreuses taches brunes et noires, dont la réunion forme un large cercle sur le milieu de l'œuf."

The late Captain Loche states that it frequents the three provinces of Algeria.

Besides inhabiting all the temperate parts of Europe, North Africa, Madeira, Asia Minor, and Persia (according to Mr. Jerdon), it must be added to the Indian list, Mr. Blyth having seen Indian examples collected by Dr. L. Stewart. I believe it is also found throughout the whole of North America. Dr. Elliott Coues, in his valuable review of the Terns of that country, says:—"I have critically compared quite a series of European and American specimens in all stages of plumage, but have been entirely unable to detect the slightest discrepancies between the birds of the two continents. The specimens before me are all absolutely identical in size and relative proportions of different parts; and the colours of those of the same age correspond minutely. There do not appear to exist the slightest characters upon which to base specific distinction."

A nest of four recent eggs, that came into my possession before being blown, were of a pure olive, profusely marked with large blotches of dark rich umber, most numerous and forming a distinct though somewhat irregular zone towards the larger end; they varied a little in size, three being an eighth of an inch longer than the fourth, and two being a sixteenth wider than the others, the largest being one inch and three-eighths in length, by one inch and a sixteenth in breadth.

There is little or no difference in the size and colouring of the sexes. The adults in summer have the bill black, the inside of the mouth flesh-colour; the irides dark brown; the tarsi, toes, and webs dull red; nails black; head, neck, breast, and belly dark or blackish lead-colour; back, wings, and tail slate-grey; vent and under tail-coverts white.

The Plate represents an adult male and female, of the natural size, and a young bird in the dress of its first autumn, about a third less.

Gould, John. 1873. "Black Tern, *Hydrochelidon nigra* [Pl. 75]." *The birds of Great Britain* 5, -. <https://doi.org/10.5962/p.324119>.

View This Item Online: <https://www.biodiversitylibrary.org/item/222497>

DOI: <https://doi.org/10.5962/p.324119>

Permalink: <https://www.biodiversitylibrary.org/partpdf/324119>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.