



ALASKA REFRESHMENTS

Butterflies of the arctic regions join in obtaining moisture from wet ground on a warm sunny day just as do tropical or temperate species of related kinds. This scene shows a roadside cut in the Tanana River Valley of Alaska, showing the primeval soil condition at the top, and the disturbed soil below. Water seepage from the peat soil above has attracted *Colias philodice vitabunda* and *Colias gigantea* males as well as an *Erebia sp.* male at the upper left center. Some blues are visible in the vicinity of the *Erebia*. Early July, 1948.

W. Hovanitz



Hovanitz, William. 1965. "Alaska refreshments." *The Journal of Research on the Lepidoptera* 4(2), 113–113. <u>https://doi.org/10.5962/p.333506</u>.

View This Item Online: https://doi.org/10.5962/p.333506 Permalink: https://www.biodiversitylibrary.org/partpdf/333506

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder Rights Holder: The Lepidoptera Research Foundation, Inc. License: <u>https://creativecommons.org/licenses/by-nc-sa/4.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.