



#### HABITAT: *ARGYNNIS CALLIPPE LAURINA*

The unsilvered form of this complex which includes *callippe*, *comstocki*, *macaria*, *laurina*, *coronis*, *rupestris* and others is found along the lower elevations of the western side of the Sierra Nevada mountain range in California. Occasional specimens of wholly or partially silvered are to be found throughout the range of the race but they are most common toward the south where populations gradually increase in the percentage of silvering until the populations may be called *macaria* in the Kern basin. Silvering picks up again at the north end of the range.

At this location, the primary local vegetation is the Digger Pine and oak woods. Adults fly after the grass is dry, late June and early July. The photograph was taken a few miles southeast of Mariposa, Mariposa Co., California, early July, 1969.

William Hovanitz





Hovanitz, William. 1968. "Habitat: *Argynnis callippe laurina*." *The Journal of Research on the Lepidoptera* 7(1), 50–50. <https://doi.org/10.5962/p.333529>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/225389>

**DOI:** <https://doi.org/10.5962/p.333529>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/333529>

#### **Holding Institution**

Smithsonian Libraries and Archives

#### **Sponsored by**

Biodiversity Heritage Library

#### **Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: The Lepidoptera Research Foundation, Inc.

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.