

I had a good view of it for 20 or 30 seconds. Points recognized were (1) size, about that of a Ring-billed Gull, (2) white and pearl-gray coloration, (3) angular outline of wings, quite different from the curved front wing-edges of a flying Gull. A moment after I saw it, it uttered its characteristic cry a second time. It uttered the same cry at least twice more before it passed out of sight over the house-tops. I had no opportunity to observe the bird through binoculars.

In connection with this identification it is pertinent to state that I had visited a breeding colony of American Caspian Terns, at Fog Island, Saguenay County, Quebec, every summer for the twelve years immediately prior to 1935, and that I had thus become very familiar with the appearance and the voice of the species. I am wholly certain that the bird recorded above was a Caspian Tern.

The day of this observation was warm, with a strong southwest wind. It was partly cloudy, but the light was good when the observation was made.

There appears to be no previous record of a Caspian Tern in the Ottawa district.—HARRISON F. LEWIS.

THE ROSS GOOSE (*Chen rossi*) IN ALBERTA.—When driving through the country east of Bittern Lake on the afternoon of October 12th, 1935, I noted two white geese feeding in a grain field, in company with a large flock of Canada Geese. At the time I believed them to be Lesser Snow Geese, small numbers of which occasional-

ly mix with the "honkers". On October 17th, Alex Burgess, a local hunter brought me a pair of very small white geese he had killed that morning as they rested on the sandy shore of Bittern Lake, about four miles south of where I had, a few days before, seen the white geese. These proved to be Ross Geese and were probably the ones I had seen on the 12th of the month. The absence of warty growths at the base of the bills would indicate they were birds of the year, probably less than four months old. Measurements of the male were: length, 22½ inches; extent, 45 inches; wing, 14½ inches; and bill, 2 inches. The female was slightly smaller. Legs and feet bluish and bill reddish. Weight of male 3¼ pounds, female 3 pounds. Mallards killed the same day weighed from three to three and a quarter pounds. The plumage of both birds had a decidedly rusty appearance, much like that commonly found on Lesser Snow Geese, a result of their frequenting iron-impregnated waters.

Wintering in California, the Ross Goose is believed to migrate, chiefly through Alberta, to and from its nesting grounds in the Arctic regions. There are, however, few records of its occurrence in this province, probably due to the fact that "wavies" are not popular, and are seldom killed by hunters. At rare intervals during the past 40 years I have heard of small white geese, no larger than a mallard, having been killed in Central Alberta, but those referred to above are the only ones I have ever seen. A more appropriate name for this diminutive "wavey" would seem to be the "Least Snow Goose."—FRANK L. FARLEY.

BOOK REVIEWS

Insect Enemies of Shade Trees, by Glenn W. Herrick. Comstock Publishing Co., Ithaca, New York, 1935. pp. 417, illus. 321, indexes, numerous bibliographies. \$5.00.

Intended primarily for people interested, either professionally or as amateurs, in the preservation of shade trees, this book forms a complete manual. Methods of repairing injured trees, materials and apparatus for the control of insect and other pests, suggestions for the fertilization of the soil, are all discussed clearly and comprehensively. Valuable notes on the

relative utility of shade trees and their susceptibility to pests are included. The American Elm heads the list for general utility, while the Ginkgo is first as far as freedom from insect pests is concerned, a fact which may account in part for the long life history of this interesting tree which is often referred to as a "living fossil" for it has existed unchanged for many millions of years and is now unknown in a wild state, being found in China and Japan in the vicinity of palaces, shrines and tombs where it is cultivated. This is a book which anyone interested in trees would do well to obtain.—D. L.



Leechman, Douglas. 1936. "Insect Enemies of Shade Trees, by Glen W. Herrick [Review]." *The Canadian field-naturalist* 50(5), 93–93.

<https://doi.org/10.5962/p.339911>.

View This Item Online: <https://www.biodiversitylibrary.org/item/89296>

DOI: <https://doi.org/10.5962/p.339911>

Permalink: <https://www.biodiversitylibrary.org/partpdf/339911>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Ottawa Field-Naturalists' Club

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.