

TROPIDORHYNCHUS CITREOGULARIS, Gould.

Fig. 1.

Trop. summo capite, dorso, uropygio, alis, caudâque brunneis, his pallidioribus, pogoniis externis remigum secundariorum olivaceo marginatis; caudâ ad apicem griseâ; nuchâ lateribus colli albescenti-griseis; mandibulâ inferiori ad basin, notâque nudâ pone oculos cœruleis; gulâ et lateribus pectoris citreis; abdomine pallidè griseo; rostro nigro; pedibus plumbeis.

Long. tot. $10\frac{1}{4}$ unc.; rostri, $1\frac{1}{4}$; alæ, $5\frac{1}{4}$; caudæ, $4\frac{1}{2}$; tarsi, $1\frac{1}{8}$.

Crown of the head, back, rump, wings, and tail brown, the two latter being lighter, and the secondaries margined externally with olive; tail slightly tipped with greyish white; base of the lower mandible, space beneath the eye, and the region of the ears naked and of a fine blue; throat and sides of the chest lemon yellow; under surface light grey; bill black; legs lead colour.

Tropidorhynchus citreogularis, Gould in Proc. of Zool. Soc., Part V., 1837.

Habitat. Australia: locality, interior of New South Wales.

TROPIDORHYNCHUS CORNICULATUS, Vig. & Horsf.

Fig. 2.

Trop. capite, collo superiore, gutturisque torquè angustocœruleis, nudis; mento, pectore rectricumque apicibus albis, hoc graciliter fusco lineato; corpore inferiore brunnescenti-griseo, subtus albescente; rostro et pedibus brunneo-nigris.

Long. tot. 13 unc.; rostri, $1\frac{5}{8}$; alæ, 6; caudæ, $5\frac{1}{2}$; tarsi, $1\frac{1}{4}$.

Crown of the head, back, and sides of the neck naked and of a blue colour; all the upper surface, wings, and tail greyish brown, the feathers of the latter tipped with greyish white; throat, chest, and under surface greyish brown; bill and feet brownish black.

SYN. *Merops corniculatus*, Lath. Ind. Orn., p. 276. No. 21.

Corbi calao, Le Vaill. Ois. d'Am. et des Indes, tom. i. p. 69. pl. 24.

Knobfronted Honey-Eater, Lath. Gen. Hist., vol. iv. p. 161. No. 6.

Tropidorhynchus corniculatus, Vig. and Horsf. in Linn. Trans., vol. xv. p. 324.—Jard. and Selb. Ill. Orn., vol. iii. pl. cxxxiii.

————— *Monachus*, Vig. and Horsf. in Ibid., vol. xv. p. 324.

Merops Monachus, Lath. Ind. Orn., Supp., p. xxxiv. No. 10.

Knobfronted Bee-Eater, White's Journ., pl. in p. 190.

Cowled Honey-Eater, Lath. Gen. Hist., vol. iv. p. 162. No. 7.

} Young?

Habitat. Australia: locality, New South Wales.



1. *Tropidorhynchus atreogularis*.
2. ----- *corniculatus*.



Gould, John and Gould, Elizabeth. 1837. "Tropidorhynchus corniculatus, Vig. & Horsf." *A synopsis of the birds of Australia, and the adjacent islands* (1), -. <https://doi.org/10.5962/p.363586>.

View This Item Online: <https://www.biodiversitylibrary.org/item/172464>

DOI: <https://doi.org/10.5962/p.363586>

Permalink: <https://www.biodiversitylibrary.org/partpdf/363586>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.