

NANODES VENUSTUS, *Vig.* & *Horsf.*

Fig. 1.

Nan. vittâ cœruleâ frontali, suprâ metallicè cœruleo angustè marginatâ ; loris lineâque supra poneque oculos splendide flavis ; vertice, dorso, uropygio, tectricibus caudæ, gulâ, pectore, lateribusque olivaceo-viridibus ; humeris tectricibusque alarum intensè cœruleis ; primariis nigris, tribus vel quatuor primis ad marginem viridi lavatis ; abdomine medio crisoque flavis ; rectricibus caudæ quatuor intermediis viridescenti-cœruleis, reliquis ad basin externè cœruleis flavo terminatis ; rostro pedibusque fuscis.

Long. tot. $8\frac{1}{2}$ unc. ; alæ, $4\frac{1}{8}$; caudæ, 5 ; tarsi, $\frac{1}{2}$.

A conspicuous bar of deep indigo blue across the forehead, bordered above by a narrow edging of light metallic blue; lores, and a stripe above and behind the eye, rich yellow; crown of the head, back, rump, upper tail-coverts, throat, chest, and flanks brownish olive green; shoulders and wing-coverts deep blue; primaries black, the outer edges of the first three or four slightly tinged with green; centre of the abdomen and under tail-coverts yellow; four middle tail-feathers greenish blue, the basal portions of the remainder beautiful blue on their outer edges, and largely tipped with fine yellow; bill and feet brown.

SYN. *Psittacus venustus*, Temm. in Linn. Trans., vol. xiii. p. 121.

— *chrysostomus*, Kuhl. Conspl. Psitt. in Nov. Act., &c., vol. x. p. 50. No. 78. t. 1.

Blue-banded Parrot, Lath. Gen. Hist., vol. ii. p. 188.

Nanodes venustus, Vig. and Horsf. in Linn. Trans., vol. xv. p. 278.

Remark. The female or young of this species I have not been able satisfactorily to determine; but judging from specimens in my possession, they approach very nearly to the female of the preceding, with this exception, that there is a slight indication of the bar on the forehead.

Habitat. Australia: locality, New South Wales.

NANODES ELEGANS, *Gould.*

Fig. 2.

MAS. *Nan. vittâ frontali purpureâ, suprâ lineâ metallicè cœruleâ marginatâ ad auriculas tendente ; loro splendide flavo ; capite, genis, dorso, tectricibusque caudæ olivaceo-viridibus aureo lavatis ; humeris cœruleis, primariis nigris, primis quatuor ad marginem viridescentibus ; secundariis alâque spuriâ nigris ; gulâ pectoreque viridescenti-flavis, hâc colore in flavum, abdomine crisoque transeunte ; abdomine centrali pallidè aurantiaco ; rectricibus caudæ duabus intermediis viridescenti-cœruleis, reliquis ad basin cœruleis, amplè flavo terminatis ; rostro pedibusque intensè fuscis.*

FœM. vel MAS JUNIOR vittam frontalem caret, et colorem habet indistinctiorem.

Long. tot. 9 unc. ; alæ, $4\frac{5}{8}$; caudæ, $5\frac{1}{4}$; tarsi, $\frac{1}{2}$.

MALE. A conspicuous bar of deep indigo blue across the forehead, bordered above by a narrow edging of light metallic blue, which is continued over the eye to the ear-coverts; space between the bill and the eye rich yellow; head, cheeks, and back, and upper tail-coverts olive green with a golden tinge; shoulders blue, primaries, secondaries, and spurious wing black; outer edges of the four first quills slightly tinged with green; throat and chest greenish yellow passing into bright yellow on the abdomen and under tail-coverts; centre of the abdomen pale orange; two middle tail-feathers greenish blue, the remainder blue at the base and largely tipped with yellow; bill and feet dark brown.

FEMALE OR YOUNG. Wants the bar on the forehead, and has only a faint indication of the other markings.


Nanodes elegans, Gould in Proc. of Zool. Soc., Part V., 1837.

Habitat. Australia: localities, New South Wales and Van Diemen's Land?

NANODES PULCHELLUS, *Vig.* & *Horsf.*

Fig. 3.

MAS. *Nan. lineâ supereiliari, genis, humeris, tectricibusque alarum minoribus, metallicè viridescenti-cœruleis ; vertice, nuchâ, corpore superiore lateribusque nitidè olivaceo-viridibus ; ad originem alarum notâ castaneâ ; primariis secundariisque externè cœruleis, internè fuscis ; pectore, abdo-*



1. *Nanodes venustus*

2. *elégans*

3. *mitchellii*



BHL

Biodiversity Heritage Library

Gould, John and Gould, Elizabeth. 1837. "Nanodes elegans, Gould." *A synopsis of the birds of Australia, and the adjacent islands* (2), --.

<https://doi.org/10.5962/p.363620>.

View This Item Online: <https://www.biodiversitylibrary.org/item/172464>

DOI: <https://doi.org/10.5962/p.363620>

Permalink: <https://www.biodiversitylibrary.org/partpdf/363620>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.