

ACANTHIZA REGULOIDES, Vig. & Horsf.

Fig. 1.

Ac. vertice, nuchâ, corpore superiore, alisque olivaceo-fuscis; uropygio, caudæ tectricibus, crissoque pallidè olivaceis; gulâ, pectoreque albis fusco adpersis; rectricibus caudæ fuscis, ad basin pallidè cervinis, omnium apicibus, externarumque pogoniis externis, fusciscenti-cervinis; rostro pedibusque fuscis.

Long. tot. $3\frac{3}{4}$ unc.; rostri, $\frac{1}{2}$; alæ, 2; caudæ, $1\frac{5}{8}$; tarsi, $\frac{3}{4}$.

Crown, back of the neck, upper surface, and wings olive brown, the feathers of the forehead tipped with a lighter colour; rump, upper and under tail-coverts pale ochre; throat and chest white, each feather with a very slight, broken margin of brown; base of all the tail-feathers pale buff, the external margin of the outer feathers and the tips of all brownish buff, the central portion blackish brown bill and feet brown.

SYN. *Acanthiza reguloides*, Vig. and Horsf. in Linn. Trans., vol. xv. p. 226.

Dwarf Warbler, var. β ? Lath. Gen. Hist., vol. vii. p. 135, no. 161.

Habitat. Australia: locality, New South Wales.

ACANTHIZA UROPYGIALIS, Gould.

Fig. 2.

Ac. capite, corpore suprâ, alisque fuscis, levitèr olivaceo lavatis; uropygio tectricibusque caudæ lætè castaneis; caudâ nigrescenti-fuscâ, latè ad apicem albo notatâ; gulâ, pectore, abdomineque medio griseis; lateribus, crissoque pallidè cervinis; rostro, pedibusque nigris.

Long. tot. $3\frac{3}{4}$ unc.; rostri, $\frac{1}{2}$; alæ, 2; caudæ, $1\frac{3}{4}$; tarsi, $\frac{3}{4}$.

Head, upper surface, and wings brown, slightly tinged with olive; the feather on the forehead tipped with a lighter colour; rump and upper tail-coverts rich reddish chestnut; tail-feathers brownish black, largely tipped with white, which on the two centre feathers is tinged with brown; throat, and chest, and centre of the abdomen greyish white; flanks and under tail-coverts buffy white; bill and feet black.

SYN. *Acanthiza uropygialis*, Gould in Proc. of Zool. Soc., Part V., 1837.

Habitat. Australia: locality, New South Wales.

ACANTHIZA CHRYSORRHŒA.

Fig. 3.

Ac. fronte nigro albo adperso; genis, gulâ, lineâque superciliari cinerescenti-albis; pectore, corporeque subtis flavescenti-albis; lateribus olivaceis; corpore superiore, alisque olivaceo-fuscis; uropygio, tectricibusque caudæ citrinis, rectricibus caudæ in medio fuscis, ad basin flavescenti-albis; externarum pogoniis externis, omniumque apicibus cineraceis; rostro, pedibusque nigrescentibus.

MAS. Long. tot. $4\frac{1}{4}$ unc.; rostri, $\frac{5}{8}$; alæ, $2\frac{5}{8}$; caudæ, $1\frac{3}{4}$; tarsi, $\frac{3}{4}$.

Forehead black, with a spot of white at the tip of each feather; cheeks, throat, and a line from the nostrils over each eye greyish white; chest and under surface yellowish white, passing into light olive brown on the flanks; upper surface and wings olive brown; rump and upper tail-coverts bright citron yellow; base of the tail-feathers white, tinged with yellow, the external margin of the outer feathers and the tips of all brownish grey, the central portion blackish brown; bill and feet blackish brown.

SYN. *Saxicola chrysorrhæa*, Quoy et Gaim. Voy. de l'Astrolabe, p. 198, pl. 10, f. 2.

Habitat. Van Diemen's Land and New South Wales.

ACANTHIZA MAGNIROSTRA, Gould.

Fig. 4.

Ac. vertice, corpore superiore, alis caudâque olivaceo-fuscis; hâc, fronteque rufescentibus; gulâ pectoreque cinereis; lateribus olivaceis; rostro nigro; pedibus brunneis.

Long. tot. $4\frac{3}{4}$ unc.; rostri, $\frac{3}{4}$; alæ, $2\frac{1}{4}$; caudæ, $1\frac{7}{8}$; tarsi, $\frac{3}{4}$.

Crown of the head, all the upper surface, wings and tail olive brown, the forehead and tail becoming rufous brown; throat and chest brownish white; abdomen greyish white, passing into bright olive green on the lower part of the flanks; bill black; feet light brown.

SYN. *Acanthiza magnirostra*, Gould in Proc. of Zool. Soc., Part V., 1837.

Habitat. Australia: locality, New South Wales.



1



2



3



4

1. *Acanthiza reguloides*.
 2. ----- *uropygialis*.
 3. ----- *chrysorrhoa*.
 4. ----- *magnirostra*.



Gould, John and Gould, Elizabeth. 1838. "Acanthiza uropygialis, Gould." *A synopsis of the birds of Australia, and the adjacent islands* (4), -. <https://doi.org/10.5962/p.363696>.

View This Item Online: <https://www.biodiversitylibrary.org/item/172464>

DOI: <https://doi.org/10.5962/p.363696>

Permalink: <https://www.biodiversitylibrary.org/partpdf/363696>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.