

Resedaceae (Mignonette Family)

Oligomeris linifolia (M. Vahl) J.F. Macbr. - Oligomeris. Uncommon, gravelly benches and sandy flats near Kelso Dunes, below 900 m. B.A. Prigge 2608 (UCR).

Rhamnaceae (Buckthorn Family)

Rhamnus ilicifolia Kellogg - Holly-leaved Red-berry. Infrequent, bouldery slopes in pinyon-juniper woodland, 1200-2100 m. J. André 3734 (GMDRC).

Rosaceae (Rose Family)

Amelanchier utahensis Koehne - Utah Service Berry. Uncommon, moist north-facing bouldery slopes, Cottonwood Basin and Snake Spring, 1300-1600 m. J. Evens 43 (GMDRC).

Coleogyne ramosissima Torrey - Blackbush. Common, dominating upper bajadas and plateaus, 1000-2000 m. J. André 3865 (GMDRC).

Fallugia paradoxa (D. Don) Endl. - Apache Plume. Infrequent, banks of canyon drainages and among boulders, 1200-2100 m. K. Hart 33 (GMDRC).

Holodiscus microphyllus Rydb. var. *microphyllus* - Rock Spiraea. Infrequent, bouldery slopes in pinyon-juniper woodland, 1800-2100 m. J. André 3943 (GMDRC).

Ivesia saxosa (E. Greene) B. Ertter - Rock Cinquefoil. Uncommon, crevices in cliff faces and boulders, 900-2100 m. J. André 3735 (GMDRC).

Prunus fasciculata (Torrey) A. Gray var. *fasciculata* - Desert Almond. Locally common, washes, occasionally on slopes, 1000-2000 m. J. André 2121 (GMDRC).

Purshia tridentata (Pursh.) DC. var. *glandulosa* (Curran) M.E. Jones - Mojave Antelope Bush. Frequent, ridges and slopes in woodland, 1300-2100 m. J. André 2102 (GMDRC).

Rubiaceae (Madder Family)

!*Galium aparine* L. - Goose Grass. Rare, in shady areas around springs on east side of range 1200-1500 m. J. Evens 98 (GMDRC).

**Galium munzii* Hilend & J. Howell - Munz's Bedstraw. Rare, cool north-facing canyon slopes, Cottonwood Basin, 1300-1600 m. B.A. Stein 36 (RSA)

Galium parishii Hilend & J. Howell - Parish's Bedstraw. Infrequent, steep rocky slopes and shady canyon bottoms, 1600-2100 m. Bradbury 286 (UC).

Galium stellatum Kellogg var. *eremicum* Hilend & J. Howell - Desert Bedstraw. Common, rocky slopes and shady canyon bottoms, 1000-2000 m. J. André 3698 (GMDRC).

Rutaceae (Rue Family)

Thamnosma montana Torrey & Fremont - Turpentine Bush. Common, gravelly bajadas in diverse shrub associations, 800-1400 m. R.F. Thorne 45381 (UCR).

Salicaceae (Willow Family)

- Populus fremontii* S. Watson ssp. *fremontii* - Fremont Cottonwood. Uncommon, from four localities: Cove and Cottonwood Springs, Bull and Willow Spr Cyns, 1300-1800 m. R.F. Thorne 45471 (UC).
- Salix exigua* Nutt. - Narrow-leaved Willow. Infrequent, clonal occurrences in deep canyon drainages and springs, 1100-1900 m. J. André 2130 (GMDRC).
- Salix gooddingii* C. Ball - Goodding's Black Willow. Rare, springs and seeps in canyons and on rocky slopes, 1200-1500 m. R.F. Thorne 45421 (UC).
- Salix laevigata* Bebb - Red Willow. Rare, moist north-facing slopes, Snake Spring (may be *S. gooddingii* mis-identified). W.L. Jepson 20532 (JEPS)
- Salix lasiolepis* Benth. - Arroyo Willow. Uncommon, rocky slopes associated with sidehill seeps, upper Granite Cove, 1600 m. J. Evens 102 (GMDRC).

Saururaceae (Lizard's-tail Family)

- Anemopsis californica* (Nutt.) Hook & Arn. - Yerba Mansa. Rare, reported from lower Bull Canyon, 1200 m. (photograph and leaf material verified by J. André 1999).

Scrophulariaceae (Figwort Family)

- Antirrhinum filipes* A. Gray - Yellow Twining Snapdragon. Occasional, on shrubs and debris, mostly in washes, below 1400 m. J. André 3955 (GMDRC).
- Castilleja angustifolia* (Nutt.) G. Don - Desert Indian Paintbrush. Occasional, upper bajadas in mixed scrub, or wooded slopes, 1200-2000 m. G.L. Clifton 24411 (GMDRC).
- Castilleja linariifolia* Benth. - Linear-leaved Paintbrush. Infrequent, gravelly bajadas, washes and rocky slopes, 1000-2100 m. B. Pitzer 4020 (UCR).
- Keckiella antirrhinoides* (Benth.) Straw var. *microphylla* (A. Gray) N. Holmgren - Desert Chaparral Beard-tongue. Infrequent, bouldery slopes, pinyon-juniper woodland, 1200-1900 m. R.F. Thorne 45443 (UC).
- Mimulus bigelovii* (A. Gray) A. Gray var. *bigelovii* - Bigelow's Monkeyflower. Occasional, sandy washes and moist canyon drainages, 800-1700 m. J. Evens 27 (GMDRC).
- Mimulus guttatus* DC. - Common Large Monkeyflower. Infrequent, moist banks of canyon drainages, seeps and springs, 1000-1800 m. B. Pitzer 3899 (UCR).
- Mimulus parishii* E. Greene - Parish's Monkeyflower. Rare, known only from wet sandy banks of Bull Creek, upper Bull Canyon, 1800 m. F. Warrick 80 (GMDRC).
- Mimulus pilosus* (Benth.) S. Watson - Downy Monkeyflower. Uncommon, streambeds or moist disturbed areas, mainly east side, 1300-1600 m. J. André 5182 (GMDRC).
- Mimulus rubellus* A. Gray - Ciliolate-toothed Monkeyflower. Uncommon, sandy slopes and banks of washes, 800-1900 m. F. Warrick 27 (GMDRC).
- Penstemon eatonii* A. Gray var. *undosus* M.E. Jones - Eaton's Firecracker. Frequent, along rocky washes and slopes, 1300-2100 m. J. André 3870 (GMDRC).
- Penstemon palmeri* A. Gray var. *palmeri* - Palmer's Penstemon. Infrequent, gravelly areas along Kelbaker Rd, Granite Pass to Kelso Dunes, 800-1300 m. J. André 3692 (GMDRC).

Penstemon pseudospectabilis M.E. Jones - Desert Penstemon. Occasional, among boulders, mostly east side, 1300-1800 m. *J. André 5087 (UCR)*. (intergrades with *P. stephensii*).

Penstemon rostriflorus Kellogg - Bridge's Penstemon. Rare, known only from rocky north-facing slope near summit of Silver Peak at 2000 m. *J. André 3821 (GMDRC)*.

**Penstemon stephensii* Brandegee - Stephens' Beardtongue. Rare, among boulders east side of range, 1300-2000 m. *J. Evens 77 (GMDRC)*. (intergrades with *P. pseudospectabilis*).

Solanaceae (Nightshade Family)

Datura wrightii Regel - Jimson Weed. Occasional, in sandy washes and along roadsides, 800-1300 m. *J. André 5708 (GMDRC)*.

Lycium andersonii A. Gray - Anderson's Desert-thorn. Frequent, sandy bajadas and moderate slopes, 1000-1900 m. *J. André 3869 (GMDRC)*.

Lycium cooperi A. Gray - Cooper's Desert-thorn. Frequent, gravelly bajadas in diverse mixed shrub associations, and slopes below 1600 m. *K. Hart 13 (GMDRC)*.

Nicotiana obtusifolia Martens & Galeotti - Desert Tobacco. Occasional, in crevices and cavities in rocks, wash margins, below 1500 m. *J. André 2124 (GMDRC)*.

Physalis crassifolia Benth. - Thick-leaved Ground Cherry. Frequent, sandy and rocky places, mostly on west and south side of range, below 1400 m. *G.L. Clifton 25200 (GMDRC)*.

Physalis hederifolia A. Gray var. *fendleri* (A. Gray) Cronq. - Ivy-leaved Ground Cherry. Infrequent, margins of washes or among rocks, 900-1900 m. *J. André 4573 (GMDRC)*.

Tamaricaceae (Tamarisk Family)

!*Tamarix ramosissima* Ledeb. - Tamarisk. Uncommon, canyon bottoms, springs and seeps, 1000-1600 m. Eradication and control program is ongoing. *B.A. Stein 76 (GMDRC)*.

Ulmaceae (Elm Family)

+*Ulmus pumila* L. - Siberian Elm. Uncommon, planted at historic homesteads but has not escaped here (has elsewhere in Mojave Desert). (*obs. J. André 1994*).

Urticaceae (Nettle Family)

Urtica dioica L. ssp. *holosericea* (Nutt.) Thorne - Hoary Nettle. Rare, one location on east side of range, in moist seep under a granitic overhang, 1350 m. *J. Evens 41 (GMDRC)*.

Viscaceae (Mistletoe Family)

- Arceuthobium divaricatum* Engelm. - Pinyon Dwarf Mistletoe. Occasional, woodlands on *P. monophylla*, 1400-2100 m. Infrequently collected. (obs. J. André 1998).
Phoradendron californicum Nutt. - Desert Mistletoe. Frequent on *Acacia* and *Prosopis*, 800-1500 m. Norris s.n. (GMDRC).
Phoradendron juniperinum A. Gray - Juniper Mistletoe. Scattered throughout range, but infrequent on *Juniperus*, 1200-2100 m. J. André 3712 (GMDRC).

Zygophyllaceae (Caltrop Family)

- Kallstroemia californica* (S. Watson) Vail - California Kallstroemia. Rare, sandy areas in washes and on flats, only from southeast side, 1250 m. J. André 6650 (UCR).
Larrea tridentata (Sessé & Moq. Ex DC.) Cov. - Creosote Bush. Common dominant, sandy or gravelly bajadas or south-facing slopes, 700-1500 m. B. Pitzer 3965 (UCR).
Tribulus terrestris L. - Puncture Vine. Rare pernicious weed, along roadways and other disturbed areas, 900-1300 m. J. André 5944 (GMDRC).

CLASS II. MONOCOTYLEDONEAE - Monocotyledons

Cyperaceae (Sedge Family)

- Carex alma* L. Bailey - Sturdy Sedge. Occasional, seeps and springs in approximately 40 localities throughout range, 1100-1900 m. J. André 3445 (GMDRC).
Eleocharis parishii Britton - Parish's Spike-rush. Infrequent, seeps and springs and abandoned livestock improvements, 1000-1800 m. J. André 3746 (GMDRC).
Scirpus pungens Vahl. - Common Threesquare. Uncommon, seeps and canyon watercourses, Bull and Budweiser Canyons, Coyote Springs, 1200-1700 m. J. André 4099 (GMDRC).

Iridaceae (Iris Family)

- Sisyrinchium halophilum* E. Greene - Nevada Blue-eyed Grass. Uncommon, seeps and springs in approximately 10 localities throughout range, 1100-1800 m. J. Evens 60 (GMDRC).

Juncaceae (Rush Family)

- Juncus bufonius* L. var. *bufonius* - Common Toad Rush. Uncommon, seeps, springs and watercourses, 1200-1700 m. B.A. Prigge 2863 (UCR).
Juncus bufonius L. var. *occidentalis* F.J. Herm. - Western Toad Rush. Rare, canyon seep at base of Budweiser Canyon, 1300 m. J. André 4086 (GMDRC).
Juncus macrophyllus Cov. - Large-leaved Rush. Rare, canyon seep at base of Carr and Budweiser Canyons, 1200 -1400 m. J. André 4156 (GMDRC).
Juncus mexicanus Willd. - Mexican Rush. Infrequent, seeps and springs and moist canyon bottoms, 1100-1800 m. J. André 4087 (GMDRC).

**Juncus nodosus* L. - Knotted Rush. Rare, known only from Coyote Spring on east side of range, 1200 m. *J. André 4046 (GMDRC)*.

Juncus xiphioides E. Meyer - Iris-leaved Rush. Infrequent, seeps and springs in approximately 30 localities throughout range, 1100-1900 m. *J. André 4584 (GMDRC)*.

Liliaceae (Lily Family)

Agave deserti Engelm. - Desert Agave. Infrequent, rocky slopes, scattered throughout east side of range, 1200-1700 m. *C.B. Wolf 10183 (RSA)*.

**Allium atrorubens* S. Watson var. *cristatum* (S. Watson) D. McNeal - Dark Red Onion. Rare, only on ridgetops, west side of range, 1800-2000 m. *J. André 4096 (GMDRC)*.

Calochortus kennedyi Porter var. *kennedyi* - Kennedy's Mariposa. Occasional, dry slopes and gravelly bajadas on far east side of range, 1100-1400 m. *J. André 3693 (GMDRC)*.

Calochortus kennedyi Porter var. *munzii* Jepson - Munz's Red Mariposa. Rare, open flats in Bull Canyon, 1450 m. *B.A. Stein s.n. (UCSC)*.

Dichelostemma capitatum (Benth.) A.W. Wood ssp. *pauciflorum* (Torrey) Keator - Desert Blue-dicks. Occasional, gravelly bajadas, east side, 1100-1600 m. *K. Hart 32 (GMDRC)*.

Hesperocallis undulata A. Gray - Desert Lily. Uncommon, aeolian sand deposits near Kelso Dunes, 800-900 m. *J. André 4213 (GMDRC)*.

Muilla transmontana E. Greene - Great Basin Muilla. Rare, sandy flats, aeolian sand deposits near Kelso Dunes, 800 m. *J. André 3628 (GMDRC)*. Note: this is a unique form that is a southern disjunct and very small-flowered perhaps warranting further evaluation.

Yucca baccata Torrey - Spanish Bayonet. Frequent, dry rocky slopes and plateaus, mostly in pinyon-juniper woodland, 1300-2100 m. *J. André 3621 (GMDRC)*.

+*Yucca brevifolia* Engelm. - Joshua Tree. Planted as ornamental in Granite Cove (1400 m), no natural populations exist in range. (*obs. J. André 1994*).

Yucca schidigera K.E. Ortgies - Mojave Yucca. Common, gravelly bajadas in diverse mixed scrub, 1000-1800 m. *J. André 5839 (GMDRC)*.

Zigadenus brevibracteatus (M.E. Jones) H.M. Hall - Desert Death Camas. Occasional, sandy bajadas, especially on east side of range, 1000-1400 m. *J. André 5061 (UCR)*.

Poaceae (Grass Family)

Achnatherum hymenoides (Roemer & Schultes) Barkworth - Indian Ricegrass. Infrequent, mainly on aeolian deposits near Kelso Dunes, 800-1000 m. *J. André 4198 (GMDRC)*.

Achnatherum parishii (Vasey) Barkworth - Parish's Needlegrass. Uncommon, gravelly or rocky slopes, 1300-2100 m. *J. André 5986 (GMDRC)*.

Achnatherum speciosum (Trin. & Rupr.) Barkworth - Desert Needlegrass. Common, rocky slopes, plateaus and gravelly pediments, 1000-2100 m. *C.B. Wolf 10118 (UC)*.

- ! *Agrostis semiverticillata* (Forsskal) C. Chr. - European Needlegrass. Uncommon, disturbed or moist areas (Coyote Spring), 1200-1500 m. J. André 3899 (GMDRC).
- Aristida adscensionis* L. - Six-weeks Three-awn. Frequent, dry open bajadas and slopes among boulders, 900-1400 m. J. André 3987 (GMDRC).
- Aristida purpurea* Nutt. var. *fendleriana* (Steudel) Vasey - Fendler Three-awn. Frequent, rocky slopes, crevices, gravelly ridgetops, 1100-2100 m. J. Evens 90 (GMDRC).
- Aristida purpurea* Nutt. var. *nealleyi* (Vasey) K.W. Allred - Neally Three-awn. Infrequent, dry south-facing slopes, 1000-1600 m. B.A. Prigge 2349 (RSA).
- Aristida purpurea* Nutt. var. *parishii* (A. Hitchc.) K.W. Allred - Parish Three-awn. Infrequent, gravelly or rocky slopes, 1000-1500 m. R.F. Thorne 50670 (RSA).
- Aristida ternipes* Cav. var. *hamulosa* (Henrard) J.S. Trent - Hook Three-awn. Rare, dry hills near Budweiser Wash at 1300 m. B.A. Prigge 2321 (RSA).
- Bothriochloa barbinodis* (Lagasca) Herter - Cane Bluestem. Uncommon, crevices in rocks, rocky shaded canyon bottoms, 1200-1400 m. J. André 3756 (GMDRC).
- Bouteloua aristidoides* (Kunth.) Griseb. var. *aristidoides* - Needle Grama. Frequent, bajadas, flats and gravelly slopes, 800-1700 m. R.F. Thorne 50627 (RSA).
- Bouteloua barbata* Lagasca var. *barbata* - Six-weeks Grama. Common after summer rains, washes, sandy bajadas, lower slopes, below 1600 m. B. Pitzer 4006 (UCR).
- ! *Bromus diandrus* Roth - Ripgut Grass. Uncommon, disturbed areas around historic settlements and livestock structures, mainly west side of range, 1100-1400 m. B. Pitzer 3917 (UCR).
- ! *Bromus hordeaceus* L. - Soft Chess. Uncommon, three isolated occurrences in disturbed sites on east side of range, 1200-1400 m. J. André 3623 (GMDRC).
- ! *Bromus madritensis* L. ssp. *rubens* (L.) Husnot - Foxtail Chess, Red Brome. Common, ubiquitous weed, widespread in all habitats, 700-2100 m. B.A. Stein 85 (GMDRC).
- ! *Bromus tectorum* L. - Cheat Grass. Common, ubiquitous weed, widespread in all habitats, 700-2100 m. J. André 5919 (GMDRC).
- Bromus trinii* Desv. - Chilean Chess. Uncommon, sandy bajadas or washes, growing in shade of large shrubs, 1200-1700 m. J. André 4908 (UCR). (mistakenly reported as non-native in Jepson Desert Manual).
- ! *Cynodon dactylon* (L.) Pers. - Crab Grass. Uncommon, moist seeps or subalkaline soils of disturbed areas (Willow Spr. Basin), 1200-1400 m. J. André 3744 (GMDRC).
- Elymus elymoides* (Raf.) Swezey ssp. *brevifolius* (J.G. Smith) Barkworth - Squirreltail. Uncommon, dry open slopes and ridgetops, 1800-2000 m. F. Warrick 75 (GMDRC).
- Elymus multisetus* (J.G. Smith) Burt Davy - Big Squirreltail. Uncommon, ridgetops and rocky crags, 1800-2150 m. F. Warrick 74 (GMDRC).
- Elymus trachycaulus* (Link) Shinn. ssp. *trachycaulus* - Slender Wheatgrass. Uncommon, rocky slopes and canyon walls, Bull Canyon only, 1400-1900 m. R.F. Thorne 51609 (RSA).
- ! *Elytrigia intermedia* (Host) Nevski ssp. *intermedia* - Intermediate Wheatgrass. Rare, unconfirmed report by G.L. Clifton, 2001.
- ! *Eragrostis cilianensis* (All.) Janchen - Stinkgrass. Infrequent, disturbed soils or in post fire recovery on hillsides, 1200-1800 m. B. Pitzer 4016 (UCR).

- Erioneuron pulchellum* (Kunth.) Tateoka - Fluff Grass. Frequent, sandy or gravelly bajadas and pediments, 800-1800 m. *Dorothy Fugle 1045 (UCSB)*.
- ! *Hordeum murinum* L. ssp. *murinum* - Wall Barley. Uncommon, disturbed areas along roadsides and buildings in Granite Cove, 1300 m. *J. André 4051 (GMDRC)*.
- Melica frutescens* Scribner - Tall Melica. Uncommon, bouldery slopes, pinyon-juniper woodland, 1300-1700 m. *Al Flink s.n. (UCSB)*.
- Melica imperfecta* Trin. - Small-flowered Melica. Infrequent, rocky slopes and bouldery areas in pinyon-juniper woodland, 1300-1600 m. *G.L. Clifton 24771 (GMDRC)*.
- Muhlenbergia asperifolia* (Nees & Meyen) L. Parodi - Scratchgrass. Rare, in Bull Cyn on moist sandy margins of creekbed, 1100 m. *J. André 3628 (GMDRC)*.
- Muhlenbergia microsperma* (DC.) Trin. - Littleseed Muhly. Uncommon, dry slopes and open sites on pediments, Willow Springs Basin, 1100-1200 m. *J. André 3954 (GMDRC)*.
- Muhlenbergia porteri* Beal. - Porter's Muhly. Frequent, crevices on cliffs and on steep rocky slopes, 900-1800 m. *S. Boyd 5143 (RSA)*.
- Muhlenbergia rigens* (Benth.) A. Hitchc. - Deergrass. Locally common, moist sandy margins of streams, canyon bottoms, 1300-2100 m. *B.A. Prigge 2323 (UCR)*.
- Phragmites australis* (Cav.) Steudel - Common Reed. Uncommon, perennial moist sand in canyon bottoms and near seeps, 1300-1800. *J. André 3742 (GMDRC)*.
- Pleuraphis rigida* Thurber - Big Galleta. Uncommon, sandy bajadas and aeolian deposits near Kelso Dunes, 700-1300 m. *G.L. Clifton 31216 (GMDRC)*.
- ! *Poa annua* L. - Annual Bluegrass. Rare waif, in moist shady drainage above Coyote Spring, and near developments in Granite Cove, 1200-1400 m. *J. André 5994 (GMDRC)*.
- Poa bigelovii* Vasey & Scribner - Bigelow's Bluegrass. Uncommon, slopes among boulders and understory of pinyon woodland, 1300-1600 m. *G.L. Clifton 11821 (GMDRC)*.
- Poa fendleriana* (Steudel) Vasey ssp. *longiligula* (Scribner & Williams) R. Soreng - Longtongue Mutton Grass. Uncommon, rocky north-facing slopes, 1500-2100 m. *G.L. Clifton 24374 (GMDRC)*.
- Poa secunda* J.S. Presl ssp. *secunda* - One-sided Bluegrass. Frequent, rocky or gravelly slopes and ridgetops, 1000-2100 m. *J. André 3939 (GMDRC)*.
- ! *Polypogon monspeliensis* (L.) Desf. - Annual Beard Grass. Uncommon, moist sandy soils near seeps and springs, 1100-1900 m. *F. Warrick 109 (GMDRC)*.
- ! *Schismus barbatus* (L.) Thell. - Mediterranean Grass. Common, sandy bajadas and flats, disturbed areas, 700-1400 m. *J. André 2099 (GMDRC)*.
- Sporobolus cryptandrus* (Torrey) A. Gray - Sand Dropseed. Uncommon, rocky to sandy washes, slopes in pinyon-juniper woodland, 1300-2100 m. *J. André 5933 (GMDRC)*.
- Tridens muticus* (Torrey) Nash - Slim Tridens. Uncommon, dry rocky slopes and in pinyon and juniper woodland, 1200-2000 m. *E.W. Lathrop 7804 (UCR)*.
- ! *Vulpia bromoides* (L.) S.F. Gray - Cheat Fescue. Rare waif, mouth of Budweiser Canyon in disturbed soils, 1300 m. *J. Evens 82 (GMDRC)*.
- Vulpia microstachys* (Nutt.) Munro var. *pauciflora* (Beal) Lonard & Gould - Awned Fescue. Uncommon, moist soils (Cottonwood Spring), 1400 m. *G.L. Clifton 26330 (GMDRC)*.

!Vulpia myuros (L.) C. Gmelin var. hirsuta Hack. - Rattail Fescue. Rare, near Cottonwood Spring, 1400 m. (obs. R.F. Thorne et. al. 1981).

Vulpia octoflora (Walter) Rydb. var. octoflora - Slender Fescue. Frequent, washes, sandy bajadas and aeolian sands near Kelso Dunes, 700-1800 m. J. André 4170 (GMDRC).

Typhaceae (Cattail Family)

Typha domingensis Pers. - Southern Cattail. Uncommon, springs and seeps and moist sandy areas in canyon bottoms, 1100-1700 m. J. André 5888 (GMDRC).



Williams, Kimberlyn et al. 2006. "Habitat correlates of *Calochortus plummerae*, a rare mariposa lily, on the campus of California State University, San Bernardino." *Crossosoma* 32(2), 75–82.

View This Item Online: <https://www.biodiversitylibrary.org/item/211111>

Permalink: <https://www.biodiversitylibrary.org/partpdf/370627>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Southern California Botanists

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.