

FOSSIL *ALDROVANDA* — ADDITIONS

DR. J. SCHLAUER

Keywords: evolution: *Aldrovanda*.


Two articles that appeared in the 1960's are not considered in the previous, otherwise accurate paper by Dr. Degreeef. As they contain important additional information they shall be mentioned in brief here.

Weyland, H., Pflug, H. D., Marinos, G., and Anastopoulos, J. 1961, Beiträge zur fossilen Flora des Braunkohlenbeckens von Megalopolis im Peloponnes (Griechenland), *Palaeontographica Abt. B Paläophytologie* 108: 93-120.

Peters, I. 1963, Die Flora der oberpfälzer Braunkohlen und ihre ökologische und stratigraphische Bedeutung, *Palaeontographica Abt. B Paläophytologie* 112: 1-50.

The first paper reports on another Tertiary find. Seeds of *A. megalopolitana* are described (on pp. 104—105, t.18 fig. 10—11) from the Pliocene of south Greece. The seeds have a rather thick testa, and especially noteworthy is an interior layer of cells that are stretched at right angles to the longitudinal axis of the seeds. This character is only observed in one other fossil species, viz. the Late Cretaceous *Palaeoalldrovanda splendens* (cf. Dr. Degreeef's paper). According to the authors, no other fossil *Aldrovanda* seeds are known in which this cell layer is preserved. For completeness, the species *A. megalopolitana* is included in Figures 1 and 2 of Dr. Degreeef's paper.

The second paper is even more intriguing. In this paper (on pp. 29-31, t. 13 fig. 74-76, t. 14 fig. 77-81), fossil laminae of *Aldrovanda inopinata* from the Upper Miocene (approximately 6 MYA) of Wackersdorf (southeast Germany) are described (Figure 1). They are rather similar to the traps of the recent *A. vesiculosa* but differ in their more cuneate base, the longer apical spur with more bristles, the (apparently) missing sensitive hairs in the central portion of the leaves and the (apparently) missing quadrifid trichomes in the marginal zone. It is a pity that this important and rather sensational find is not cited in large portions of the more recent literature dealing with *Aldrovanda* fossils.



Aldrovanda inopinata Peters

Figure 1: *Aldrovanda inopinata* Peters, reconstruction of lamina. Reprinted with permission by Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, Germany.



Schlauer, Jan. 1997. "Fossil Aldrovanda--Additions." *Carnivorous plant newsletter* 26(3), 98–98. <https://doi.org/10.55360/cpn263.js888>.

View This Item Online: <https://www.biodiversitylibrary.org/item/237796>

DOI: <https://doi.org/10.55360/cpn263.js888>

Permalink: <https://www.biodiversitylibrary.org/partpdf/375262>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

IMLS LG-70-15-0138-15

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: International Carnivorous Plant Society

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.