

Review of the infrageneric classification of *Adenanthos* (Proteaceae)— the subsections of *Adenanthos* are polyphyletic

Francis J. Nge^{1,2,5} and Kevin R. Thiele^{3,4}

¹School of Biological Sciences, Faculty of Science,
The University of Adelaide, Adelaide, SA 5000, Australia

²State Herbarium of South Australia, Adelaide, SA 5005, Australia

³Western Australian Herbarium, Biodiversity and Conservation Science,

Department of Biodiversity, Conservation and Attractions,
Locked Bag 104, Bentley Delivery Centre, Western Australia 6983

⁴School of Biological Sciences, University of Western Australia,
35 Stirling Hwy, Crawley WA 6009

⁵Corresponding author, email: francisnge@yahoo.com

SHORT COMMUNICATION

Adenanthos Labill. (Proteaceae) is an endemic Australian genus comprising 31 species (Nge *et al.* 2021). Most species are restricted to southwest Western Australia, with two in South Australia. The genus was divided into two sections, sect. *Eurylaema* Benth. and sect. *Adenanthos* Benth., by Bentham (1870) based on anther and style morphology (Figure 1).

Nelson (1978), in the most recent full revision of the genus, erected a subsection (subsect. *Anaclastos* E.C.Nelson) within sect. *Adenanthos* (thus also establishing the autonymic subsect. *Adenanthos*) based on the single character of perianth length (Figure 1).

Adenanthos sect. *Eurylaema* Benth.

One anther sterile; style-end flattened, much broader than style

(3 species: *A. detmoldii* F.Muell., *A. barbiger* Lindl., *A. obovatus* Labill.)

Adenanthos sect. *Adenanthos*

All anthers fertile; style-end conical, not broader than style

(All remaining species)

subsect. *Anaclastos* E.C.Nelson

Perianth 10–15 mm long

(3 species: *A. apiculatus* R.Br., *A. dobagii* E.C.Nelson, *A. drummondii* Meisn.)

subsect. *Adenanthos*

Perianth > 15 mm long

(All remaining species)

Figure 1. Nelson's (1978) infrageneric classification of *Adenanthos*, with diagnostic characters and species numbers.

Nge *et al.* (2021) have shown that the two sections are reciprocally monophyletic, but the two subsections of sect. *Adenanthos* are polyphyletic, with strong support, in Bayesian and Maximum Likelihood phylogenetic analyses using nuclear and chloroplast gene sequences. The three species of *A.* subsect. *Anaclastos* are nested within other clades comprising taxa belonging to *A.* subsect. *Adenanthos* (Figure 2).

Polyphyly of the subsections in the chloroplast topology could be attributed to introgression and chloroplast capture, which was shown to be very common within *Adenanthos* (Nge *et al.* 2021). However, polyphyly is also strongly supported in analyses based on 35 nuclear gene regions, and these are expected to provide a more accurate estimation of the species phylogeny. The nuclear topologies of Nge *et al.* (2021), from both coalescent and concatenated analyses, are largely congruent with morphological delimitations of species in *Adenanthos*.




Figure 2. The floral character used to delimit *Adenanthos* subsect. *Anaclastos* (perianth length) mapped onto a Maximum Likelihood RAxML phylogram of *Adenanthos*, based on 35 nuclear genes, using the 'anc.ML' function in the phytools package in R (R Core Team 2016). Perianth length (mm) is coloured from blue–red (short–long). The three species in *A.* subsect. *Anaclastos* are highlighted with green squares. Bootstrap support values are indicated at each node (very strongly supported branches [100 bootstrap] are represented by '*', weakly supported branches [< 50 bootstrap] are represented by '-'). The tree was sourced and adapted from Nge *et al.* (2021).

Short and long perianths have both evolved multiple times independently in the genus (Figure 2), presumably as adaptations to different pollinators. The placement of species in the two subsections based on whether the perianth is 10–15 mm long (subsect. *Anaclastos*) or > 15 mm long (subsect. *Adenanthos*) is not supported phylogenetically. Accordingly, the sections, but not the subsections, are accepted here as infrageneric taxa in *Adenanthos* and subsect. *Anaclastos* is reduced to a synonym of sect. *Eurylaema*.

Revised classification

Adenanthos Labill., *Pl. Nov. Holl.* i. 28. t. 36 (1804). *Lecto: Adenanthos cuneatus* Labill., *fide* E.C. Nelson, *Brunonia* 1: 316 (1978) [as *A. cuneata*].

Adenanthos sect. **Eurylaema** Benth., *Fl. Austral.* 5: 350 (1870). *Lecto: Adenanthos obovatus* Labill., *fide* E.C. Nelson, *Brunonia* 1: 322 (1978) [as *A. obovata*].

Adenanthos Labill. sect. **Adenanthos**

Adenanthos subsect. *Anaclastos* E.C. Nelson, *Brunonia* 1(3): 332 (1978), *syn. nov.* *Type: Adenanthos apiculatus* R.Br.

Acknowledgements

We thank Laurence Haegi and Barbara Rye for their comments, which improved the manuscript.

References

- Bentham, G. (1870). *Flora Australiensis*. (Reeve & Co.: London.)
- Nelson, E.C. (1978). A taxonomic revision of the genus *Adenanthos* (Proteaceae). *Brunonia* 1: 303–406.
- Nge, F.J., Biffin E, Thiele, K.R. & Waycott, M. (2021). Reticulate evolution, ancient chloroplast haplotypes, and rapid radiation of the Australian plant genus *Adenanthos* (Proteaceae). *Frontiers in Ecology and Evolution* 8: 616741.
- R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.



Nge, Francis Jason and Thiele, Kevin. 2021. "Review of the infrageneric classification of *Adenanthos* (Proteaceae)—the subsections of *Adenanthos* are polyphyletic." *Nuytsia: journal of the Western Australian Herbarium* 32, 59–61. <https://doi.org/10.58828/nuy01000>.

View This Item Online: <https://www.biodiversitylibrary.org/item/322286>

DOI: <https://doi.org/10.58828/nuy01000>

Permalink: <https://www.biodiversitylibrary.org/partpdf/376403>

Holding Institution

Western Australian Herbarium

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Western Australian Herbarium

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.