١

RELATIONSHIPS BETWEEN FREE CUPRIC ION CONCENTRATIONS IN SEA WATER AND COPPER METABOLISM AND GROWTH IN CRAB LARVAE

BRENDA M. SANDERS^{1,2} AND KENNETH D. JENKINS^{1,3}

¹Molecular Ecology Institute, California State University, Long Beach, California, 90840; ²Duke University Marine Laboratory, Beaufort, North Carolina, 28516; and ³Department of Biology, California State University, Long Beach, California, 90840

ABSTRACT

Crab larvae (*Rhithropanopeus harrisii*) were exposed to a range of free cupric ion concentrations, $[Cu^{2+}]$, regulated in sea water by a metal chelate buffer system. We found a biphasic relationship between intracellular copper distribution and $[Cu^{2+}]$ in sea water. At $[Cu^{2+}]$ within the ambient range $(10^{-12.4} \text{ to } 10^{-10.6} M)$, cytosolic copper was associated with both metallothionein (MT) and high molecular weight (HMW) ligands, and was independent of external $[Cu^{2+}]$. At higher $[Cu^{2+}]$, copper was also associated with very low molecular weight (VLMW) ligands, and accumulated in this ligand pool and the MT pool as external $[Cu^{2+}]$ increased. In marked contrast, copper in the HMW ligand pool did not correlate with $[Cu^{2+}]$ in sea water over the entire range of exposures. Reductions in larval growth occurred at greater than estimated ambient $[Cu^{2+}]$ and correlated with copper accumulation in the MT and VLMW pools.

INTRODUCTION

The concentrations and subcellular distributions of metals can provide valuable information on an organism's capacity to adapt to accumulated metals. As a consequence, it has been suggested that the more subtle aspects of metal toxicity in aquatic organisms can be more accurately estimated by examining the distribution of metals among the various intracellular ligand pools (Bayne *et al.*, 1980). However, to understand the ecological significance of these data we must also be able to relate this information on metal metabolism to effects on the organism, the population, and the community (Sanders *et al.*, 1983).

In a previous study on larvae of the crab *Rhithropanopeus harrisii* (Sanders *et al.*, 1983) we used metal-chelate buffer systems to control Cu speciation since the biological availability of Cu is related to the concentration of the free cupric ion, $[Cu^{2+}]$, rather than to the total or chelated Cu concentration (Sunda and Guillard, 1976; Anderson and Morel, 1978; Jackson and Morgan, 1978; Zamuda and Sunda, 1982). These buffers enabled us to expose larvae throughout zoeal development to a range of calculated $[Cu^{2+}]$ while the concentrations of potentially competitive metals were kept constant. In that study we found that most cytosolic Cu was associated with metallothionein, a cysteine-rich metal-binding protein whose synthesis is induced by metals (Hildebrand *et al.*, 1979). This ubiquitous protein has been implicated in metal uptake, metabolism, and detoxication in vertebrates and invertebrates (Richards and Cousins, 1976; Brown *et al.*, 1977; Li *et al.*, 1980; Roesijadi, 1980; Jenkins *et al.*, 1982). We found that the concentration of Cu-thionein in crab larvae was related

Received 16 April 1984; accepted 18 September 1984.

to $[Cu^{2+}]$ in sea water, and an increase in Cu-thionein accumulation correlated with inhibition of larval growth.

In this study we have followed changes in high (HMW) and very low molecular weight (VLMW) Cu binding ligands as well as Cu-thionein. Also, we examined these ligand pools over a wider range of $[Cu^{2+}]$. We report relationships between $[Cu^{2+}]$, Cu-thionein, and larval growth which are similar to our previous findings. In addition, we found that the Cu concentration within the VLMW pool could also be related to the $[Cu^{2+}]$ in sea water, and that increases of Cu in this pool correlated with inhibition of larval growth. In contrast to a current hypothesis (Brown, 1977), we found no correlation between Cu associated with HMW ligands and either external $[Cu^{2+}]$ or inhibition of larval growth. Finally, we found that increases of Cu in the MT and VLMW pools, and inhibition of larval growth, occurred at $[Cu^{2+}]$ just beyond the estimated ambient range in the estuary where the crabs were collected.

MATERIALS AND METHODS

Gravid females of the mud crab *Rhithropanopeus harrisii* were collected in the Newport River estuary near Beaufort, North Carolina. They were kept in tanks of running sea water and transferred to culture bowls when their eggs were ready to hatch. Newly hatched larvae were exposed to a range of $[Cu^{2+}]$ with metal-nitrotriacetic acid (NTA) buffer systems in sea water (Sunda *et al.*, 1978). The larvae were cultured in the laboratory until they molted to the megalopa stage as described by Costlow *et al.* (1966). Buffered sea water was changed daily. There were 10 larvae per culture bowl and five replicates for each treatment (Sanders *et al.*, 1983).

The metal-NTA buffer system was made up in 35 parts per thousand sea water which had previously been diluted with distilled water to 20 parts per thousand and contained 10^{-4} M NTA, 4.6×10^{-8} M ZnCl₂, 2.4×10^{-8} M CoCl₂, 1.0×10^{-8} M MnCl₂, 1.0×10^{-7} FeCl₃, and 1.0×10^{-9} M Na₂MoO₄. NaOH was added at 4 $\times 10^{-4}$ M to adjust the pH to 8.0 ± 0.1 . Copper was added as Cu-NTA to achieve concentrations of 1.8×10^{-8} M to 5.7×10^{-4} M. Free ion concentrations for Cu²⁺ were computed from metal ion-NTA equilibria (Sunda *et al.*, 1978). These concentrations were based on added metal and ranged from $10^{-12.4}$ M to $10^{-7.9}$ M.

After they molted to the megalopa stage the larvae were sampled, rinsed with distilled water, and lyophilized immediately. Survival and length of time to megalopa were determined as described by Costlow *et al.*, 1966. The freeze-dried megalopa were weighed to the nearest 0.1 μ g (Sanders *et al.*, 1984).

Replicate samples were pooled, rehydrated, and homogenized in 0.02 *M* tris-HCl (pH 7.4) with an acid-washed Teflon pestle tissue grinder. The homogenate was centrifuged at $100,000 \times g$, and the resulting supernatant was filtered by centrifugation through a 0.2 μ m nylon filter. An aliquot (100 μ l) of this filtered cytosol was chromatographed on a Toyo Soda steric exclusion (SEC) HPLC column (TSK SW 3000; Sanders *et al.*, 1983). Copper concentrations in the eluant fractions were determined by graphite furnace atomic absorption spectrophotometry (Sanders *et al.*, 1983). Chromatographic data were analyzed by correlation analysis, linear and orthogonal polynomial regression analysis, and analysis of variance (Sokal and Rohlf, 1981). Our confidence limits were set at 0.05.

RESULTS

Cytosolic Cu was associated with three major pools of ligands (Fig. 1): (1) high molecular weight ligands (HMW; $\geq 20,000$ daltons), the majority of which elute with or just behind the void volume; (2) a metallothionein peak (MT) which has an



FIGURE 1. Cytosolic distribution of copper in larvae of the crab *Rhithropanopeus harrisii* exposed to free cupric ion concentrations in sea water. HMW, MT, and VLMW represent high molecular weight, metallothionein, and very low molecular weight pools, respectively. The bars represent fractions which were combined for each pool.

apparent molecular weight of 10,000 daltons, and (3) very low molecular weight ligands (VLMW; <5000 daltons) which included molecules too small to be resolved by the column.

There was a biphasic relationship between $[Cu^{2+}]$ in sea water and Cu-thionein and LMW Cu. At $[Cu^{2+}]$ from $10^{-12.68}$ to $10^{-10.6}$ *M* copper metabolism appeared independent of external $[Cu^{2+}]$ (Fig. 2a). At these concentrations there was no significant correlation between external $[Cu^{2+}]$ and Cu-thionein. Copper associated with the VLMW pool was not detectable at environmental concentrations below $10^{-11.0}$ *M*. At concentrations greater than $10^{-10.6}$ *M*, however, Cu in the metallothionein and the VLMW pools correlated significantly with increases in $[Cu^{2+}]$ in sea water (F = 220; d.f. = 1, 2; F = 365; d.f. = 1, 2). This copper accumulation was reflected in total cytosolic Cu (F = 363; d.f. = 1, 2).

The relationship between Cu associated with the HMW ligands and $[Cu^{2+}]$ in sea water contrasted with those of the other pools (Fig. 2b). Although Cu associated with these ligands was significantly above background (F = 1096; d.f. = 8, 36), it did not correlate with $[Cu^{2+}]$ in sea water and no accumulation phase was observed (Fig. 2b).

We examined the relationships between $[Cu^{2+}]$ in sea water, survival, duration of the zoeal stages, and megalopa weight to determine the physiological affect of changes in intracellular Cu distribution. Free cupric ion concentrations did not correlate with survival or the duration of zoeal development at concentrations less than $10^{-8.9}$ M. At the two highest concentrations tested $(10^{-7.9} M \text{ and } 10^{-8.4} M)$, all larvae died within three days of exposure. Megalopa weight did correlate with $[Cu^{2+}]$ with significant linear (F = 53; d.f. = 1, 36) and quadratic components (Fig. 2c; F = 50; d.f. = 1, 36). Megalopa weight was greatest for larvae exposed to $10^{-11.0} M [Cu^{2+}]$ and decreased rapidly at higher concentrations.

Copper in the MT and VLMW ligand pools correlated with megalopa weight (Fig. 3; $r^2 = 0.676$; $r^2 = 0.826$). Megalopa weight increased with increases in Cu-



FIGURE 2. (a) Cytosolic distribution of copper expressed in micromoles per kilogram (wet weight) of tissue in *R. harrisii* megalopa exposed to a range of values of $p[Cu^{2+}]$ (-log of free cupric on concentration) throughout the duration of larval development. Total, MT, and LMW represent total cytosolic copper, metallothionein, and very low molecular weight ligand pools, respectively. (b) As described above for high molecular weight (HMW) ligand pool. (c) Dry weights of megalopa described above. The vertical lines are one standard error of the mean for five replicates.

thionein until a concentration of 72.6 μ m Cu kg⁻¹ wet weight (10^{-11.0} *M* exposure), and decreased rapidly at higher concentrations. Although Cu was undetectable in the VLMW pool at the three lowest [Cu²⁺], it accumulated at higher concentrations. Megalopa weight decreased rapidly with increases of Cu in this pool. In contrast to the MT and VLMW ligand pools, Cu in the HMW pool did not correlate with weight over the entire range of exposures.



FIGURE 3. Vector diagrams of megalopa dry weight and copper associated with the total cytosol and each cytosolic ligand pool (expressed as micromoles per kilogram dry weight of tissue) in *R. harrisii* megalopa. Total, HMW, MT, and VLMW represent total cytosolic copper, high molecular weight, metallothionein, and very low molecular weight ligands pools, respectively. The arrows delineate the $p[Cu^{2+}]$ (-log free cupric ion concentration) to which the larvae were exposed throughout development from the lowest, to the highest exposures ($p[Cu^{2+}]$ from 12.4 to 8.9).

DISCUSSION

Cytosolic copper accumulation and distribution

The cytosolic distributions of Cu among the HMW, MT, and VLMW ligand pools were similar to those previously reported for *R. harrisii* (Fig. 1; Sanders *et al.*, 1983) and for other crab species (Olafson *et al.*, 1979b; Overnell, 1982). These three metal binding pools have also been reported in other marine invertebrates (Coombs, 1974; Howard and Nickless, 1977, 1978; Jenkins *et al.*, 1982; Frazier and George, 1983).

In our fractionation procedure the HMW pool contained a heterogeneous group of Cu ligands. Most of these ligands are soluble proteins. A number of these proteins, including Cu metalloenzymes and the respiratory pigment hemocyanin, have specific The chemical characterization of metallothioneins in crabs includes amino acid analysis in two species (Olafson *et al.*, 1979a; Overnell, 1982) and amino acid sequencing for one species (Lerch *et al.*, 1981, 1982). Characterization of the MT pool of *R. harrisii* was described in our previous study (Sanders *et al.*, 1983). As with other crab (Olafson *et al.*, 1979a; Overnell and Trewhella, 1979) and mammalian (Kagi and Nordberg, 1979) metallothioneins, it resolves as two major metal binding ligands when fractionated by ion exchange chromatography.

The chemical identity and origin of the VLMW ligand pool is not known but it appears to be made up of a heterogeneous group of Cu complexes. Amino acids may represent an important component of this pool, but these data have yet to be confirmed (Coombs, 1974; Howard and Nickless, 1977, 1978). Frazier and George (1983) have suggested that *in vivo* these VLMW ligands may be associated with granules which become solubilized during homogenization and extraction. Granules containing high concentrations of Cu are present in the hepatopancreas of another crustacean (Weiser and Klima, 1969) but it is not clear if this is the source of the VLMW ligands in crab larvae.

Copper regulation and metabolism

The relationship between $[Cu^{2+}]$ in sea water and cytosolic Cu fell into two discrete patterns (Fig. 2). This biphasic response can be related to the ambient $[Cu^{2+}] (\leq 10^{-10.4} M)$ estimated by Sunda and Gillespie (1979) for estuarine waters where the crabs were collected. The first phase occurred at a range of $[Cu^{2+}]$ which is equal to or less than the estimated ambient concentration $(10^{-12.4} M \text{ to } 10^{-10.6} M)$. In this range of 1.8 orders of magnitude, Cu in the HMW, MT, and VLMW ligand pools were independent of $[Cu^{2+}]$. These data support our previous observations (Sanders *et al.*, 1983) and indicate that Cu accumulation and distribution among the three ligand pools is precisely regulated in crab larvae at exposures at or below ambient $[Cu^{2+}]$.

When larvae are exposed to $[Cu^{2+}]$ which are greater than the ambient concentration $(10^{-10.2} M \text{ to } 10^{-8.9} M)$, a second phase, one of accumulation, was observed (Fig. 2a). In this phase total cytosolic Cu increased with increases in $[Cu^{2+}]$ in sea water. This accumulation was the result of increases in both Cu-thionein and Cu in the VLMW ligand pool. Accumulation of total cytosolic Cu and Cu-thionein in *R. harrisii* larvae exposed to greater than ambient $[Cu^{2+}]$ has been reported previously (Sanders *et al.*, 1983). In that study cytosolic Cu accumulation and Cu-thionein concentration approached apparent saturation at $[Cu^{2+}]$ just below those which resulted in larval death. This saturation, however, was based on a minimum of data points. In contrast, the data presented here showed no evidence of saturation and the concentrations of total Cu and Cu-thionein attained prior to larval mortality were several fold higher than in the previous experiment (Fig. 3a).

In marked contrast to the other ligand pools, Cu in the HMW ligand pool did not correlate with $[Cu^{2+}]$ in sea water (Fig. 2b). The lack of Cu accumulation in this pool suggests that crab larvae can regulate subcellular Cu distribution over the 3.5 orders of magnitude range of $[Cu^{2+}]$ used in this experiment. Also, the simultaneous increase of Cu in the MT and VLMW ligand pools suggests that they may function as Cu reservoirs which limit nonspecific binding to the HMW ligands.

Free cupric ion concentration and larval growth

In order to understand the ecological significance of Cu accumulation and cytosolic distribution, we examined three parameters: survival, duration of zoeal development,

and megalopa weight. However, only changes in megalopa weight correlated with $[Cu^{2+}]$ (Fig. 2c). Maximum megalopa weight occurred within the range of the estimated ambient $[Cu^{2+}]$ and decreased rapidly at concentrations beyond this range. This pattern of a hormetic overshoot in weight followed by a rapid decrease is an inherent characteristic of stress etiology which has been reported previously in *R. harrisii* larvae (Laughlin *et al.*, 1981; Sanders *et al.*, 1983; 1984). These growth data support our previous observation that fluctuations in ambient $[Cu^{2+}]$ in sea water could limit crab larval growth (Sanders *et al.*, 1983). Cupric ion concentrations near the ambient range also limit growth and nutrient uptake in phytoplankton and bacteria (Jackson and Morgan, 1978; Fitzwater *et al.*, 1982; Sunda and Ferguson, 1983; Jenkins *et al.*, 1984). Thus, natural variations in $[Cu^{2+}]$ could regulate productivity in a wide range of estuarine and marine species through the inhibition of important metabolic processes. The effects of this inhibition could, in turn, modify population dynamics and community composition (Sunda *et al.*, 1981).

Copper accumulation and larval growth

It has been suggested that as long as metals can be sequestered in the MT pool the organism is protected against the toxicity of these metals (Brown *et al.*, 1977, 1983; Jenkins *et al.*, 1982; Nolan and Duke, 1983). However, our data indicate that when larvae were exposed to $[Cu^{2+}]$ beyond the ambient range, the accumulation of Cu in the MT pool was correlated with growth inhibition (Fig. 3).

Although the nonspecific binding of metals to HMW ligands is proposed to be a major mechanism of metal toxicity (Winge *et al.*, 1973; Brown, 1977; Pruell and Engelhardt, 1980), our data show no correlation between Cu in the HMW ligand pool and toxic effects on growth (Fig. 3). Several explanations may account for this apparent discrepency: (1) the techniques used in this study can only detect changes in total Cu and, therefore, do not necessarily distinguish changes in the level of nonspecific binding; (2) the mechanism of metal toxicity may involve the nonspecific binding of Cu to membrane-bound metalloenzymes or macromolecules in other subcellular compartments; and (3) finally Cu uptake and accumulation may modify the uptake and metabolism of other essential metals (*e.g.*, Zn) which may in turn affect larval growth (Bremner and Campbell, 1980; Sunda *et al.*, 1981; Cousins, 1983).

Of the three cytosolic ligand pools, it was the accumulation of Cu in the VLMW pool which was most indicative of toxic effects on the organism. The detection of Cu in this pool was concomitant with perturbations of larval growth. As the Cu concentration in the VLMW pool increased, larval growth rapidly decreased. Cadmium accumulation has been reported in the VLMW pool of the oyster (*Ostrea edulis*) during chronic Cd exposure (Frazier and George, 1983). Although the authors suggest that sequestering cadmium in this pool may protect the oyster from Cd toxicity, they provide no information on the relationship between this accumulation and sublethal effects at the organismic level.

These data emphasize the importance of understanding the interactions among all cytosolic ligand pools. They also underscore the need to correlate these interactions with sensitive indicators of stress (such as growth) before the toxic effects of metals on an organism can be estimated from data on subcellular metal distributions. Finally, these data indicate that changes in intracellular metal metabolism and growth in crab larvae occur at the upper range of the ambient $[Cu^{2+}]$ in the estuary, suggesting that $[Cu^{2+}]$ in sea water may limit larval growth.

ACKNOWLEDGMENTS

We thank W. G. Sunda for his help in developing the metal buffers, J. D. Costlow for his help with larval culture techniques, C. Norwood for statistical assistance, and P. C. Denny, P. A. Denny, and P. Oshida for critical review of earlier drafts of this manuscript. We also thank T. Marshall and J. F. Alfafara for technical assistance throughout this study. This study was supported by grants from the Ecological Division of the Office of Health and Environmental Research, Department of Energy, to Duke University, and to California State University, Long Beach. This publication is contribution JS-84-002 of the Molecular Ecology Institute, California State University, Long Beach.

LITERATURE CITED

- ANDERSON, D. M., AND F. M. MOREL. 1978. Copper sensitivity of Gonyaulax tamarensis. Limnol. Oceanogr. 23: 283–295.
- BAYNE, B. L., D. A. BROWN, F. L. HARRISON, AND P. P. YEVICH. 1980. Mussel Health. Pp. 163–235 in The International Mussel Watch, Report on a Workshop at Barcelona, Spain, December 1978, E. Goldberg, ed. National Academy of Science, Washington, DC.
- BREMNER, I., AND J. K. CAMPBELL. 1980. The influence of dietary copper intake on the toxicity of cadmium. NYAS 355: 319–332.
- BROUWER, M., C. BONAVENTURA, AND J. BONAVENTURA. 1983. Metal ion interactions with Limulus polyphemus and Callinectes sapidus hemocyanins: Stoichiometry and structural and functional consequences of calcium (II), cadmium (II), zinc (II), and mercury (II) binding. Biochemistry 22: 4713-4723.
- BROWN, D. A. 1977. Increases of Cd and the Cd:Zn ratio in the high molecular weight protein pool from apparently normal liver of tumor-bearing flounders (*Parophrys vetulus*). Mar. Biol. 44: 203–209.
- BROWN, D. A., C. A. BAWDEN, K. W. CHATEL, AND T. R. PARSONS. 1977. The wildlife community of Iona Island jetty, Vancouver, B.C., and heavy-metal pollution effects. *Environ. Conserv.* 4: 213– 216.
- BROWN, D. A., R. W. GOSSETT, P. HERSHELMAN, H. A. SCHAEFER, K. D. JENKINS, AND E. M. PERKINS. 1983. Bioaccumulation and detoxification of contaminants in marine organisms from southern California coastal waters. Pp. 171–190 in *Waste Disposal in the Oceans*, D. F. Soule and D. Walsh, eds. Westview Press, Boulder, Colorado.
- COOMBS, T. L. 1974. The nature of zinc and copper complexes in the oyster. Ostrea edulis. Mar. Biol. 28: 1-10.
- COSTLOW, J. D., AND C. G. BOOKOUT. 1971. The effect of cyclic temperatures on larval development in the mudcrab *Rhithropanopeus harrisii*. Pp. 211–220 in *Fourth European Marine Biology Symposium*, D. J. Crisp, ed. Cambridge University Press, London.
- COSTLOW, J. D., C. G. BOOKOUT, AND R. J. MONROE. 1966. Studies on the larval development of the crab *Rhithropanopeus harrisii* (Gould). I. The effects of salinity and temperature on larval development. *Physiol. Zool.* **39**: 81–100.
- COUSINS, R. J. 1983. Metallothionein—aspects related to copper and zinc metabolism. J. Inherited Metab. Dis. 6 Suppl. 1: 15-21.
- FITZWATER, S. E., G. A. KNAUER, AND J. H. MARTIN. 1982. Metal contamination and its effect on primary production measurements. *Limnol. Oceanogr.* 27: 544–551.
- FRAZIER, J. M., AND S. G. GEORGE. 1983. Cadmium kinetics in oysters—a comparative study of *Crassostrea* gigas and Ostrea edulis. Mar. Biol. **4**: 55–62.
- HILDEBRAND, C. E., R. A. TOBEY, E. W. CAMPBELL, AND M. D. ENGER. 1979. A cadmium-resistant variant of the Chinese hamster (CHO) cell with increased metallothionein induction capacity. *Exp. Cell Res.* **124**: 237–246.
- HOWARD, A. G., AND G. NICKLESS. 1977. Heavy metal complexation in polluted molluscs. II. Oyster (Ostrea edulis and Crassostrea gigas) Chem. Biol. Interact. 17: 257-263.
- HOWARD, A. G., AND G. NICKLESS. 1978. Heavy metal complexation in polluted molluscs. III. Periwinkles (Littorina littorea), cockles (Cardium edule) and scallops (Chlamys opercularis). Chem. Biol. Interact. 23: 227-231.
- JACKSON, G. A., AND J. J. MORGAN. 1978. Trace metal-chelator interactions and phytoplankton growth in seawater media: Theoretical analysis and comparison with reported observations. *Limnol. Oceanogr.* 23: 268–282.
- JENKINS, K. D., D. A. BROWN, G. P. HERSHELMAN, AND W. C. MEYER. 1982. Contaminant in white croakers Genyonemus lineatus (Ayers 1855) from the southern California bight: I Trace metal detoxification/toxification. Pp. 177–196 in Physiological Mechanisms of Marine Pollutant Toxicity, W. B. Vernberg, A. Calabrese, F. P. Thurberg, F. J. Vernberg, eds. Academic Press, New York.
- JENKINS, K. D., D. A. BROWN, P. S. OSHIDA, AND E. M. PERKINS. 1982. Cytosolic metal distribution as an indicator of toxicity in sea urchins from the southern California bight. *Mar. Pollut. Bull.* 13: 413-421.

- JENKINS, K. D., B. M. SANDERS, AND W. G. SUNDA. 1984. Metal regulation and toxicity in aquatic organisms. Pp. 277-288 in *Mechanisms of Drug and Toxin Action*, T. Singer, T. Mansour, R. Ondarza, eds. Academic Press, New York.
- KAGI, J. H. R., AND M. NORDBERG. 1979. *Metallothionein*. Birkhauser Verlag, Basel, Boston, Stuttgart. 378 pp.
- LAUGHLIN, R. B., JR., J. NG, AND H. E. GUARD. 1981. Hormesis: A response to low environmental concentrations of petroleum hydrocarbons. *Science* 211: 705-707.
- LERCH, K., D. AMMER, AND R. W. OLAFSON. 1981. Amino acid sequence of crab metallothionein. FEBS Lett. 126: 165-168.
- LERCH, K., D. AMMER, AND R. W. OLAFSON. 1982. Crab metallothionein, primary structures of metallothioneins 1 and 2. J. Biol. Chem. 257: 2420-2423.
- LI, T. Y., A. J. KRAKER, C. F. SHAW III, AND D. H. PETERING. 1980. Ligand substitution reactions of
- metallothioneins with EDTA and apo-carbonic anhydrase. *Proc. Natl. Acad. Sci.* 77: 6334-6338. NOLAN, C. V., AND E. J. DUKE. 1983. Cadmium accumulation and toxicity in *Mytilus edulis:* involvement of metallothioneins and heavy-molecular weight protein. *Aguat. Toxicol.* 4: 153-163.
- OLAFSON, R. W., R. G. SIM, AND K. G. BOTO. 1979a. Isolation and chemical characterization of the heavy metal-binding protein metallothionein from marine invertebrates. *Comp. Biochem. Physiol.* 62B: 407–416.
- OLAFSON, R. W., A. KEARNS, AND R. G. SIM. 1979b. Heavy metal induction of metallothionein synthesis in the hepatopancreas of the crab *Scylla serrata*. *Comp. Biochem. Physiol.* **62B**: 417–423.
- OVERNELL, J. 1982. Copper metabolism in crabs and metallothionein: *in vivo* effects of copper II on soluble hepatopancreas metal binding components in the crab *Cancer pagurus* containing varying amounts of cadmium. *Comp. Biochem. Physiol.* **73B:** 555–564.
- OVERNELL, J., AND E. TREWHELLA. 1979. Evidence for the natural occurrance of (cadmium, copper) metallothionein in the crab *Cancer pagurus*. Comp. Biochem. Physiol. 64C: 69-76.
- PRUELL, R. J., AND F. E. ENGELHARDT. 1980. Liver cadmium uptake, catalase inhibition and cadmium thionein production in the killifish (*Fundulus heteroclitus*) induced by experimental cadmium exposure. *Mar. Envir. Res.* 3: 101–111.
- RICHARDS, M. P., AND R. J. COUSINS. 1976. Influence of parenteral zinc and actinomysin D on tissue zinc uptake and the synthesis of a zinc-binding protein. *Bioinorg. Chem.* 4: 215-224.
- ROESIJADI, G. 1980. Influence of copper on the clam *Protothaca staminea*: effects on gills and occurrence of copper-binding proteins. *Biol. Bull.* **158**: 233–247.
- SANDERS, B. M., K. D. JENKINS, W. G. SUNDA, AND J. D. COSTLOW. 1983. Free cupric ion activity in seawater: effects on metallothionein and growth in crab larvae. *Science* 222: 53–55.
- SANDERS, B. M., R. B. LAUGHLIN, JR., AND J. D. COSTLOW. 1984. Growth regulation in larvae of the mud crab, *Rhithropanopeus harrisii*. In *Crustacean Issues, Vol. 2: Crustacean Growth, A. M.* Wenner, ed. A. A. Balkema, Rotterdam. (In press.)
- SOKAL, R. R., AND F. J. ROHLF. 1981. Biometry. W. H. Freeman and Company, San Francisco. 859 pp.
- SUNDA, W. G., AND R. L. FERGUSON. 1983. Sensitivity of natural bacterial communities to additions of copper and to cupric ion activity: a bioassay of copper complexation in seawater. P. 871 in *Trace Metals in Sea Water*, C. Wong, E. Boyle, K. W. Bruland, J. D. Burton, and E. D. Goldberg, eds. Plenum Publishing Corp., New York.
- SUNDA, W. G., AND P. A. GILLESPIE. 1979. The response of a marine bacterium to cupric ion and its use to estimate cupric ion activity in seawater. J. Mar. Res. 37: 761-777.
- SUNDA, W. G., AND R. R. GUILLARD. 1976. The relationship between cupric ion activity and the toxicity of copper to phytoplankton. J. Mar. Res. 34: 511-529.
- SUNDA, W. G., R. T. BARBER, AND S. A. HUNTSMAN. 1981. Phytoplankton growth in nutrient rich seawater: importance of copper-manganese cellular interactions. J. Mar. Res. 39: 567–586.
- SUNDA, W. G., D. W. ENGEL, AND R. M. THUOTTE. 1978. Effect of chemical speciation on toxicity of cadmium to grass shrimp, *Palaemonetes pugio*: Importance of free cadmium ion. *Environ. Sci. Technol.* 12: 409-413.
- WEISER, W., AND J. KLIMA. 1969. Compartmentalization of copper in the hepatopancreas of isopods. *Mikroskopie* 24: 1-9.
- WINGE, D., J. KRASNO, AND A. V. COLUCCI. 1973. Cadmium accumulation in rat liver: correlation between bound metal and pathology. Pp. 500–501 in *Trace Element Metabolism in Animals*, Vol. 2, W. G. Hoetkstra, J. W. Suttie, H. E. Ganther, and W. Mertz, eds. University Press, Baltimore.
- ZAMUDA, C. D., AND W. G. SUNDA. 1982. Bioavailability of dissolved copper to the American oyster Crassostrea virginica. I. Importance of chemical speciation. Mar. Biol. 66: 77-82.



Biodiversity Heritage Library

Sanders, Brenda M and Jenkins, Kenneth D. 1984. "RELATIONSHIPS BETWEEN FREE CUPRIC ION CONCENTRATIONS IN SEA WATER AND COPPER METABOLISM AND GROWTH IN CRAB LARVAE." *The Biological bulletin* 167, 704–712. <u>https://doi.org/10.2307/1541421</u>.

View This Item Online: https://doi.org/10.2307/1541421 Permalink: https://www.biodiversitylibrary.org/partpdf/40402

Holding Institution MBLWHOI Library

Sponsored by MBLWHOI Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: University of Chicago License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.