Case 3386.

Pseudocoenia d'Orbigny, 1850 (Coelenterata, Scleractinia): proposed conservation of usage by the designation of a lectotype for the type species

H. Löser

Universidad Nacional Autónoma de México, Instituto de Geología, Estación Regional del Noroeste, Apartado Postal 1039, 83000 Hermosillo, Sonora, México (e-mail: loeser@paleotax.de)

Abstract. The purpose of this application, under Article 74.1 of the Code, is to conserve the name *Pseudocoenia* d'Orbigny, 1850 in its accustomed usage by designating a new lectotype for its type species *Pseudocoenia bernardina* d'Orbigny, 1850. The present lectotype of the type species contradicts the original description of the type species as well as the concept of the genus as indicated by its author and as currently used.

Keywords. Nomenclature; taxonomy; Coelenterata; Scleractinia; *Pseudocoenia*; *Pseudocoenia* bernardina; Jurassic-Cretaceous; corals.

- 1. The genus *Pseudocoenia* (Coelenterata, Scleractinia, STYLINIDAE) was established by d'Orbigny (1850, p. 33). His original diagnosis of the genus is short but to the point for people who know Mesozoic corals: 'Ce sont des *Cryptocoenia* à huit systèmes, au lieu de six' [It is a *Cryptocoenia* with eight systems instead of six]. *Cryptocoenia* was established earlier by d'Orbigny (1849, p. 7) as a '*Tremocoenia*, sans columelle styliforme' [a *Tremocoenia* without any styliform columella].
- 2. The genus *Pseudocoenia* was characterised as a *Cryptocoenia* with eight instead of six septal systems, i.e. the number of septa in one generation (or cycle) is eight or a multiple (16, 32) of eight. When the genus was established, d'Orbigny (1850, p. 34) listed seven species: *Pseudocoenia suboctonis*, *Pseudocoenia bernardina*, *Pseudocoenia ramosa*, *Pseudocoenia digitata*, *Pseudocoenia octonis*, *Pseudocoenia elegans* and *Pseudocoenia ramosa*. In some cases the presence of eight septal systems is repeated in the species diagnosis. Currently a total of 23 Jurassic (Lathuilière, 1989) and one Cretaceous species (Löser, 2000) are assigned to the genus. The genus is widely distributed in the Late Jurassic and is rare in the Early Cretaceous. All seven originally assigned species have equal claim to be designated as the type species, since there is no Article in the Code which gives priority to the first species in the list.
- 3. The type species of *Pseudocoenia* was designated by Wells (1936, p. 128) according to Article 69.1 of the Code (Type species by subsequent designation): *Pseudocoenia bernardina* d'Orbigny, 1850 (p. 34). The type locality according to d'Orbigny is 'France, Landeyron'.
- 4. Pseudocoenia bernardina d'Orbigny, 1850 is characterised as 'Espèce voisine de l'octonis par ses huit chambres aux cellules, mais dont celles-ci sont d'un tiers plus

petites' [species close to octonis by its eight chambers in the cell, but these are one third smaller].

- 5. Four syntypes of *Pseudocoenia bernardina* d'Orbigny, 1850 exist in the Coll. d'Orbigny, Muséum National d'Histoire Naturelle (MNHN) Paris:
 - (1) MNHN (Paris), typothèque, R9199 (ex Coll. d'Orbigny 4472); 'Pseudocoenia bernardiana d'Orbigny' Corallien, Landeyron (Ain); figured in Cottreau, 1931, pl. 61, fig. 6; small specimen with one polished surface. Septa in a decameral system. Styliform columella.
 - (2) MNHN (Paris), Coll. d'Orbigny, 4472 no label; Landeyron (Ain); poorly preserved small specimen without polished surface which possibly belongs to the same species as R9199.
 - (3) MNHN (Paris), Coll. d'Orbigny, 4472a; no label; Tonnerre (Yonne); a mould of a plocoid coral.
 - (4) MNHN (Paris), Coll. d'Orbigny, 4472b; no label; Châtel-Censoir (Yonne); a large specimen with a thin section. Septa in an octomeral system. No columella.
- 6. From these syntypes, Wells (1936, p. 128) designated the specimen of number 4472 as lectotype. He excluded the specimens numbered 4472a and 4472b by writing 'Type No. 4472 [non 4472a, b]'. Wells was probably influenced by (1) this specimen was depicted by Cottreau (1931), who first illustrated - at least in part - the fossil corals of the d'Orbigny Collection, (2) the identical type locality to the locality indicated by d'Orbigny, and (3) the opinion expressed by other authors, like Milne-Edwards & Haime (1851, p. 78), that Pseudocoenia was a junior synonym of Stylina Lamarck, 1816. The last has to be seen in a historical context. Milne-Edwards and Haime believed that all 'Stylina without columella' were only poorly preserved and that the absence of the columella was caused by an 'accident dependent on the process of fossilisation'. Milne-Edwards and Haime saw no reason why corals comparable to Stylina but without columella should be separated. Milne-Edwards (1857, p. 235) mentioned the eight systems in Pseudocoenia (not ten, as in the lectotype designated by Wells), but did not see any reason why Pseudocoenia or analogous genera (such as the above-mentioned Cryptocoenia, among other forms without columella) should be regarded as being different from Stylina.
- 7. Wells's designation contradicts (1) d'Orbigny's diagnosis of the species *Pseudocoenia bernardina* d'Orbigny, 1850, (2) the definition of the genus *Pseudocoenia*, and (3) most of the originally included species, as far as type material is available. The genus *Pseudocoenia* in its accustomed usage is understood as belonging to stylinids without columella, and with short and/or rudimentary septa in symmetry of eight. The current lectotype of *Pseudocoenia bernardina* d'Orbigny, 1850 belongs to a stylinid genus (i.e. *Stylina*) with ten septal systems and a styliform columella. When Wells (1936) designated the lectotype, he did not comply with d'Orbigny's intention (1850). Instead of admitting a coral comparable to *Cryptocoenia* but with an 8-part septal symmetry, Wells obviously sought to synonymize genera, including *Pseudocoenia*. *Pseudocoenia* is still in use in recent literature in its original understanding; its synonymy with *Stylina* is widely ignored (Beauvais, 1964; Eliášová, 1981; Errenst, 1990; Helm et al., 2003; Helm, 2005; Kuzmicheva, 2002; Liao & Xia, 1994; Pandey & Fürsich, 2003; Roniewicz, 1966, 1976; Turnšek, 1972).
- 8. Alloiteau (1948, p. 704) revised the genus *Pseudocoenia*. The proposed designation of *Pseudocoenia suboctonis* d'Orbigny, 1850 as the type species is invalid being

preceded by Well's designation. Alloiteau based his investigations on type material of *Pseudocoenia suboctonis*, emending the characteristics of the genus. He was followed by Beauvais (1964). Later Roniewicz (1966, 1976) and Eliášová (1981) modified the generic concept and also accepted corals with symmetry other than eight. In the past fifteen years this has caused some confusion, mainly when dealing with Cretaceous corals, where one and the same species has been successively assigned to *Adelocoenia* d'Orbigny, 1849, *Convexastrea* d'Orbigny, 1849, *Cryptocoenia* d'Orbigny, 1849, *Cyathophora* Michelin, 1842, *Orbignycaenia* Alloiteau, 1948 or *Pseudocoenia* d'Orbigny, 1850.

- 9. In order to conserve the name *Pseudocoenia* d'Orbigny, 1850 in its accustomed use, it is proposed that all previous type fixations for its type species *Pseudocoenia bernardina* d'Orbigny, 1850 be set aside and that a lectotype consistent with current usage be designated; a suitable specimen would be specimen 4472b, Coll. d'Orbigny, MNHN (Paris).
- 10. The International Commission on Zoological Nomenclature is accordingly asked:
 - (1) to use its plenary power to set aside all previous type fixations for the nominal species *Pseudocoenia bernardina* d'Orbigny, 1850 and to designate specimen 4472b, Coll. d'Orbigny, MNHN (Paris) as the lectotype;
 - (2) to place on the Official List of Specific Names in Zoology the name *bernardina* d'Orbigny, 1850, as published in the binomen *Pseudocoenia bernardina* and as defined by the lectotype 4472b, Coll. d'Orbigny, MNHN (Paris) designated in (1) above.

Acknowledgements

The proposal was submitted to various specialists on Mesozoic corals; minor corrections were provided by B. Lathuilière (Nancy) and D.K. Pandey (Jaipur).

References

- **Alloiteau, J.** 1948. Polypiers des couches albiennes à grandes trigonies de Padern (Aude). Bulletin de la Société Géologique de France. Sér. 5, **18**: 699–738.
- **Beauvais, L.** 1964. Étude stratigraphique et paléontologique des formations à madréporaires du Jurassique supérieur du Jura et de l'Est du Bassin de Paris. *Mémoires de la Société Géologique de France. N.S.*, **43**: 1–287.
- Cottreau, J. 1931. Types du prodrome de paléontologie stratigraphique universelle (8). *Annales de Paléontologie*, **20**(1): 133–172.
- Eliášová, H. 1981. Sous-ordre Stylinina Alloiteau, 1952 (Hexacorallia) des calcaires de Stramberk (Tithonien, Tchécoslovaquie). Sbornik geologickych vied, Paleontologie, 24: 117–133.
- Errenst, C. 1990. Das korallenführende Kimmeridgium der nordwestlichen iberischen Ketten und angrenzender Gebiete (1). *Palaeontographica. Abt. A*, **214**(3/6): 121–207.
- **Helm, C.** 2005. Riffe und fazielle Entwicklung der florigemma-Bank (Korallenoolith, Oxfordium) im Süntel und östlichen Wesergebirge (NW-Deutschland). *Geologische Beiträge Hannover*, 7: 3–339.
- Helm, C., Reuter, M. & Schülke, I. 2003. Die Korallenfauna des Korallenooliths (Oxfordium, Oberjura, NW-Deutschland): Zusammensetzung, Stratigraphie und regionale Verbreitung. Paläontologische Zeitschrift, 77(1): 77–94.
- **Kuzmicheva**, **E.I.** 2002. Morfologiya skeleta, sistema i evolyutsiya skleraktinii [Skeletal morphology, systematics and evolution of Scleractinia]. *Trudy Paleontologicheskogo instituta*, **286**: 1–211.

Lathuilière, B. 1989. Répertoire objectif des coraux jurassiques. 76 pp. Presses universitaires, Nancy.

Liao, Wei-hua & Xia, Jin-bao. 1994. Mesozoic and Cenozoic scleractinian corals from Tibet. Palaeontologia Sinica (Zhongguo-gushengwu-zhi), 184: 1–252.

Löser, H. 2000. Répertoire of Species. Catalogue of Cretaceous Corals, vol. 1. 137 pp. CPress Verlag, Dresden.

Milne-Edwards, H. 1857. Histoire naturelle des coralliaires ou polypes proprement dits (1, 2). viii, 326 pp., 633 pp. Librairie encyclopédique de Roret, Paris.

Milne-Edwards, H. & Haime, J. 1851. A monograph of the British fossil corals (2): Corals from the oolitic formations. *Palaeontographical Society monographs*, 5: 74–146.

Orbigny, A. d'. 1849. Note sur les polypiers fossiles. 12 pp. Masson, Paris.

Orbigny, A. d'. 1850. Prodrôme de paléontologie stratigraphique universelle des animaux mollusques et rayonnés (2). 428 pp. Masson, Paris.

Pandey, D.K. & Fürsich, F.T. 2003. Jurassic corals of east-central Iran. Beringeria, 32: 3–138.
Roniewicz, E. 1966. Les madréporaires du Jurassique supérieur de la bordure des monts de Sainte-croix, Pologne. Acta Palaeontologica Polonica, 11(2): 157–264.

Roniewicz, E. 1976. Les scléractiniaires du Jurassique supérieur de la Dobrogea centrale Roumanie. *Palaeontologia Polonica*, **34**: 17–121.

Turnšek, D. 1972. Zgornjejurske korale iz juzne Slovenije. Razprave Slovenska akademija znanosti in umetnosti, (4)15(6): 147–265.

Wells, J.W. 1936. The nomenclature and type species of some genera of recent and fossil corals. *American Journal of Science*, (5)31(182): 97–134.

Acknowledgement of receipt of this application was published in BZN 63: 154.

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the Executive Secretary, I.C.Z.N., c/o Natural History Museum, Cromwell Road, London SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).



Löser, H. 2007. "Case 3386 Pseudocoenia d'Orbigny, 1850 (Coelenterata, Scleractinia): proposed conservation of usage by the designation of a lectotype for the type species." *The Bulletin of zoological nomenclature* 64, 79–82.

View This Item Online: https://www.biodiversitylibrary.org/item/107007

Permalink: https://www.biodiversitylibrary.org/partpdf/42795

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: International Commission on Zoological Nomenclature

License: http://creativecommons.org/licenses/by-nc-sa/3.0/

Rights: https://biodiversitylibrary.org/permissions

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.