


PROCEEDINGS
OF THE
BIOLOGICAL SOCIETY OF WASHINGTON

DRACHMASTER BULLISI NEW GENUS AND SPECIES
OF OPHIDIASTERIDAE (ECHINODERMATA:
ASTEROIDEA), WITH A KEY TO THE
CARIBBEAN SPECIES OF THE FAMILY

BY MAUREEN E. DOWNEY

Smithsonian Institution, Washington, D.C.




Two specimens of a small starfish collected by the Bureau of Commercial Fisheries vessel M/V *Oregon*, one off Trinidad and the other off Martinique, represent another genus of the large family Ophidiasteridae. Of the 23 recognized genera in this family, at least six are represented in the Caribbean and the Gulf of Mexico. Eight or more species of Ophidiasteridae have been reported from this same area; however, in the course of a study of a very large collection of asteroids from this region, only six of the species (in addition to the new one) have been found. This essentially tropical family is principally a shallow water one, only three species in the family occurring at depths below 200 fathoms. Indeed, most species are found in very shallow water and they are usually reef-dwelling or reef-associated. They are remarkably abundant in the Indo-Pacific region and especially in the vicinity of Torres Strait. H. L. Clark (1921) has very thoroughly reviewed the family and described several new species from Australia. Two new monotypic genera have been added to the family since that time, one from the Indo-Pacific and one from Florida.

Drachmaster new genus

Arms with internal plates connecting the outer plates in transverse series.

Etymology: Generic name—masculine, from the Greek, drachme, target, and aster, star.





FIGS. 1-4. 1, Cross-section of half a ray, showing arrangement of plates. 2, External view of transverse arrangement of plates. 3, Pedicellaria (erect). 4, Alveolus in which pedicellaria rests. Abbreviations: C = carinal plate; Ar = adradial plate; S = superomarginal; I = inferomarginal; A = first actinolateral; B = second actinolateral; Ad = adambulacral plates; Ip = internal plates.

***Drachmaster bullisi* new species**

Figures 1-6

Disc small, rather flat. Anal pore central, surrounded by a number of enlarged spiniform granules; madreporite irregularly triangular or elliptical, sunken below general disc surface, and covered with deep gyri. Arms long, cylindrical, not much tapered, and terminate bluntly. All plates covered with moderately thick skin closely beset with small squamiform or minutely spinous granules of uniform size (because of this skin covering, it was not possible to determine the exact arrangement of the disc plates, but it seems to be quite like that of *Ophidiaster*). Plates on arms smooth, slightly tumid, in 11 very regular longitudinal rows (exclusive of adambulacral plates), and also in regular transverse rows. Plates in transverse series connected internally by small tumid plates (the principal feature by which *Drachmaster* is distinguished from all other genera of Ophidiasteridae) (Fig. 1). Eight longitudinal rows of conspicuous papular areas, with from one to three papulae per area. Viewed externally, from top of arm to adambulacrals, plates in transverse

**Fig. 5****Fig. 6**

FIGS. 5-6. 5, *Drachmaster bullisi* (Holotype, USNM E11359). Aboral view. 6, Oral view, same specimen.

series are: (1) carinal, (2) adradial, (3) superomarginal, (4) inferomarginal, (5) first actinolateral, and (6) two second actinolaterals, side by side, each corresponding to an adambulacral plate (fig. 2).

Adambulacral armature of two short, blunt, somewhat flattened furrow spines, connected near their bases by skin, and a much longer,

flat, more acute outer spine. Small spines around oral margin of jaw plates similar to and continuous with furrow spines; behind these, on actinal surface of jaw plates, two pairs of large spines, like outer adambulacral spines; distalmost pair accompanied each by a small spinelet between furrow spines and larger outer spine. Internally, a large flange on upper edge of ambulacral plates overlaps upper edge of ambulacral plate proximal to it. Superambulacral plates present.

On smaller specimen, but one pedicellaria, on abactinal surface in one interradial area; on rays, several incipient pedicellariae between adambulacral and inferomarginals. Larger specimen abundantly provided with pedicellariae; one on upper edge of each superomarginal plate. Pedicellariae rare below inferomarginals, but occur frequently on abactinal surface. Valves of pedicellariae slender, elongate, minutely toothed, flattened laterally, and terminate in prominent hyaline hook which crosses hook of opposite valve when pedicellariae are erect (Fig. 3). Alveoli in which pedicellariae rest elongate-diamond-shaped, with raised lips (Fig. 4).

Ocular plates large, tumid; terminal portion of plate covered with large low rounded tubercles.

R/r—6/1. Holotype: R = 33 mm, r = 4.5 mm. Paratype: R = 14 mm, r = 4.5 mm.

The type material is in the U.S. National Museum.

Material examined: Holotype (USNM E11359), "Oregon" Station 5970, 11°13'N, 60°52'W, 34 fms., March 1966 (one arm missing, another regenerating). Paratype (USNM E11360), "Oregon" Station 5923, 15°36'N, 61°15'W, 40 fms., March 1966 (one arm removed for dissection).

Etymology: In honor of Harvey R. Bullis, Jr., Director, Exploratory Fishing and Gear Research Base, Bureau of Commercial Fisheries, Pascagoula, Mississippi.

DISCUSSION

H. L. Clark's (1921) excellent review of the Ophidiasteridae renders unnecessary an extensive discussion of the other genera within this family. His very useful key need not be altered to accommodate *Drachmaster*, as the major character separating this new genus from the other genera in the family is the presence of the internal plates connecting the plates in transverse series.

A KEY TO THE CARIBBEAN SPECIES OF OPHIDIASTERIDAE

1. With internal plates connecting the outer plates in transverse series *Drachmaster bullisi*
- Without internal plates 2
2. Complete covering of imbricating scales *Copidaster lymani*
- Without covering of imbricating scales 3
3. Abactinal plates in regular longitudinal series 4
- Abactinal plates not in regular longitudinal series 7

4. Papular areas regularly present below inferomarginals 5
 Papular areas not present below inferomarginals (isolated papulae
 sometimes present) *Tamaria floridae*
5. Papular areas in eight series 6
 Papular areas in ten series *Hacelia superba*
6. Three rows of spines on adambulacrals *Ophidiaster bayeri*
 Two rows of spines on adambulacrals *Ophidiaster guildingi*
7. Adambulacral armature spiniform, subprismatic; arms triangular
 in cross-section (in adult) *Narcissia trigonaria*
 Adambulacral armature granuliform; arm cylindrical 8
8. Large tumid plates occurring randomly on disc and arms
 *Linckia nodosa*
 No tumid plates *Linckia guildingi*

ACKNOWLEDGMENTS

I wish to thank Dr. Harvey R. Bullis and the Fish and Wildlife Service for providing the material on which this paper is based. I also appreciate the help of Dr. Mary Gardiner, whose knowledge of Greek was called upon to aid me with the etymology of *Drachmaster*. Dr. David. L. Pawson, of the Smithsonian Institution, was, as always, patient and helpful with suggestions during the preparation of the manuscript.

BIBLIOGRAPHY

- CLARK, A. H. 1948. Two new starfishes and a new brittle-star from Florida and Alabama. *Proc. Biol. Soc. Washington*, vol. 61, pp. 55-66, 1 pl.
- CLARK, A. M. 1967. Notes on asteroids in the British Museum (Natural History). V. *Nardoa* and some other ophidiasterids. *Bull. Brit. Mus. (Nat. Hist.) Zoology*, vol. 15, no. 4, pp. 169-198, 6 pls.
- CLARK, H. L. 1921. The Echinoderm Fauna of Torres Strait: its composition and its origin. Carnegie Institution of Washington, Dept. of Marine Biology, vol. X, viii, 223 pp., 38 pls.
- FISHER, W. K. 1911. Asteroidea of the North Pacific and adjacent waters. Pt. I. Phanerozonia and Spinulosa. *U.S. Nat. Mus. Bull.* 76, vi, 419 pp., 122 pls.
- GRAY, J. E. 1841. A synopsis of the genera and species of the class Hypostoma (Asterias, Linn.). *Ann. Mag. nat. Hist.* 6: 175-184; 275-290.
- SLADEN, W. P. 1889. Report on the Asteroidea collected by H.M.S. "Challenger" during the years 1873-1876. Vol. XXX, xli: 893 pp., 117 pls.



Downey, Maureen E. 1970. "Drachmaster bullisi new genus and species of Ophidiasteridae (Echinodermata: Asteroidea), with a key to the Caribbean species of the family." *Proceedings of the Biological Society of Washington* 83, 77–82.

View This Item Online: <https://www.biodiversitylibrary.org/item/107535>

Permalink: <https://www.biodiversitylibrary.org/partpdf/44881>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Biological Society of Washington

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.