# Giant Ringtail Possums (*Marsupialia*, *Pseudocheiridae*) and Giant Koalas (*Phascolarctidae*) from the Late Cainozoic of Australia

# MICHAEL ARCHER, KAREN BLACK AND KERRY NETTLE

School of Biological Science, University of New South Wales, Sydney N.S.W. 2052

Archer, M., Black, K. and Nettle, K. (1997). Giant Ringtail Possums (Marsupialia, Pseudocheiridae) and Giant Koalas (Phascolarctidae) from the late Cainozoic of Australia. Proceedings of the Linnean Society of New South Wales 117: 3-16

While seven pseudocheirids have been described from the late Oligocene to early Miocene and five from the Plocene of Australia, none are known to have been confined to the Pleistocene. We describe here a giant ringtail, *Pseudokoala cathysantamaria* n. sp., from the early Pleistocene Portland Local Fauna (lower Nelson Bay Fm) of Victoria. We review the generic distinction of the Miocene–Pliocene *Corracheirus* Pledge, 1992 and conclude that it is a junior synonym of *Pseudokoala* Turnbull and Lundelius, 1970. The palaeohabitat for all species of *Pseudokoala* appears to have been rainforest. A second pseudocheirid in the Portland Local Fauna is most similar in morphology and size to the still-living *Pseudocheirus peregrinus*, a species whose habitat range also includes rainforest. We describe the first known upper and additional lower molars for the giant koala *Cundokoala yorkensis* Pledge, 1992, from a Pleistocene deposit in Wellington Caves, New South Wales, a taxon previously only known from Pliocene sediments of South Australia. The Portland giant ringtail and Wellington giant koala add two more families to the already extensive list that exhibited Pleistocene gigantism. Previously, only one other lineage (tree kangaroos) of arboreal mammals has been known to have undergone gigantism.

Manuscript received 6 June 1996, accepted for publication 23 October 1996.

KEYWORDS: Koalas, Phascolarctidae, Pleistocene gigantism, Pseudocheiridae, Quaternary, ringtail possums.

### **INTRODUCTION**

The record for ringtail possums (Pseudocheiridae) begins in the late Oligocene. From sediments of this age in South Australia, Woodburne et al. (1987a) and Pledge (1987b) describe four species of *Pildra*, two of *Marlu* and one of *Paljara*. Archer (1992) reports over twelve additional unnamed taxa from the Oligo-Miocene sediments of Riversleigh, northwestern Queensland. Archer and Bartholomai (1978) note a Pseudochirops from the late Miocene of Alcoota, Northern Territory. Turnbull and Lundelius (1970; augmented by Turnbull et al. 1987) describe Pseudokoala from the early Pliocene Hamilton Local Fauna as well as two species of 'Pseudocheirus' which may (Archer 1984) represent the otherwise modern genus Petauroides. Another species of Pseudochirops, from the early Pliocene Bluff Downs Local Fauna of northeastern Queensland, is under study (B. Mackness and M. Archer). Pledge (1992) describes a giant species of Corracheirus from the late Miocene to Pliocene Curramulka Local Fauna of southern South Australia as well as a single tooth of another large pseudocheirid from the Plio-Pleistocene Cement Mills Local Fauna of southeastern Queensland. Until now, however, no giant pseudocheirids have been reported from undoubted Pleistocene sediments. The tooth described here as Pseudokoala cathysantamaria, from early Pleistocene

deposits at Portland, Victoria, is the only ringtail possum confined to the Pleistocene and the only giant ringtail known from the Quaternary.

Fossil koalas (Phascolarctidae) are similarly known from late Oligocene and younger sediments. Stirton (1957), Stirton, Tedford and Woodburne (1967), Woodburne et al. (1987b) and Springer (1987) describe six species in the Oligo-Miocene genera *Perikoala, Litokoala* and *Madakoala*. Black and Archer (in press) name another Miocene genus from Riversleigh and Black (in prep.) reports additional Miocene taxa from Riversleigh. Pledge (1992) describes a Miocene to Pliocene species of the giant *Cundokoala*. Bartholomai (1968) describes a large Plio-Pleistocene Queensland species of the modern genus *Phascolarctos*. Pledge (1987a) describes another large *Phascolarctos* from the early Pliocene of South Australia. We describe here isolated teeth, from Pleistocene deposits of Wellington Caves, New South Wales, that appear to represent the same Miocene/Pliocene South Australian species described by Pledge (1992) as *Cundokoala yorkensis*. The Wellington material includes the first known upper molars for this taxon.

Material described is registered in the palaeontological collections of three institutions as follows: AM F numbers represent the Australian Museum fossil collection; P numbers represent the palaeontological collections of the Museum of Victoria; SAM P numbers represent the palaeontological collections of the South Australian Museum. Dental terminology used here follows Luckett (1993) for molar homology, Flower (1867) for premolar homology and Archer (1984) and Woodburne et al. (1987a,b) for molar morphology.

### SYSTEMATICS

Class Mammalia Linneaus, 1758 Superorder Marsupialia Illiger, 1811 Order Diprotodontia Owen, 1866 Superfamily Phalangeroidea Weber, 1928 Family Pseudocheiridae Winge, 1893

Pseudokoala Turnbull and Lundelius, 1970

# Type species:

Pseudokoala erlita Turnbull and Lundelius, 1970

### Additional species:

Pseudokoala curramulkensis (Pledge, 1992) and Pseudokoala cathysantamaria n. sp.

### Revised generic distribution:

Corra Lynn Cave, Yorke Peninsula, South Australia; Hamilton Local Fauna, southwestern Victoria; Portland Local Fauna, Nelson Bay Formation, Portland, Victoria.

### Revised age range for the genus:

The Hamilton Local Fauna is interpreted (Rich et al. 1991) to be early Pliocene in age. The Curramulka Local Fauna is interpreted (Pledge 1992) to be late Miocene to Pliocene in age; the Nelson Bay Formation is interpreted (Flannery and Hann 1984) to be early Pleistocene in age.

# Revised generic diagnosis:

Species of *Pseudokoala* are distinguished from *Marlu praecursor* by their continuous postmetacristid and preentocristid, and posthypocristid which does not extend to the

lingual side of the crown. They are distinguished from *M. kutjamarpensis* by their continuous postmetacristid and preentocristid, connection of the cristid obliqua to the metacristid and their truncated posthypocristid. They are distinguished from species of *Pildra* and *Paljara* in their lack of an entostylid, presence of a buccal buttress on the protoconid, *en echelon* overlap of the postmetacristid and preentocristid, large size, and truncated posthypocristid. They are distinguished from species of *Pseudocheirus*, *Petauroides* and *Hemibelideus* in having a truncated posthypocristid, being much larger in size, having a second buccal buttress on the protoconid, not having steep entoconid blades and in lacking a protostylid. They are distinguished from species of *Pseudochirops* in having a truncated posthypocristid, being larger in size, having a second buccal buttress on the protoconid, blades, having a second buccal buttress on the protoconid, not having steep entoconid blades, having a truncated posthypocristid, being larger in size, having a second buccal buttress on the protoconid, not having steep entoconid blades, having the cristid obliqua connected to the metacristid, and in lacking a protostylid.

### Pseudokoala curramulkensis (Pledge, 1992)

### Emendation of dental homology for the paratype and referred specimen:

Pledge (1992) notes three specimens including the Holotype. The homology of the teeth (using the M1-4 molar homology system of Luckett 1993) given in table 5 needs emendation as follows. SAM P29901 (referred specimen) is a right dentary fragment with  $M_2$  (not  $M_1$ ) and alveoli for  $M_1$  and  $M_{3-4}$ . SAM P31792 (Paratype) is a left dentary fragment with  $M_{2-3}$  (not  $M_{1-2}$ ) and alveoli for  $M_1$  and  $M_4$ .

### Pseudokoala cathysantamaria n. sp.

# Holotype:

P173650, an isolated  $LM_1$  (Fig. 1).

### Type Locality:

Interpreted to be Nelson Bay, Portland, Victoria, although no specific locality details are recorded in the Museum of Victoria Palaeontology register (B. Thompson, pers. comm. to M. Archer), nor are there any details on the specimen label. However, it is registered within a series of fossils, including P173649 (a dentary of *Pseudocheirus* sp. cf. *P. peregrinus*; see below), for all of which the locality data is Nelson Bay, Portland, Victoria.

### Age:

Marine invertebrates from the Lower Nelson Bay Formation, the probable source of the Holotype, are early Pleistocene in age (Flannery and Hann 1984).

# Etymology:

In honour of Cathy Santamaria for her constant interest and much appreciated encouragement for palaeontological research in Australia.

	N	Measurements of Pseudokoala material. All measurements in millimetres.												
Specimen	P3 L	W	M1 L	AW	PW	M2 L	AW	PW	M3 L	AW	PW	M4 L	AW	PW
MV P173650	Dist	(Supil)	10.7	5.20	5.60	marth	-log	sing, si	1.1.1		15110	in the second	120	
SAM P26542			10.9	4.51	5.20									

TABLE 1

PROC. LINN. SOC. N.S.W., 117. 1997



Figure 1. A-A', Holotype of *Pseudokoala cathysantamaria* n.sp, Museum of Victoria P173650, occlusal stereopair of left  $M_1$ , from the early Pleistocene of Portland, Victoria; B, *Pseudokoala erlita*, NMV P54159, left  $M_1$ , occlusal stereopair; C, *Pseudocheirus* sp. cf. *peregrinus*, Museum of Victoria P173649, a right dentary fragment with  $M_{1-4}$ , from the early Pleistocene of Portland, Victoria . Bar indicates 5mm.

### Species diagnosis:

*Pseudokoala cathysantamaria* is a giant ringtail possum with molar teeth at least 20% larger than those of *P. erlita* (Turnbull et al. 1987) but comparable in size (Table 1) to *P. curramulkensis* (Pledge 1992; see below). It is also distinguished from *P. erlita* by having an extra anterior vertical buttress (as well as a posterior one) on the buccal flank of the protoconid, an anterobuccal vertical trigonid cleft anterior to the extra vertical buttress, an anterolingual basal cingulum and cingular pocket, a lingually convex metacristid, posterior elongation of the postentocristid, and mesostylids between the bases of the hypoconid and protoconid. Although the missing anterior half of  $M_1$  for *P. curramulkensis* prohibits comparisons in this area, the tooth of *P. curramulkensis* is smaller, appears to lack at least the posterior mesostylid, may lack the anterior flexure on the cristid obliqua, and has a much shorter postentocristid. In this feature, the three species appear to form a gradient with the postentocristid of *P. erlita* the least extended and that of *P. cathysantamaria* the most.

# Description:

The Holotype (Fig. 1) is the least worn of any *Pseudokoala* specimen known enabling details of dental morphology in the genus to be clarified.  $M_1$  has a large, medially situated protoconid (as in *P. erlita*) which is connected by a steep anterior paracristid to a tiny possible paraconid (not evident in *P. erlita* and unknown for *P. curramulkensis*), and by a gently inclined posterior metacristid to a poorly distinguished metaconid (as in at least *P. erlita*). Two prominent, vertical buccal buttresses subtend the protoconid. There is a tiny, anterolingual basin defined by a small, basal anterolingual cingulum, an anterobuccal cingula and a marked, vertical, anterobuccal trigonid cleft. Between the cristid obliqua and posterior buccal buttress are two basal mesostylids.

The following features of *P. cathysantamaria* are also evident (where they can be checked) in the holotype of P. curramulkensis and P. erlita. There is no protostylid. At the posterior end of the metacristid is a topographically defined metastylid immediately buccal to the anterior end of the preentocristid. The cristid obliqua does not run directly to the metacristid but instead veers anterobucally to tangentially contact the metacristid. There are pronounced crenulations on the lingual side of the cristid obliqua. There is en echelon overlap of the postmetacristid over the preentocristid. The paracristid, metacristid and postentocristid are elongate and together define a composite blade that extends the length of the whole tooth. The entoconid blades are not steeply inclined. There is no entostylid. The posthypocristid is truncated at the lingual end before it reaches the postentocristid. It stops far short of the posterolingual corner of the tooth, leaving the posterior end of the postentocristid (which extends posteriorly further than the same crest in the other species of *Pseudokoala*) to define the edge of the crown. In terms of relative height on the crown, the protoconid is just higher than the metaconid which is higher than the entoconid which is higher than the metastylid which is higher than the hypoconid which is higher than the paraconid.

### Discussion:

Estimates of body size for *Pseudokoala cathysantamaria*, based on tooth and body size in *Phascolarctos cinereus*, suggest that this ringtail may have weighed 9–10 kg, far larger than any living ringtail possum. However, it is only slightly larger than the Curramulka giant ringtail described by Pledge (1992) as *Corracheirus curramulkensis*. Pledge (1992) suggests that species of *Corracheirus* (viz. *C. curramulkensis*) are differentiated from those of *Pseudokoala* (viz. *P. erlita*) by three features: 1, larger size; 2, having the postprotocristid continuous with the cristid obliqua rather than extending to the metastylid; and 3, having a continuous entocristid with a simple metastylid flexure.

Large size is not normally regarded to be a basis for establishment of a monotypic genus (in this case for *curramulkensis* Pledge). Considering the second feature, there are three specimens of P. curramulkensis noted in the type description: the Holotype with part of M<sub>1</sub> and M<sub>2-4</sub>, and SAM P29901 with M<sub>2</sub> which support the distinction with the postprotocristid appearing to make no contact with the metastylid; and SAM P31792 with  $M_{2-3}$  where the postprotocristid bypasses (but is touched by) the cristid obliqua to make contact with the metastylid flexure. Consequently, this feature appears to be variable and does not distinguish species of Corracheirus from those of Pseudokoala. Considering the third distinguishing feature, the Holotype and SAM P29901 do support the suggestion that the preentocristid is continuous with metacristid via a simple metastylid flexure. However, the condition in SAM P31792 is much closer to the P. erlita condition in that there is only a very tenuous direct connection linking the preentocristid and the metacristid via a fine and low bridge of enamel buccal to the metastylid. In this specimen, the preentocristid and metacristid really overlap en echelon, with the preentocristid passing anterobuccal to the posterior end of the metacristid to make contact with the postprotocristid, a condition closer to that seen in P. erlita.



Figure 2. Cundokoala yorkensis, referred material from Wellington Caves, New South Wales. A-A', AMF98885, a left  $M^1$ . B-B', AMF98886, a right  $M_1$ . C-C', AMF98889, a right  $M_2$ . D-D', AMF98890, a left  $M_3$ . Bar indicates 5mm.

Conversely, comparison of SAM P31792 with the  $M_3$  or  $_4$  of *P. erlita* illustrated in Turnbull, Rich and Lundelius (1987, Fig. 2C) suggests basic overall similarity in all key features (except, possibly, the degree to which the metacristid is lingually concave and the degree of posterior development of the postentocristid; see below). *Corracheirus curramulkensis, P. erlita* and *P. cathysantamaria* also exhibit a striking synapomorphy of  $M_1$ : truncation of the posthypocristid such that it stops well short of the lingual side of the crown and does not contact any other structure. This condition differs from all other pseudocheirids where the posthypocristid closely approximates the postentocristid (or the entostylid) at or near the lingual margin of the tooth. All three taxa also exhibit a tendency to posteriorly extend the postentocristid, a feature best-developed in  $M_1$ . Finally, all three taxa appear to be united in their tendency towards gigantism. For these reasons, we suggest that *Corracheirus* Pledge, 1992 is better regarded as junior synonym of *Pseudokoala* Turnbull and Lundelius, 1970 which then contains three species: *erlita* Turnbull and Nettle (this paper), *erlita* being the type species of the genus.

Within *Pseudokoala*, *P. cathysantamaria* appears to be the sister-group of *P. curra-mulkensis*, these two sharing as synapomorphies extreme gigantism and greater posterior development of the postentocristid. Although the trigonid is unknown for *P. curra-mulkensis*, that of *P. cathysantamaria* shows two prominent buccal buttresses in contrast to one in *P. erlita*, the second buttress being an autapomorphic condition (or perhaps a synapomorphic condition shared with *P. curramulkensis*). The single buttress of *P. erlita* may be the homologue of the vertical protostylid ridge in the same position in species of *Marlu* (or the homologue of the protostylid in, e.g., species of *Pseudochirops*, *Pseudocheirulus*).

Species of *Pseudokoala* share distinctive features of  $M_1$  with species of *Marlu* Woodburne, Tedford and Archer, 1987 including: failure of the cristid obliqua to <u>directly</u> contact the postmetacristid (a deep crevice intervening); a crenulated cristid obliqua; *en echelon* overlapping of the postmetacristid and the preentocristid (in *M. kutjamarpensis* but not *M. praecursor*); a vertical buttress but no protostylid on the buccal flank of the protoconid; no entostylid; and gently inclined (rather than steeply inclined) entoconid blades. This intergeneric relationship was originally suggested by Woodburne, Tedford and Archer (1987; Fig. 23) on the basis of synapomorphies exhibited primarily by the upper molars. The new species *P. cathysantamaria* exhibits the same combination of synapomorphies, lending support to the hypothesis of a *Pseudokoala/Marlu* clade.

Marked differences between species of *Marlu* and *Pseudokoala* include the truncated posthypocristid of species of *Pseudokoala* and their better-developed entoconid blades. Differences in the nature of the connection between the preentocristid and metacristid vary within at least *P. curramulkensis* (see above) making intergeneric contrasts here of questionable value.

In terms of patristic relationships, the early Miocene *Marlu kutjamarpensis* exhibits no features that would preclude it being ancestral to the Pliocene/Pleistocene species of *Pseudokoala*.

# Pseudocheirus Ogilby, 1837 Pseudocheirus sp. cf. P. peregrinus (Boddaert, 1785)

Specimen:

P173649, a right dentary with  $M_{1-4}$  (Fig. 1).

### Locality:

Nelson Bay, Portland, Victoria. Label data also records: 'Monash Univ. field trip (1973?)' and '# 280 004'.

# Age:

The marine invertebrate assemblage of the Lower Nelson Bay Formation, source of P173649, is early Pleistocene in age (e.g., Flannery and Hann 1984).

### Description:

P173649 is a Pseudocheirus peregrinus-sized ringtail possum (Table 2) with overall similarity to P. peregrinus and P. occidentalis. It is, however, distinguished from these modern species (most individuals but not all, these features being somewhat variable in at least P. occidentalis) in that: the preentocristid is connected to the metastylid on  $M_{2-4}$ . and the metaconid on  $M_1$  is less distinct. It is distinguished from *Pseudocheirulus her*bertensis in that: the preentocristid is connected to the metastylid; the M<sub>1</sub> lingual parastylid is not developed; and there is no notch between the postmetacristid and metastylid. It is distinguished from P. caroli and P. forbesi in that: the preentocristid is connected to the metastylid; the M1 lingual parastylid is not developed; the preentocristid is strongly bladed; and there is no notch between postmetacristid and metastylid. It is distinguished from *P. canescens* and *P. mayeri* in that: the preentocristid is connected to the metastylid; the preentocristid is strongly bladed; and there is no notch between postmetacristid and metastylid. It is distinguished from Hemibelideus lemuroides in that: the buccal shelf is not prominent; and the M1 metaconid is less distinct. It is distinguished from *Petauroides volans* in that: the preentocristid is connected to the metastylid; there is no entostylid ridge; and the M<sub>1</sub> metaconid is less distinct. It is distinguished from Pseudochirops albertisii in: not having a protostylid basin on M1; lacking a prominent buccal shelf; lacking an entostylid ridge; lacking a posterobuccal trigonid basin; and in having a less distinct  $M_1$  metaconid. It is distinguished from P. corinnae in that: the preentocristid is connected to the metastylid; it lacks a protostylid basin on M<sub>1</sub>; it lacks a prominent buccal shelf; it lacks an entostylid ridge; it has no posterobuccal trigonid basin; and the M<sub>1</sub> metaconid is less distinct. It is distinguished from *P. cupreus* in: not having a protostylid basin on  $M_1$ ; lacking a prominent labial shelf, the  $M_4$  cristid obliqua is not angulate, no entostylid ridge, no posterobuccal trigonid basin; and in having a less distinct metaconid on  $M_1$ . It is distinguished from P. archeri in: not having a protostylid basin on  $M_1$ ; having an  $M_4$  cristid obliqua that is not angulate; lacking an entostylid ridge; lacking a posterobuccal trigonid basin; and in having a less distinct metaconid on M<sub>1</sub>. It is distinguished from *Petropseudes dahli* in: not having a protostylid basin on M<sub>1</sub>; lacking a prominent buccal shelf; lacking a posterobuccal trigonid basin; and in having a less distinct metaconid on M1. It is distinguished from the species of all extinct ringtail genera in having: a well-developed M<sub>1</sub> protostylid; a preentocristid connected to the metastylid; a strongly bladed preentocristid; an entoconid that is positioned anterior to the hypoconid; a protoconid positioned lingual to the midline of M<sub>1</sub>; and a less distinct metaconid on  $M_1$ .

Measurements of <i>Pseudocheirus</i> material. All measurements in millimetres.														
Specimen	P3 L	W	M1 L	AW	PW	M2 L	AW	PW	M3 L	AW	PW	M4 L	AW	PW
MV P173649			4.16	1.96	2.22	4.06	2.15	2.30	4.05	2.15	2.27	4.36	1.88	2.15
AM M4046	3.51	1.58	3.98	1.74	2.29	3.84	2.12	2.54	4.03	2.37	2.60	4.74	2.43	2.30

TABLE 2

### The early Pleistocene habitat of Portland:

The presence of an ektopodontid and *Pseudokoala cathysantamaria* in the Portland assemblage suggests a rainforest component in the palaeoenvironment. All ektopodon-

tids occur in assemblages that have been interpreted (e.g. Archer, Hand and Godthelp 1995) to represent rainforest communities or communities that include rainforest: the late Oligocene Ditjimanka and Ngama Local Faunas, South Australia; the early and middle Miocene assemblages of Riversleigh, Queensland; the ?early or middle Miocene Kutjamarpu LF, South Australia; and the early Pliocene Hamilton LF, Victoria. The late Miocene to Pliocene Curramulka LF of South Australia, which contains *Pseudokoala curramulkensis*, is also regarded (Pledge 1992) to represent wet sclerophyll forest and/or rainforest, as is the Hamilton assemblage (Turnbull and Lundelius 1970, Turnbull, Rich and Lundelius 1987) which contains *P. erlita*. The occurrence in the Portland assemblage of *Pseudocheirus* sp., *cf. P. peregrinus* does not conflict with this interpretation because, although primarily an inhabitant of dense understorey vegetation, some modern populations of *P. peregrinus* extend well into rainforest.

### Order Diprotodontia Owen, 1866

Suborder Vombatiformes Woodburne, 1984 Infraorder Phascolarctomorphia Aplin and Archer, 1987 Family Phascolarctidae Owen, 1839

Cundokoala yorkensis Pledge, 1992

### Holotype:

SAM P24904, left dentary with  $M_{1-4}$  and alveolus for P3.

### **Referred** material

(Fig. 2) and (Table 3): SAM P24905, partial left M<sub>4</sub> in its alveolus. AM F98885, L M<sup>1</sup>; AM F98886, R M<sub>1</sub>; AM F98887, L M<sub>2</sub>; AM F98888, L M<sub>2</sub>; AM F98889, R M<sub>2</sub>; and AM F98890, L M<sub>3</sub>.

Specimen	Р3	himer	M <sup>1</sup>	M <sup>1</sup>			M <sup>2</sup>			M <sup>3</sup>			M <sup>4</sup>		
	L	W	L	AW	PW	L	AW	PW	L	AW	PW	L	AW	PW	
AM F98885	0000	W Gro	12.1	12.3	11.0	in have	RICIOS	ene lavo	e origi	smin	alai h	ILS. SA	10. 20		
Tagailen 190	P <sub>3</sub> L	W	M <sub>1</sub> L	AW	PW	M <sub>2</sub> L	AW	PW	M <sub>3</sub> L	AW	PW	M <sub>4</sub> L	AW	PW	
P24904	usobu	guild g	11.8	7.0	8.1	12.9	8.1	8.1	12.9	8.0	8.2	12.7	7.4	7.3	
AM F98886			11.5	6.5	8.0										
AM F98887						11.5	6.9	6.8							
AM F98888						11.2	7.2	7.0							
AM F98889						11.3	7.1	6.9							
AM F98890									11.4	7.3	6.7				

### TABLE 3

Measurements of all known specimens of *Cundokoala yorkensis*. All measurements in millimetres. Abbreviations: L, length; AW, anterior width; PW, posterior width.

### Revised distribution:

*Cundokoala yorkensis* is known from Corra Lynn Cave, the Curramulka Local Fauna, Yorke Peninsula, South Australia and now Wellington Caves, New South Wales.

### Revised age range:

The Curramulka Cave site of the typical material is interpreted by Pledge (1992) to be Miocene to Pliocene in age. The exact locality and age for all but one (AM F98886) of the Wellington Caves specimens is unknown. They were collected from the surface of spoils piles dumped outside the newly restored entrance to the Phosphate Mine. The material on these piles had been obtained during excavation of tourist paths in the Wellington Caves complex in late 1995. The source of the material for each pile was not certainly known but thought by excavation workers at the site to be various localities within the Phosphate Mine and Bone Cave. Although most of the taxa recovered to date from the Wellington Caves complex of sediments have been interpreted to be Pleistocene in age, Hand, Dawson and Augee (1988), L. Dawson et al. (in prep.) and Osborne (1983) have demonstrated that some deposits in the Wellington Caves complex (e.g., in the entrance doline of Big Sink and others in the Phosphate Mine) are Pliocene in age. Therefore the age of the *Cundokoala yorkensis* material obtained from the spoils piles could be Pleiocene, Pleistocene or both.

AM F98886, however, was obtained by one of us (MA) in company with H. Godthelp, A. Gillespie, A. Musser et al. from a newly-excavated pathway in the Phosphate Mine. Material excavated at the same time included an isolated lower molar of *Diprotodon* sp. *cf. D. optatum* (only known from the Pleistocene), a dentary of *Protemnodon* sp. *cf. P. roechus* (only known from the Pleistocene), abundant dentary and maxilla fragments of *Aepyprymnus rufescens* (Pleistocene and living species) and a dentary of *Onychogalea unguifera* (living species). Although other taxa obtained at the same time have yet to be identified, nothing contradicts a Pleistocene age for this assemblage. We would conclude, pending a thorough analysis of the rest of the fauna from this deposit, that Wellington Caves *C. yorkensis* is Pleistocene in age, giving the species a Pliocene-Pleistocene age range.

### **Revised diagnosis**

Cundokoala yorkensis differs from all other phascolarctids: in being larger; having higher-crowned teeth; and in having a relatively short, massive dentary. It differs from all other phascolarctids except species of Phascolarctos in having: a larger paraconule and neometaconule on M<sup>1</sup>; a more lingually positioned protoconid and a larger protostylid on  $M_1$ ; well-developed lingual columnar stylids on the metaconid and entoconid of  $M_{1_2}$ 4; and a well-developed buccal cingulum and metastylid fold on  $M_{1-4}$ . It differs from species of Litokoala in: lacking the posterobuccal crest which extends from the apex of the metaconid in M<sub>2-4</sub>: lacking an anteriorly displaced entoconid (relative to the hypoconid) on M<sub>4</sub>; lacking an anteriorly displaced metaconid (relative to the protoconid) on the  $M_4$ ; and in having the postprotocristid and cristid obliqua of  $M_4$  meeting in the transverse median valley on M<sub>4</sub> (in contrast to the parallel arrangement seen in Litokoala kanunkaensis) It differs from species of Madakoala and Perikoala: in having: a larger parastyle on M1; a more crenulate, less linearly-oriented paraconule and neometaconule on M<sup>1</sup>; a paraconule that connects anterobuccally to the anterior cingulum on M<sup>1</sup>; an entostylid ridge on M1-4; a more lingual junction of the postprotocristid and cristid obliqua; and in lacking protoconid-metaconid and hypoconid-entoconid crests.

### Description

The new Wellington Caves materials (AM F98885 to AM F98890) augment understanding about the morphology of this species. AM F98885 is an unworn, relatively square selenodont  $M^1$  that tapers posteriorly. It is morphologically very similar to the  $M^1$  of species of *Phascolarctos* and differs mainly in being relatively wider anteriorly. The paracone and metacone are similar in height and are the tallest cusps on the tooth followed by the metaconule and protocone. The apex of the protocone lies lingually opposite that of the paracone. The apex of the metaconule lies lingually opposite and slightly anterior to that of the metacone. The lingual bases of the paracone and metacone and the anterolingual base of the protocone are highly crenulate. The protocone and metacone and their associated crests are slightly obliquely (anterolingually) oriented. The tooth is bisected bucco-lingually by a deep transverse median valley and antero-posteriorly by a relatively deep longitudinal valley. The buccal tooth margin is mildly convex sloping anterolingually at the anterolingual tooth margin and curving posterolingually around the buccal margin of the metacone. The lingual bases of the protocone and metaconule slope gently towards the base of the crown. The lingual bases of the paracone and metacone slope more steeply into the longitudinal tooth valley. The triangular buccal surface of the paracone is reduced relative to the metacone as is the paracone buccal margin. The buccal basin of the paracone is closed and deep. The buccal basin of the metacone is comparatively shallower but remains closed. A ridge-like stylar shelf extends along the length of the buccal tooth margin. The relative heights of stylar cusps in descending order are as follows: stylar cusp C, B, A, D and E. The preparacrista, postparacrista, premetacrista and postmetacrista which make up the buccal selene of the tooth, are distinct linear crests which extend from the apices of the paracone and metacone respectively. The lingual selene is offset posteriorly and is composed of a relatively linear preprotocrista and premetaconule crista and a crescentic postmetacrista and highly crescentic postprotocrista. The postprotocrista and premetaconulecrista meet in the transverse median valley at a point slightly lingual to the longitudinal valley. The parastyle is poorly developed. It exists as a slight swelling of the anterior cingulum at the anterobuccal corner of the tooth. A shallow pocket is created between the anterior cingulum and the anterior base of the paracone. Lingually, the pocket is defined by a small, crenulate, but distinct paraconule which lies at the anterolingual base of the paracone. A short, non-cuspate, spur-like protostyle which originates from the preprotocrista at a point slightly lingual to the longitudinal valley, extends posteriorly along the buccal base of the protocone, terminating opposite the protocone apex. A well developed anterolingual paracrista connects the paraconule posteriorly to the apex of the paracone. A similarly welldeveloped posterolingual paracrista extends from the apex of the paracone into the junction of the transverse median and longitudinal valleys wherein it divides into two spurs which become part of the crenulation pattern of the tooth.

A small, crenulate, non-cuspate neometaconule lies in the longitudinal tooth valley at the anterolingual base of the metacone. It is connected to the metacone base by a weak anterolingual metacrista, which originates approximately half way down the base of the metacone. A posterolingual metacrista is absent. A crenulate, cingulum-like shelf runs along the anterior and anterolingual base of the protocone forming a shallow pocket between the cingulum and the protocone base. A weak anterolingually directed crest extends from the apex of the protocone to meet the anterolingual cingulum. A similar, but deeper pocket occupies the anterolingual base of the metaconule and is bounded by the posterior base of the protocone anteriorly, the anterior base of the metaconule posteriorly, the junction of the premetaconule crista and postprotocrista buccally, and a short, crescentic lingual cingulum lingually. The lingual cingulum effectively closes off the lingual exit of the transverse median valley and is continuous with a weak crest that extends anterolingually from the apex of the metaconule. Weak buccally directed spurs extend from the apices of the protocone and metaconule, fading down their respective bases towards the longitudinal valley. A well developed posterior cingulum is continuous with the postmetaconulecrista lingually and buccally and meets stylar cusp E at the posterobuccal tooth corner.

The  $M_1$  of the holotype is poorly worn and the anterolingual tooth corner is missing. Consequently, much of the crown morphology of the trigonid has been lost. In AMF98886 the preprotocristid is a well defined, linear crest which terminates anteriorly in a small paraconid. The preprotostylidcristid extends anterolingually to terminate at the base of the paraconid. Pledge (1992), in his description of the holotype  $M_1$ , suggests that *C. yorkensis* differs from all other phascolarctids in having a fine anterobuccal spur of the preprotocristid meet the preprotostylidcristid at the anterior tooth margin. Its absence in AM F98886 suggests this is a variable feature within the species, and the prepro-

tocristid spur is most probably part of the crenulation pattern of the tooth. A well-developed posterobuccal ridge extends from the protostylid apex in AM F98886. This crest is only vaguely discernible in the holotype. The buccal cingulum is better developed in AM F98886. The postprotocristid meets the anterior base of the cristid obliqua at a slightly more lingual position than in the holotype. The lingual columnar stylids of the metaconid and entoconid are large, crenulate and well preserved in AM F98886 whereas they are only represented by slight swellings in the holotype. The entostylid is a well-developed, cuspate swelling on the terminus of the postentocristid at the posterolingual tooth corner in AM F98886. AM F98887, AM F98888 and AM F98889 (M<sub>2</sub>) are morphologically similar to but smaller than the M2 of the holotype. Again, the holotype is highly worn, the apices of all major cusps are missing and the lingual tooth margin is damaged such that the columnar stylids are not preserved and the entostylid is missing. In contrast, all of the Wellington Caves M2 specimens show little or no wear. The columnar stylids are large, particularly that of the metaconid. The postentocristid curves lingually to the posterolingual tooth corner where it terminates in a well developed entocristid. The entoconulid, like in the holotype, is poorly developed. Well developed lingually directed crests extend from the apices of the protoconid and hypoconid and terminate just prior to reaching the longitudinal valley. These crests are not present in the holotype. The M<sub>2</sub>, AM F98890, is largely unworn and is slightly smaller than the M<sub>3</sub> of the holotype. The postmetaconid, preentocristid and postentocristid are more arcuate than in the holotype. The lingual buttresses of the metaconid and entoconid are better developed however the buccal bases of these cusps are more greatly expanded in the holotype. The metastylid is larger and the metastylid fold is more pronounced. The lingual faces of the protoconid and hypoconid are more steeply sloping and the lingually directed crests associated with these cusps are well developed (however they are absent in the holotype). Following an analysis of variation in dentitions of the modern species, it is evident that the above mentioned morphological differences between the referred material and the holotype fall within the boundaries of normal intraspecific variation.

# DISCUSSION

*Pseudokoala cathysantamaria* is an enormous ringtail possum and *Cundokoala yorkensis* an enormous koala. With the addition of these two to the Pleistocene of Australia, all families of Australian herbivorous mammals are now known to have had giants in the Pleistocene megafauna. They also significantly increase the ranks of known *arboreal* megafaunal species. Apart from one modestly large koala (*Phascolarctos stirtoni* Bartholomai, 1968), the only undoubted megafaunal arboreal species previously known from the Pleistocene was *Bohra paulae* Flannery and Szalay, 1982, a gigantic tree kangaroo from Wellington Caves, New South Wales.

### ACKNOWLEDGMENTS

We thank the Australian Research Committee, the University of New South Wales, the Queensland Museum, National Estate Programs (Queensland), the Riversleigh Society, Queensland National Parks and Wildlife Service, Australian Geographic Society, Royal Zoological Society of New South Wales, Linnean Society of New South Wales and private individuals for support that has enabled discovery and preparation of the diverse koalas and ringtail possums of Riversleigh. Henk Godthelp, Anna Gillespie and Anne Musser (all of the University of New South Wales) helped collect material from Wellington Caves. Armstrong Osborne (University of Sydney) and Michael Augee (University of New South Wales) organised the opportunity to examine the Wellington deposits. Mathew Crowther (University of New South Wales) confirmed the identifications of other Wellington mammals excavated with the tooth of *Cundokoala yorkensis*. Neville Pledge (South Australian Museum) and Tom Rich (Museum of Victoria) made materials available for study. Betty Thompson (Museum of Victoria) confirmed collection details of Portland materials. Suzanne Hand and Jenni Brammall (University of New South Wales) read and constructively criticised a draft of this paper.

### REFERENCES

- Archer, M. (1984). The Australian marsupial radiation. In 'Vertebrate zoogeography and evolution in Australasia' (Eds M. Archer and G. Clayton) pp. 633–808. (Hesperian Press: Perth).
- Archer, M. (1992). Ringtail possums (Pseudocheiridae, Marsupialia) from the Tertiary deposits of Riversleigh. *The Beagle* **9**: 257–258.

Archer, M. and Bartholomai, A. (1978). Tertiary mammals of Australia: a synoptic review. Alcheringa 2: 1-19.

- Archer, M., Hand, S.J., and Godthelp, H. (1995). Tertiary environmental and biotic change in Australia. In 'Paleoclimate and evolution, with emphasis on human origins' (Eds E.S. Vrba, G.H. Denton, T.C. Partridge, L.H. Burckle) pp. 77–90. (Yale University Press: New Haven).
- Bartholomai, A. (1968). A new fossil koala from Queensland and a reassessment of the taxonomic position of the problematical species, *Koalemus ingens* DeVis. *Memoirs of the Queensland Museum* **15**: 65–73.
- Black, K. and Archer, M. (in press). *Nimiokoala*, a new genus and species of koala (Marsupialia, Phascolarctidae) from Riversleigh, northwestern Queensland, with a review of *Litokoala* Stirton, Tedford and Woodburne, 1967. *Memoirs of the Queensland Museum*.
- Flannery, T.F. and Hann, L. (1984). A new macropodine genus and species (Marsupialia: Macropodidae) from the Early Pleistocene of southwestern Victoria. *Australian Mammalogy* 7: 193–204.
- Flannery, T.F. and Szalay, F. (1982). *Bohra paulae*, a new giant fossil tree kangaroo (Marsupialia, Macropodidae) from New South Wales, Australia. *Australian Mammalogy* **5**: 83–94.
- Flower, W.H. (1867). On the development and succession of teeth in the Marsupialia. *Philosophical Transactions of the Royal Society, London* **157**: 631–641.
- Hand, S.J., Dawson, L., and Augee, M. (1988). Macroderma koppa, a new Tertiary species of false vampire bat (Microchiroptera: Megadermatidae) from Wellington Caves, New South Wales. Records of the Australian Museum 40: 343–351.
- Luckett, W.P. (1993). An ontogenetic assessment of dental homologies in therian mammals. In 'Mammal phylogeny: Mesozoic differentiation, multituberculates, monotremes, early therians, and marsupials' (Eds F.S. Szalay, M.J. Novacek, M.C. McKenna) pp. 182–204. (Springer-Verlag: New York).
- Osborne, R.A.L (1983). Cainozoic stratigraphy at Wellington Caves, New South Wales. *Proceedings of the Linnean Society of New South Wales* **107**: 131–147.
- Pledge, N.S. (1987a). *Phascolarctos maris*, a new species of koala (Marsupialia: Phascolarctidae) from the Early Pliocene of South Australia. In 'Possums and opossums: studies in evolution' (Ed. M. Archer) pp. 327–330. (Surrey Beatty and Sons Pty Ltd: Sydney).
- Pledge, N.S. (1987b). Pseudocheirids (Marsupialia: Pseudocheiridae) from the Middle Miocene Ngama Local Fauna of northern South Australia. In 'Possums and opossums: studies in evolution' (Ed. M. Archer) pp. 639–679. (Surrey Beatty and Sons Pty Ltd: Sydney).
- Pledge, N.S. (1992). The Curramulka Local Fauna: a new late Tertiary fossil assemblage from Yorke Peninsula, South Australia. *The Beagle* **9**: 115–142.
- Rich, T.H., Archer, M., Hand, S.J., Godthelp, H., Muirhead, J., Pledge, N.S., Flannery, T.F., Woodburne, M.O., Case, J.A., Tedford, R., Turnbull, W.D., Lundelius Jr, E.L., Rich, L.S.V., Whitelaw, M.J., Kemp, A. and Rich, P.V. (1991). Appendix 1: Australian Mesozoic and Tertiary terrestrial mammal localities. In 'Vertebrate palaeontology of Australasia' (Eds P. Vickers-Rich, J.M. Monaghan, R.F. Baird, T.H. Rich ) pp. 1005–1058. (Pioneer Design Studio: Melbourne).
- Springer, M.S. (1987). Lower molars of *Litokoala* (Marsupialia: Phascolarctidae) and their bearing on phascolarctid evolution. In 'Possums and opossums: studies in evolution' (ed M. Archer) pp. 319–325. Surrey Beatty and Sons Pty Ltd: Sydney.
- Springer, M.S. (1993). Phylogeny and rates of character evolution among ringtail possums (Pseudocheiridae: Marsupialia). Australian Journal of Zoology **41**: 273–291.
- Stirton, R.A. (1957). A new koala from the Pliocene Palankarinna Fauna of South Australia. *Records of the South Australian Museum* 13: 71–81.
- Stirton, R.A., Tedford, R.H. and Woodburne, M.O. (1967). A new Tertiary formation and fauna from the Tirari Desert, South Australia. *Records of the South Australian Museum* **15**: 427–462.
- Turnbull, W.D. and Lundelius Jr, E.L. (1970). The Hamilton Fauna, a late Pliocene mammalian fauna from the Grange Burn, Victoria, Australia. *Fieldianna: Geology* **19**: 1–163.
- Turnbull, W.D., Rich, T.H.V. and Lundelius Jr, E.L. (1987). Pseudocheirids (Marsupialia: Pseudocheiridae) of the Early Pliocene Hamilton Local Fauna, southwestern Victoria. In 'Possums and opossums: studies in evolution' (Ed. M. Archer) pp. 689–713. (Surrey Beatty and Sons Pty Ltd: Sydney).
- Woodburne, M.O., Tedford, R.A. and Archer, M. (1987a). New Miocene ringtail possums (Marsupialia: Pseudocheiridae) from South Australia. In 'Possums and opossums: studies in evolution' (Ed. M. Archer) pp. 681–688. (Surrey Beatty and Sons Pty Ltd: Sydney).
- Woodburne, M.O., Tedford, R.A., Archer, M. and Pledge, N.S. (1987b). *Madakoala*, a new genus and two species of Miocene koalas (Marsupialia: Phascolarctidae) from South Australia, and a new species of *Perikoala*. In 'Possums and opossums: studies in evolution' (Ed. M. Archer) pp. 293–317. (Surrey Beatty and Sons Pty Ltd: Sydney).





Archer, Michael, Black, Karen H., and Nettle, K. 1997. "Giant ringtail possums (Marsupialia, Pseudocheiridae) and giant koalas (Phascolarctidae) from the late Cainozoic of Australia." *Proceedings of the Linnean Society of New South Wales* 117, 3–16.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/108631</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/47486</u>

Holding Institution MBLWHOI Library

**Sponsored by** Boston Library Consortium Member Libraries

**Copyright & Reuse** Copyright Status: In copyright. Digitized with the permission of the rights holder. License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.