TWO NEW SPECIES OF WOOD-BORING MOTHS FROM WEST AFRICA.

By Sir GEORGE F. HAMPSON, Bart.

Duomitus armstrongi, sp. nov. (Plate xvii, fig. 8).

♀. *Head* and *thorax* white ; antennae black ; palpi and sides of head black-brown ; prothorax and sides of metathorax with tufts of long black hair slightly glossed with metallic blue, the patagia with black spots near base and black tips ; pectus, legs and abdomen black-brown, the last with the dilated anal segment whitish, the rest of abdomen with tufts of long black hair on dorsum and at sides slightly glossed with metallic blue. *Forewing* grey, the base suffused with black-brown, the rest of wing with numerous black-brown spots which tend to become bars on terminal half below the cell and vein 4. *Hindwing* black-brown, with traces of deeper black-brown spots on terminal half.

Forewing with a forked veinlet in end of cell, vein 3 from long before end of cell, the median nervure bent upwards slightly just beyond it, vein 5 from above angle, 6 from well below upper angle, 7, 8 from angle, 9 absent, 10 from before angle, 11 from middle of cell ; hind wing with a forked veinlet in end of cell, vein 3 from long before angle, 5 from just above angle, 6, 7 coincident, 8 free. *Exp.* 44 mill.

GOLD COAST : Aburi (*L. Armstrong*) 1 ♀, type. The larva bores in the stem of coffee.

Genus MELISOMIMAS.

Melisomimas, Jord., Entom., xl, p. 127 (1907).

Melisomimas metallica, sp. nov.

Melisomimas grandis, Jord., Entom., xl, p. 127 (1907), non descr.—nec *Melisa grandis*, Holland.

Head, *thorax* and *abdomen* black, the tegulae with some white hairs at base, the patagia with some white hairs at base of upper edge ; coxae with some white hairs, the femora, tibia and tarsi white at tips ; abdomen with some metallic blue suffusion. *Forewing* brilliant metallic blue, the basal area suffused with black ; a more or less prominent tuft of white hair below base of cell. *Hindwing* brilliant metallic blue, some black suffusion at base of inner area ; some white in submedian fold below end of cell and sometimes some white beyond the cell above and below vein 5. *Exp.* 30–44 mill.

SIERRA LEONE : (*Major P. Smith*) 1 ♀ ; S. NIGERIA : Ibadan Distr. (*Dr. W. A. Lamborn*)* 2 ♀, type.

* [Bred from larvae boring in the bark of an *Albizia*.—ED.]

FURTHER NOTES ON THE BIONOMICS OF *TABANUS DITAENIATUS*, MACQ., AND *TABANUS TAENIOLA*, P. DE B.

By HAROLD H. KING, F.L.S., F.E.S.

(Government Entomologist, Anglo-Egyptian Sudan; Wellcome
Tropical Research Laboratories, Khartoum.)

(PLATE XXVI.)

In 1909, the writer collected egg-masses of the seroot *Tabanus taeniola*, P. de B., and reared some of the resulting larvae to approaching maturity, but failed to obtain the pupa. The following year he took a number of immature larvae of *Tabanus ditaeniatus*, Macq., and reared them to the adult stage. The various stages in the life-cycles of these two Tabanids, with the exception of the pupa of the former and the egg-mass of the latter species, were figured and described in this Bulletin.* The puparium of *T. taeniola* and egg-masses of *T. ditaeniatus* which are here figured and described were obtained during the summer of 1913.

***Tabanus taeniola*, P. de B.**

A nearly mature larva of this species was taken in wet soil at the edge of Khor Barboy, near Tonga, on the White Nile, on 20th June. It fed readily on earthworms till about 25th July, when it appeared to be mature. On 8th August it was found to have pupated; the eyes were coloured, but the rest of the pupa was yellowish-white. Later, the wings and other appendages darkened, and finally the abdominal markings of the adult could be distinguished. An adult male emerged on 19th August.

Pupa (Plate xxvi. figs. 3-5).—Length, 21 mm. Colour yellowish, thoracic tubercles and abdominal spiracles tinged with brown, the former bearing hairs. On the posterior third of each of the second to the seventh abdominal segments, inclusive, is a ring of backwardly projecting spines, shortest on the second segment and longest on the seventh, the spines yellowish, the ridge from which they arise chestnut-brown. The eighth segment terminates in a coronet of six teeth, chestnut-brown in colour, darker at the tips, approximately equal in size, arranged in a circle, the lateral teeth nearer to the dorsal than to the ventral teeth. Ventrally placed to this coronet is a transverse row of tiny teeth and dorso-laterally situated on either side is a comb of five teeth, the central ones being the longest and slightly longer than those on the ventral row.

***Tabanus ditaeniatus*, Macq.**

The egg-masses (Plate xxvi, figs. 1, 2) of this Tabanid were plentiful on grass growing in rain pools in the vicinity of Khor Barboy and at the junction of Khor Felus and the Sobat River. In shape they are very variable, some being long and narrow, others short and broad. Of the specimens collected the longest measured 20.5 mm. by 3 mm. and the shortest 8.5 mm. by 4 mm. The eggs are not covered with a secretion as are those of *T. biguttatus*, Wied. When freshly deposited the egg-mass is probably white to yellowish-white in colour, but all those taken varied from light to dark brown.

*Bull. Ent. Res., i, p. 102, and i, pp. 265-268.

The act of oviposition was not witnessed. The time occupied from the hatching of the egg to the emergence of the adult varied considerably. From an egg-mass which hatched on 24th June the first adult was obtained on 2nd November; while from another egg-mass which hatched on 25th July an adult was obtained on 11th September.

Of the egg-masses collected at Khor Barboy many were parasitised by an undetermined species of hymenopteron—probably a *Telenomus*.

Egg.—Length about 1.25 mm., of the typical Tabanid shape. As the embryo develops the ends of the egg darken, giving the egg-mass when viewed from a distance a general dark appearance.

Egg-masses of *Tabanus ditæniatus*.

Posterior view of eighth abdominal segment of pupa of *Tabanus tæniola*.

Lateral view of pupa
of *T. tæniola*.

Lateral view of anal segment of
pupa of *T. tæniola*.

NOTES ON AFRICAN CHALCIDOIDEA.—I.

By JAMES WATERSTON, B.D., B.Sc.,

Imperial Bureau of Entomology, London.

The following descriptions are based on material submitted by various collectors to the Imperial Bureau of Entomology. The types of the new species will be deposited in the British Museum.

Family AGAONIDAE.

In an interesting series of Fig Insects, comprising both caprifiers and inquilines, which have been forwarded by Dr. G. D. H. Carpenter from Uganda, are two species belonging to this family and representing respectively its main divisions.

The Sycophagine species appears to demand a new genus for its reception. Amongst its many noteworthy features, the most remarkable is the extreme development of the sub-apical spur of the first tibia into an organ comparable with the appendage to the mandibles so characteristic of the Agaonines.

For this new genus, the name *Sycoecus* (σῦκον, οἶκος) is proposed. *Sycoecus* affords a really extraordinary instance of homoplasy. The tibia is externally furnished with heavy, tooth-like spines, which, in conjunction with the flattened, small process, form a digging apparatus exactly similar to the mandible and appendage of *Pleistodontes*, etc. In these instances, therefore, the two sub-families have adopted the same device, but by modifying different organs. The tissue of the ovary, etc., of the fig is probably first dug into and torn by the teeth and then cut and swept aside by the lamina, which is in each case movable horizontally.

Both the Sycophagine (a few) and the Agaonine (very plentiful) occurred in the same species of fig, and sometimes together. There can, I think, be no doubt that the Agaonine species belongs to the genus *Agaon*, Dalm. (1818), about which little appears to be known at first hand. Dr. Carpenter's specimens may even be *A. paradoxum*, Dalm. (Svensk. Vet-Akad. Handl., xxxix, p. 69, 1818), but the thorax appears to be longer than in the genotype, while the petiole and first abdominal segment are maculate or banded. The original description is insufficient for specific determination. Having unsuccessfully tried to view the type, I have thought it better, meanwhile, to separate the Uganda examples from the Sierra Leone species.

Ashmead's table (Mem. Carn. Mus., i, no. 4, p. 233, 1904) should be corrected. In *Agaon* the head is less than $2\frac{1}{2}$ times as long as broad; the antennae have 11 not 12 joints; the third joint shows a distinct process, and the mandibles have 4 and not 3 teeth. *Agaon* appears to be most closely related to *Pleistodontes*, Saund. (Trans. Ent. Soc. Lond., 1883, p. 8), but the latter genus differs conspicuously in the antennae and legs.

***Agaon fasciatum*, sp. nov. (figs. 1 and 2).**

General colour pale yellow-brown, with darker head and sheath of ovipositor, first tergite with a medianly interrupted dark band.

♀. *Head* flattened greatly, $1\frac{1}{2}$ – $1\frac{3}{4}$ times as long as broad. Eyes reaching vertex, rather less than one-half length of head, placed mainly at the side and widely apart.

Vertex V-shaped, the sides clothed with short hairs ; frons with sharp median sulcus, the sides contiguous to one-third from vertex, widening thence to scrobes, V-shaped in transverse section, with scattered short hairs except behind the scapes. Clypeus produced in a rounded median projection, between which and the scrobes are many short, stiff hairs. Antennae (fig. 1) 11-jointed, set at less than one-fourth from mouth edge ; scrobes widely apart, narrow, slanting inwards and upwards, face triangularly excavated from this level to the edge of the clypeus ; scapes large, opposed to one

Fig. 1. *Agaon fasciatum*, sp. nov. ; a, antenna ; b, terminal sense-organ.

another along their much flattened triangular faces, with (at about one-half) a strong outwardly-directed process, which bears six short bristles ; second joint transverse, short, with six bristles ; third joint flat and round basally, thereafter drawn out apically to a sharp point, with one short, stout bristle ; 4th joint cylindrical, short, rising from a depression in the third near the base, having 3 short bristles. Joints 5-8 (of which 5 and 6, and 7 and 8 respectively, are subequal) are similarly built ; each bears near the base five or six very long tubular bristles and one or two more near the apex ; 9 is half as wide again as 8 and bears 2 rows of the long bristles, as do 10 and 11 ; it is apically produced into a neck ; 10 is produced into a neck in both directions ; the 11th segment bears a sense organ composed of about ten short leaf-like hairs.

Mouth-parts (fig. 2, a, b).—Mandibles powerful, symmetrical, with 4 teeth on inner edge ; one pair (apical and sub-apical) are equal, a third behind them shorter, a fourth small, with sometimes a trace of another behind. Under surface of the mandible with over a dozen sharp ridges ; on the inner superior angle there is a strong broadened process for muscle attachment. Hinged to the mandible, but not interfering with its mobility, is a very long, backwardly directed and apically slightly outcurved rasping blade, reaching to the occiput. This appendage breaks easily from the mandible ;

it is sharply serrate on the outer edge, and less distinctly so on its inner aspect; the lateral serrations are the terminal units of parallel rows (about 50) of teeth, numbering 15–20 per row. Maxilla laterally compressed and appearing like a blade between the saws attached to the mandibles; stipes in the form of a long, narrow strip (6:1), broadest near the apex; galea with two bristles; palpus entirely absent. Labium like a laterally compressed (5:1) spoon; palpi not certainly present; in their place and possibly homologous with them are two approximated slightly raised clear spots which give rise to a bristle; two minute distal bristles.

Fig. 2. *Agaon fasciatum*, sp. nov.; a, mandible and serrate appendage; a', mandible, under surface; b, trophi; c, metathoracic stigma; d, hind leg; d', apex of hind tibia; e, caudal stigma; f, cells at junction of submarginal and marginal veins; g, radius; h, receptaculum seminis; i, apex of ovipositor.

Thorax.—Noteworthy in this region are the reduction of the pronotum, the development of the prothoracic spiracle, the mesonotum and the propodaeum. Pronotum short, angularly emarginate behind, consisting of two triangular sclerites connected by a narrow, membranous area. On each triangle there is an irregular row of bristles along the inner edge, a few more on the outer, and one or two on the surface near the posterior angle. Prothoracic spiracle with a strong, terminal chitinous cap, which, along with part of the trachea, projects clear of the side; this structure and the tegula are probably the two thorns or spines with which Dalman says his insect is provided on the sides of the thorax. Mesonotum triangular before the suture, coming to a point anteriorly in the middle of the prothorax; no definite parapsidal furrows; 4–6 short bristles posteriorly, on each side of an incomplete median line.

Tegulae conspicuous. Axillae bare, externally rectangular. Scutellum with a median impressed triangular area; over 20 short bristles in two irregular rows before the suture and mainly inside the triangle. Metanotum distinct, ribbon-like; the posterior margin broadly excised in the middle, with 4–6 minute bristles at each side. Propodaeum similar to the metanotum, but about twice as broad. Spiracle oval, with the anterior end narrower and debouching on a distinct stalk or ridge, so that the whole structure is pear-shaped (fig. 2, c); 5 short bristles behind the spiracle. Mesophragma long, narrow and distinctly entering the abdominal cavity. Prosternum diamond shaped, slightly truncate anteriorly, while the posterior angle is bifid and free from the thoracic surface. Mesosternal area anteriorly concave, with a strong median ridge, and posteriorly with six minute bristles on each side.

Abdomen.—Petiole very broad and dark, almost coextensive with the propodaeum. The abdomen conic-ovate and compressed, with V-shaped hollow on basal segments. First tergite almost unchitinised medianly, dark at the sides, with about 30 minute bristles on each side. The other tergites bare, but nearer the spiracle are one or two minute bristles. Spiracle (fig. 2, e) more than twice as long as broad, with a sharp narrow indentation on the lower proximal edge at one-fourth from base. Sternites with the chaetotaxy obsolescent, but on the ploughshare-shaped fifth sternite are some short hairs in two ventral rows. Stylet narrow; longer on the upper edge, where are 4 hairs, and 2 others at the apex. The ovipositor is from three-fourths to as long as the abdomen; the piercing part slightly longer than the sheath, which is black, edged with scattered bristles; apex of saw with 2 strong teeth (fig. 2, i).

Wings (fig. 2, f, g).—Forewing, length, 2.5 mm.; breadth, 1.1 mm. Hyaline, with a slight cloud below the upturning of the submarginal vein and in a line from the frenulum to the base. There are three faint venae spuriae from the end of the radius and two others from the base of the wing. The submarginal, marginal, postmarginal and radius are in the proportion:—8:3:5:2. Hindwing, length, 1.35 mm.; breadth, .25 mm., with cilia nearly half as long as the breadth; remarkably long in proportion to the breadth, with rounded apex. Hooks of retinaculum on a distinct elevation. Submarginal vein broad basally, but tapering off before the hooks at about three-fourths. Colour clear, slightly clouded at the end of the single vein.

Fore legs.—Coxae compressed, $2\frac{1}{2}$ times as long as broad; 6–8 bristles along anterior edge, chiefly on anterior two-thirds; bare externally, save on the upper or basal posterior third, where are 6–9 short spines; on the posterior edge and along the adjacent inner aspect, especially on the basal third, are numerous soft hairs; on the inside also are 3 longer apical bristles. Trochanter short, with 4–5 ventral bristles. Femur one-third longer than coxa, narrow on basal half, then expanded; greatest width one-fourth the length; on the outer surface are about a dozen bristles, which increase in length towards the apex; on the apical ventral region are numerous scattered hairs, besides a few on the anterior edge and a regular row (short) on or near the ventral. Tibia one-third of the femur in length; $2\frac{1}{2}$ times as long as broad, with 2 short stout outer apical spines, one dorsal, the other ventral; many strong scattered bristles on the outer aspect and a few on the inner; the pre-apical ventral bristle not stoutly developed. Tarsus nearly thrice

Waterston, James. 1914. "Notes on african Chalcidoidea.—I." *Bulletin of entomological research* 5, 249–258.

<https://doi.org/10.1017/S0007485300031862>.

View This Item Online: <https://www.biodiversitylibrary.org/item/111123>

DOI: <https://doi.org/10.1017/S0007485300031862>

Permalink: <https://www.biodiversitylibrary.org/partpdf/49643>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.