Genus OCHLEROTATUS, Arrib., Edw.

O. hirsutus, Theo.

Theobald (Union of S. Afr., Second Rept. Vet. Research, p. 328) has criticised my statement that his *Culex hirsutum* and *C. transvaalensis* are one and the same species. I have re-examined the specimens, and find no reason to modify my previous opinion, except in one particular. The male and female types of *C. hirsutum* belong to different species, and it was perhaps owing to his confusion of the two that Theobald was unable to agree with my findings. The female type must be taken as the type of the species, the male (the specimen figured in Bull. Ent. Res. ii, p. 249) being really *O. nigeriensis* (Theo.). These species, as previously pointed out, can be readily distinguished by the colour of the wing-scales, but for their better definition, I now give figures of the male genital claspers of each (fig. 1, *d* and *e*).

Unrecorded localities for *O. hirsutus* are: BRITISH EAST AFRICA: Moyale, 1. v. 1913 (Dr. G. R. H. Chell); PORTUGUESE WEST AFRICA: Bihé (Dr. C. Wellman).

O. cheili, sp. nov.

♀. Head clothed with yellowish-ochreous narrow curved and upright forked scales, only a few flat ones at the sides. Basal joint of antennae with small flat ochreous scales. Palpi and proboscis entirely black-scaled. Thorax with the integument dark brown, clothed with light ochreous brown scales at the sides, darker brown ones in the middle. Scutellum with narrow pale ochreous scales. Abdomen marked somewhat as in *O. dorsalis*: it is dark brown, with ochreous basal bands on each segment and with a continuous median longitudinal stripe of the same colour; towards the apical margins of the fifth, sixth and seventh segments this stripe broadens out considerably. Scales of venter ochreous. Legs: femora and tibiae strongly mottled with dark brown and pale ochreous scales, the femora entirely pale on their inner or posterior surfaces, except towards the tip. Tarsi of all the legs dark brown, the first joint with a fairly broad but ill-defined pale ring at the base, second and third joints with very narrow basal pale rings, fourth joint with a mere trace of such a ring; fifth entirely dark. Claws toothed (hind pairs missing). Wings with dark brown scales, and a few scattered pale ones on the median series; scales of lateral series linear. Upper fork-cell with its base nearer the apex of the wing than the lower.

Length about 5 mm.

BRITISH EAST AFRICA: Dido, 30. xi. 1911 (Dr. G. R. H. Chell), 3♀.

These specimens answer in many respects to Neveu-Lemaire's description of *Taeniorhynchus africanus*, but there are several discrepancies, and on the whole it seems most probable that Neveu-Lemaire's species was only *O. dorsalis*, some disagreement notwithstanding. The present species is named in honour of the collector, the only person who has up to the present collected mosquitos in this interesting district of British East Africa.

O. bevisi, sp. nov.

♀. Head clothed mainly with pale ochreous scales, but the usual spot of dark brown ones is present on each side. Scales narrow and curved in the middle and round the eyes, broad and flat at the sides, the latter reaching not far short of the

middle line. Palpi blackish brown, with creamy tips. Proboscis dark brown at the base and apex, ochreous in the middle two-thirds beneath and at the sides. First two joints of antennae with small flat ochreous scales, the basal joint ochreous. *Thorax* dark brown, clothed with narrow brown and ochreous-brown scales without any definite pattern. Scutellar scales narrow. Flat ochreous scales on the pleurae. *Abdomen* dark brown, each segment with whitish basal lateral spots and median bands not connected with the spots. Venter whitish ochreous, the apical margins of the apical segments black. *Legs*: femora without any intersprinkling of light and dark scales, pale knee-spots very distinct. Tibiae blackish, on the front and middle legs largely pale on their outer and posterior faces; hind tibiae with a well-marked pale spot at the apex on the outside, which is about equal in length to the breadth of the tibia. Tarsi blackish; on all the legs there is a narrow ochreous ring embracing the tip of the first and the base of the second joint, and a still narrower ring embracing the tip of the second and the base of the third. All the claws toothed. *Wings* clothed with dark brown scales, those of the lateral series linear. Bases of fork-cells level.

Length about 5 mm.

NATAL: Umbilo, Durban; 1 ♀, 16. v. 1914 (type); 2 ♀, 13. ix. 14 and 1 ♀, 25. ix. 1914 (*L. Bevis*).

Readily distinguished from all other African species by the tarsal markings. Although small pale rings are present on these, the species seems to belong to the *dentatus*-group.

***O. albocephalus*, Theo.**

A good series of specimens of both sexes of this species have been received from Durban (*E. C. Chubb* and *L. Bevis*). Previously it was only known in the male sex from West Africa. It proves to be of considerable interest owing to a marked difference of scale characters between the two sexes. In the male, the flat scales of the head extend up to the middle line in front, while in the female the middle area of the head is clothed with narrow scales; the scutellar scales in the male are broad, in the female narrow. No such sexual difference has been noticed in any other mosquito.

As I have previously pointed out (*Bull. Ent. Res.* iii, p. 21), the figure which Theobald gives (*Mon. Cul.* v, p. 206) as representing the male genitalia of *O. punctothoracis*, really depicts those of *O. albocephalus*. This figure, however, is by no means accurate, and a fresh one is therefore given herewith (fig. 1, c).

The female *O. albocephalus* can be distinguished from *O. quasiunivittatus*, which it otherwise closely resembles, by the much larger white spot at the apex of the hind tibiae, which is quite twice as long as the width of the tibia.

***O. quasiunivittatus*, Theo.**

This species occurs side by side with *O. albocephalus* in Durban, a few females having been sent in by Mr. Chubb. What appears to be a good specific character, is the size of the pale spot on the hind tibia; this is very distinct, and is about equal in length to the diameter of the apex of the tibia, *i.e.* it is about as broad as long. The thoracic scaling is like that of the female *O. albocephalus*. The fact that this

species and *O. dentatus* differ markedly in the male genitalia has already been referred to, and the present opportunity is taken of figuring this difference (figs. 1, *a*, *b*). There are other well-marked genitalic distinctions between *O. quasiunivittatus* and *O. dentatus* besides those in the claspers, but as the latter are most readily seen (being usually visible even in a dried specimen), they alone have been figured.

Fig. 1. Male claspers of :—(*a*) *Ochlerotatus quasiunivittatus*, Theo. ; (*b*) *O. dentatus*, Theo. ; (*c*) *O. albocephalus*, Theo. ; (*d*) *O. nigeriensis*, Theo. ; (*e*) *O. hirsutus*, Theo.

***O. dentatus*, Theo.**

This species also apparently occurs at Durban together with *O. albocephalus* and *O. quasiunivittatus*, though only female specimens have been received. It seems to differ from the other two in the thorax being darker, the pale scales having a tendency to a linear arrangement, and in the very small pale spot at the tip of the hind tibiae, which is never more than about half as long as broad. The best specific character is, however, the structure of the male genitalia (fig. 1, *b*). The species has been received from various localities in British East Africa.

***O. pembaensis*, Theo.**

Aedes pembaensis, Theo., Mon. Cul. ii, p. 235 (1901).

Skusea pembaensis, Theo., Mon. Cul. iii, p. 291 (1903).

Verrallina? *pembaensis*, Theo., Mon. Cul. v, p. 495 (1910).

Howardina? *pembaensis*, Edw., Bull. Ent. Res. iii, p. 13 (1912).

Aedes (*Skusea*) *pembaensis*, Edw., Bull. Ent. Res. iv, p. 49 (1913).

As indicated above, this species has undergone many nomenclatorial vicissitudes, and it is not certain that it has even yet found a final resting-place, as it is one of those intermediate forms which make the classification of the *Aedes* group so difficult. Specimens representing both sexes have lately been sent in by Dr. W. M. Aders from Zanzibar, where the larvae were found living in stagnant water, 31. iii. 1914. The male palpi are as long as the proboscis, and though the last two joints are scarcely swollen, they are hairy and bent downwards as usual in *Ochlerotatus*, so that in spite of the simple claws of the female the species is probably better placed

1915. "Diagnoses of new bornean Culicidae." *Bulletin of entomological research* 5, 283–285. <https://doi.org/10.1017/S0007485300030315>.

View This Item Online: <https://www.biodiversitylibrary.org/item/111123>

DOI: <https://doi.org/10.1017/S0007485300030315>

Permalink: <https://www.biodiversitylibrary.org/partpdf/49646>

Holding Institution

Smithsonian Libraries

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.