VII.—On a supposed new Species of *Rhiostoma* from Borneo, and Notices of Two other Species of Shells from Palawan.
By Lieut.-Col. H. H. GODWIN-AUSTEN, F.R.S. &c.

THE *Rhiostoma* I now describe was collected by Mr. Boxall and was sent by him to Dr. Hungerford, who asked me to describe and name it after its discoverer; but I received it too late to bring into my paper on Bornean shells published in the Proc. Zool. Soc., June 1889, p. 332.

Rhiostoma Boxalli, sp. n. (Fig. 1, *a*, *b*, *c*.)

Loc. Near Kina Balu (*Boxall*). In Coll. Dr. Hungerford.

Shell discoid, widely and perspectively umbilicated; sculpture smooth; colour dark madder-brown, with a few white blotches on the second and third whorls; spire scarcely raised above the last whorl, apex rounded; suture deep; whorls 4, round, rapidly increasing, the last separated for a

Fig. 1.

Rhiostoma Boxalli.

a. Specimen from Kina Balu; *b*, *c.* Ditto from Palawan.

short distance from the penultimate; the sutural tube short, horizontal; aperture oblique, circular; peristome double, the inner simple, the outer expanded at right angles to the body-whorl, and curved slightly backwards on the outer margin and with a slight nick near the sutural tube.

Size: maj. diam. 24.0, min. diam. 16.25, alt. axis 5.75 mm.

This species comes nearest to *R. iris*, G.-A. (described in the Proc. Zool. Soc., June 1889, p. 343), in the expanded peristome; the general form differs, it is not so turbinate or streaked with zigzag markings.

One specimen of this species occurred in a collection of land-shells sent to me by Mr. John Whitehead from the island of Palawan.

Lagocheilus similis, var., Edgar Smith. (Fig. 2, *a*, *b*.)

Among the shells collected by Mr. John Whitehead in Palawan and sent to me there are three specimens of what I

Fig. 2.

Lagocheilus similis, var., Edgar Smith.

take to be the above species, only much better grown. When in perfect preservation it is covered with a brown epidermis having well-displayed transverse, regular, fine liration and the periphery adorned with a strong hairy fringe.

The largest specimen measures:—major diam. 18·75, minor 15·0, alt. axis 7·5 mm.

Cassidula bicolor, sp. n. (Fig. 3, *a*, *b*.)

Palawan. Collected by Mr. John Whitehead.

This is a smaller shell than *sulculosa*, Mouss., to which it is nearest as regards the form of the aperture, but it is not so strongly formed. The body-whorl is crossed by shallow

Fig. 3.

Cassidula bicolor.

longitudinal plications. The colour is a very dark sepia-brown, with two broad, parallel, very white bands, which extend to the outer margin of the peristome. I could find nothing like it in the British Museum collection.

Godwin-Austen, Henry Haversham. 1893. "VII.—On a supposed new species of *Rhiostoma* from Borneo, and notices of two other species of shells from Palawan." *The Annals and magazine of natural history; zoology, botany, and geology* 12, 32–33. <https://doi.org/10.1080/00222939308677570>.

View This Item Online: <https://www.biodiversitylibrary.org/item/53308>

DOI: <https://doi.org/10.1080/00222939308677570>

Permalink: <https://www.biodiversitylibrary.org/partpdf/51213>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.