

PROCEEDINGS  
OF THE  
CALIFORNIA ACADEMY OF SCIENCES

Vol. 50, No. 3, pp. 39–114.

December 9, 1997

SPECIES CATALOG OF THE NEUROPTERA, MEGALOPTERA,  
AND RAPHIDIOPTERA OF AMERICA NORTH OF MEXICO

By

Norman D. Penny

Department of Entomology, California Academy of Sciences,  
San Francisco, CA 94118

DEC 18 1997

Woods Hole, MA 02543

Phillip A. Adams

California State University, Fullerton, CA 92634

and

Lionel A. Stange

Florida Department of Agriculture, Gainesville, FL 32602

Woods Hole Biological Laboratory /  
Oceanographic Institution  
Library

The 399 currently recognized valid species of the orders Neuroptera, Megaloptera, and Raphidioptera that are known to occur in America north of Mexico are listed and full synonymies given. Geographical distributions are listed by states and provinces. Complete bibliographic references are given for all names and nomenclatural acts. Included are two new junior homonyms indicated, seven new taxonomic combinations, two new changes of rank, fourteen new synonymies, three new lectotype designations, and one new name.

Received March 20, 1996. Accepted June 3, 1997.

The recent publication of *Nomina Insecta Nearctica, A Check List of the Insects of North America* (Poole 1996) has given us a listing of North American Neuropterida (Neuroptera + Megaloptera + Raphidioptera) species for the first time in more than a century. However, for anyone trying to identify these species, the literature is scattered and obscure. Only with the greatest difficulty is a non-specialist able to keep up with current usage of names. With this lack of reference materials in mind, the present catalog has been developed to allow the non-specialist and specialist alike to grasp the limits and diversity of these groups. The primary literature has

been consulted whenever possible, as well as *Zoological Record*, and appropriate monographic revisions published up to 1 January 1997.

A number of taxonomic changes are incorporated into this catalog: there are two new junior homonyms indicated, seven new taxonomic combinations, two new changes of rank, fourteen new synonymies, three new lectotype designations, and one new name. However, the classification of neuropteroid insects is constantly being modified and enhanced as we understand the group better. Thus, the authors know that the genus "Mantispa" is currently being redefined by Kevin Hoffman, and the genus "Brachynemu-

"rus" was recently redefined by Stange (1994). Such studies enhance our knowledge of these groups and are a normal part of taxonomic development and maturity, but at the same time they require that this list, as with any catalog, be used in conjunction with other recent taxonomic literature to ensure the most up-to-date treatment of any particular taxon. To help maintain currency, we have listed most recent treatments of included taxa at the end of each family.

The geographical coverage of this catalog includes continental United States (including Alaska with its offshore islands), and Canada. Greenland has been excluded. The southern limits are the political boundaries of the United States, excluding the Caribbean and all offshore islands, such as the Bahamas, not belonging to the United States. The Hawaiian Islands have been excluded because they belong to another faunal region, as well as having a current, on-line faunal listing through the Hawaiian Biological Survey URL address (<http://www.bishop.hawaii.org/bishop/HBS/>).

This portion of the Nearctic Region, excluding Mexico, currently contains 399 valid species, distributed in the following families:

Family	Genera	Species
<b>Neuroptera</b>		
Ascalaphidae	3	8
Berothidae	1	10
Chrysopidae	15	81
Coniopterygidae	8	55
Dilaridae	1	2
Hemerobiidae	6	61
Ithonidae	1	1
Mantispidae	4	15
Myrmeleontidae	17	94
Polystoechotidae	2	2
Sisyridae	2	6
<b>Raphidioptera</b>		
Inocellidae	1	3
Raphidiidae	2	18
<b>Megaloptera</b>		
Corydalidae	7	19
Sialidae	1	24
Total	71	399

Distributional information is credited to an author only if a state or province is specifically mentioned in the referenced work. Thus, distributional records associated with some species

are sparse, even though earlier authors may have mentioned such regions as "Atlantic Coastal States" or "Maritime Provinces." New distributional data introduced in this catalog are credited to the contributing author.

Some of the published records must be viewed with caution, particularly among older literature. Where we know records are wrong, such as with *Chrysoperla externa* in Canada, we have indicated this. However, other records which may have been correct are listed without comments under each species treatment.

Type label information has come from three sources. Where possible, information has been transcribed from the type specimens. Much label information has also been cited from the original literature. A valuable third source has been the electronic type catalog at Harvard University's Museum of Comparative Zoology (gopher: //huh.harvard.edu:179/7), which houses the extensive Banks collection.

There are a surprisingly large number of North American types in the collection at the Museum of Comparative Zoology, Harvard University. In addition to the expected types of Hagen, Banks, and Carpenter, there are also primary types described by Schneider, Burmeister, Smith and Navás, among others. Information about the types at this museum are now available as a listing on internet.

Within the main body of the catalog the orders are arranged: Neuroptera, Raphidioptera, and finally Megaloptera. Families are arranged alphabetically. The included species are arranged within a taxonomic infrastructure of subfamilies and tribes in the larger families, with included genera and species organized alphabetically. Each valid species includes in sequential order: reference to the original description; original generic placement; label data of type specimens and type depository; synonyms, their label data and type depository; taxonomic status changes with appropriate references; and finally geographical distribution by state or province within the defined area. Acronyms for states/provinces are alphabetized and followed by a superscript number that keys to the citation at the end of the entry, e.g., AZ<sup>6</sup> (Say, 1824:305).

Generic synonymies were published by Oswald and Penny (1991) and are not repeated here.

## ABBREVIATIONS

The state and province abbreviations used throughout this paper are the two letter codes used by the U. S. Postal Service for the United States and Canada. Collection abbreviations are as follows:

AMNH = American Museum of Natural History, New York, NY

ANSP = Academy of Natural Sciences, Philadelphia, PA

BMNH = The Natural History Museum, London, England

CAS = California Academy of Sciences, San Francisco, CA

CIC = Cawthon Institute Collection, New Zealand

CMNH = Carnegie Museum of Natural History, Pittsburgh, PA

CNC = Canadian National Collection, Ottawa, Canada

CORN = Cornell University, Ithaca, NY

DEIC = Institut für Pflanzenschutzforschung, Eberswalde, Germany

HMB = Museum für Naturkunde der Humboldt Universität, Berlin, Germany

INHS = Illinois State Natural History Survey, Champaign, IL

IRSNB = Institut Royal des Sciences Naturelles de Belgique, Brussels, Belgium

LACM = Los Angeles County Natural History Museum, Los Angeles, CA

MACN = Museo Argentino de Ciencias Naturales, Buenos Aires, Argentina

MCZ = Museum of Comparative Zoology, Harvard University, Cambridge, MA

MLP = Museo de La Plata, La Plata, Argentina

MNHN = Muséum National d'Histoire Naturelle, Paris, France

MZB = Museu de Zoologia, Barcelona, Spain

MZH = Universitetets Zoologiska Museum, Helsingfors, Finland

NMV = National Museum of Victoria, Melbourne, Australia

NRS = Naturhistoriska Riksmuseet, Stockholm, Sweden

OHSTU = Ohio State University Insect Collection, Columbus, OH

PSU = Pennsylvania State University Insect Collection, State College, PA

ROM = Royal Ontario Museum, Ontario, Canada

SDNHM = San Diego Natural History Museum, San Diego, CA

SEM = Snow Entomological Museum, Lawrence, KS

USNM = National Museum of Natural History, Smithsonian Institution, Washington, DC

UCB = University of California, Berkeley, CA

UCD = University of California, Davis, CA

YPM = Yale University Peabody Museum, New Haven, CT

ZIL = Zoological Institute, Lund, Sweden

ZMC = Zoologiske Museum, Copenhagen, Denmark

ZMH = Zöologisches Museum, Hamburg, Germany

ZMUS = Zoological Museum, University of Oslo, Norway

ZSBS = Zoologische Sammlungen des Bayerischen Staates, Munich, Germany

## ASCALAPHIDAE

Ascalaphidae, or owlflies, as adults are large crepuscular aerial predators. For some species the daily flight period is very short, less than a half hour before total darkness in the evening (MacNeill 1962). Eggs are laid on the tips of twigs (Henry 1977), often near streams. The larvae live on the soil surface or on leaves, where they await passing prey. Adults are strong fliers and superficially resemble dragonflies, except for their distinctive long antennae. Males and females are often dimorphic in wing coloration, wing shape, and even sometimes antennal length. The world fauna was monographed by Weele (1909a). Shetlar's (1977) unpublished Ph.D. thesis is the most recent treatment of North American species. Only three genera are known from North America.

## HAPLOGLENIINAE

*Ascalobyas* Penny, 1982:395

**Ascalobyas albistigma** (Walker), 1853:452 [*Ascalaphus*].

Holotype (sex unknown): Honduras (BMNH).

TAXONOMY.—To *Haploglenius* (see McLachlan, 1871:236); to *Byas* (see Weele, 1908:30); to *Ascalobyas* (see Penny, 1982:395).

DISTRIBUTION.—USA: TX (Vogtsberger, 1990:158).

**Ascaloptynx** Banks, 1915:350

**Ascaloptynx appendiculatus** (Fabricius), 1793:96 [*Ascalaphus*].

Holotype female: USA, "Carolinas" (BMNH). = *Ptynx juvenilis* McLachlan, 1871:239. Type(s) (sex unknown): USA, Texas, Belfrage (BMNH). **NEW SYNONYMY.**

= *Ptynx furciger* McLachlan, 1891:509. Five syntypes (3 males, 2 females): USA, Arizona, H. K. Morrison (BMNH). **NEW SYNONYMY.**

TAXONOMY.—To *Haploglenius* (see Rambur, 1842:363); to *Ptynx* (see McLachlan, 1871:239); to *Neuroptynx* (see Banks, 1907c:32); to *Ascaloptynx* (see Banks, 1915a:350); (*P. juvenilis* = *A. appendiculatus*, *P. furciger* = *A. appendiculatus* (see Shetlar, 1977:99)).

DISTRIBUTION.—USA: AL<sup>5</sup>, AR<sup>5</sup>, AZ<sup>2</sup>, FL<sup>3</sup>, GA<sup>3</sup>, MO<sup>4</sup>, MS<sup>6</sup>, NC<sup>3</sup>, OK<sup>5</sup>, SC<sup>5</sup>, TN<sup>5</sup>, TX<sup>1</sup>, VA<sup>5</sup> (<sup>1</sup>McLachlan, 1871:239, as *P. juvenilis*; <sup>2</sup>McLachlan, 1891:509; <sup>3</sup>Weele, 1908:58, 59; <sup>4</sup>Froeschner, 1947:129; <sup>5</sup>Shetlar, 1977:102; <sup>6</sup>Lago & Testa, 1989:12).

REMARKS.—The synonymy of *Ptynx juvenilis* and *Ptynx furciger* with *Ascaloptynx appendiculatus* was first suggested by Shetlar in 1977. However, his Ph.D. thesis was never published and therefore cannot be used for nomenclatorial purposes. The present publication is the first official record of these synonymies.

ASCALAPHINAE

**Ululodes** Currie in Smith, [1900] 1899:57

**Ululodes arizonensis** Banks, 1907a:275. Three syntypes (sex unknown): USA, Arizona, Palmerlee, IX, C. R. Biederman; same data, except VII (two specimens) (MCZ). **NEW STATUS.**

TAXONOMY.—To *U. quadrimaculata arizonensis* (see Weele, 1908:110); to *U. arizonensis* (see Shetlar, 1977:142).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NM<sup>2</sup>, NV<sup>3</sup>, TX<sup>2</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1907a:275; <sup>2</sup>Shetlar, 1977:147, 148); <sup>3</sup>Penny, **NEW STATE RECORDS**).

REMARKS.—The elevation of *Ululodes arizonensis* to species rank was first suggested by Shetlar in 1977. However, his Ph.D. thesis was never published and therefore cannot be used for nomenclatorial purposes. The present publication is the first official record of this status change.

**Ululodes bicolor** (Banks), 1895a:521 [*Ulula*].

Three syntypes (sex unknown): Mexico, Baja California Sur, San José del Cabo; Sierra El Taste, IX (3 specimens) (MCZ).

TAXONOMY.—To *Ululodes* (see Weele, 1908:105).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>1</sup>, LA<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Shetlar, 1977:138; <sup>2</sup>Penny, **NEW STATE RECORDS**).

**Ululodes floridana** (Banks), 1906a:99 [in *Suhpalacsas* as "Suhpalasca"]

Holotype male: USA, southern Florida (MCZ). **REVISED STATUS.**

TAXONOMY.—To subspecies of *Ululodes quadrimaculata* (see Weele, 1908:111); to species status (see Shetlar, 1977:139).

REMARKS.—The elevation of *Ululodes floridana* to species rank was first suggested by Shetlar in 1977. However, his Ph.D. thesis was never published and therefore cannot be used for nomenclatorial purposes. The present publication is the first official record of this status change.

DISTRIBUTION.—USA: FL<sup>1</sup>, GA<sup>2</sup>, SC<sup>2</sup> (<sup>1</sup>Banks, 1906a:99; <sup>2</sup>Shetlar, 1977:141).

**Ululodes macleayana** (Guilding), 1825:140 [*Ascalaphus*].

Holotype male: Antilles, Saint-Vincent, VI.1824, Guilding (BMNH).

= *Ascalaphus senex* Burmeister, 1839:1001. Syntype(s) unknown: locality unknown (collection unknown).

= *Ulula microcephala* Rambur, 1842:359.

= *Ascalaphus avunculus* Hagen, 1861:238.

Syntype(s) unknown: Cuba, Poey (USNM).

= *Ululodes macleayana hageni* Weele, 1908:101.

TAXONOMY.—To *Ulula* (see McLachlan, 1871:247); to *Ululodes* as synonym of *U. hyalina* (see Banks, 1907c:32); as valid species (see Weele, 1908:101).

DISTRIBUTION.—USA: AL<sup>4</sup>, AR<sup>4</sup>, FL<sup>1</sup>, GA<sup>4</sup>, KS<sup>2</sup>, LA<sup>4</sup>, MO<sup>3</sup>, MS<sup>4</sup>, NC<sup>5</sup>, NJ<sup>1</sup>, OK<sup>4</sup>, SC<sup>4</sup>, TX<sup>1</sup> (<sup>1</sup>Weele, 1908:99 as subspecies *U. m. hageni*; <sup>2</sup>Smith, 1925:170; <sup>3</sup>Froeschner, 1947:129;

<sup>4</sup>Shetlar, 1977:116–120; <sup>5</sup>Penny, NEW STATE RECORD).

**Ululodes nigripes** Banks, 1943:79.

Holotype (sex unknown): USA, Texas, Davis Mountains, D. J. & J. N. Knoll (OHSTU).

DISTRIBUTION.—USA: TX (Banks, 1943:79).

**Ululodes quadripunctatus** (Burmeister), 1839:1001 [Ascalaphus]. NEW STATUS

Type(s) (female): USA, New York (MCZ).

= *Ascalaphus quadrimaculata* (Say), 1824:305. A junior primary homonym of *Ascalaphus quadrimaculatus* Lichtenstein, 1796:192. Holotype female: USA, Pennsylvania (MCZ).

= *Colobopterus excisus* Hagen, 1887:193. Holotype male: USA, Florida, 1862, Uhler (MCZ).

= *Ulula albifrons* Banks, 1901c:172. Two syntype females: USA, Arizona, Phoenix, 20.IX, Kunze (MCZ).

TAXONOMY.—To *Ulula* (see McLachlan, 1871:247); to *Ululodes* (see Currie, 1900:57); (*A. quadripunctatus* = *U. quadrimaculata*; *U. albifrons* = *U. quadrimaculata albifrons*) (see Weele, 1908:109); (*C. excisus* = *U. quadrimaculata*) (see Beatty & Runner, 1970:153).

DISTRIBUTION.—CAN: ON<sup>13</sup>. USA: AL<sup>12</sup>, AR<sup>14</sup>, AZ<sup>6</sup>, DC<sup>4</sup>, DE<sup>12</sup>, FL<sup>3</sup>, IA<sup>12</sup>, IL<sup>12</sup>, IN<sup>8</sup>, KS<sup>7</sup>, KY<sup>12</sup>, LA<sup>12</sup>, MD<sup>2</sup>, MI<sup>12</sup>, MO<sup>10</sup>, MS<sup>12</sup>, NC<sup>9</sup>, NE<sup>14</sup>, NJ<sup>5</sup>, NY<sup>2</sup>, OH<sup>12</sup>, OK<sup>12</sup>, PA<sup>1</sup>, SC<sup>12</sup>, TN<sup>12</sup>, TX<sup>12</sup>, VA<sup>12</sup>, WS<sup>11</sup>, WV<sup>14</sup> (Say, 1824:305; <sup>2</sup>Hagen, 1861:239, as *U. quadripunctata*; <sup>3</sup>Hagen, 1887:193, as *C. excisus*; <sup>4</sup>Banks, 1892:361; <sup>5</sup>Smith 1900:57; <sup>6</sup>Banks, 1901c:172; <sup>7</sup>Smith, 1925:170, as *C. excisus*; <sup>8</sup>Montgomery & Trippel, 1933:260, as *C. excisus*; <sup>9</sup>Brimley, 1938:32; <sup>10</sup>Froeschner, 1947:129; <sup>11</sup>Throne, 1972:121; <sup>12</sup>Shetlar, 1977:124–127; <sup>13</sup>Garland & Marshall, 1980:637; <sup>14</sup>Penny, NEW STATE RECORDS).

### BEROTHIDAE

Berothidae, or beaded lacewings, are so called because of the development of secretion encrusted setae on wings of females. Adults of most North American species are immediately recognizable by the falcate shape of the forewing. Females lay stalked eggs on wood surfaces, often near termite nests. Larvae are inquilines in dry-wood termite nests, where they feed by injecting a paralyzing chemical into the termite, and then sucking out the body fluids of the immobile prey.

The only genus of berothids known from North America is *Lomamyia* Banks 1904. The species were reviewed and a key provided by Carpenter (1940). Several additional undescribed species are known from North America.

### **Lomamyia** Banks, 1904c:209

**Lomamyia banksi** Carpenter, 1940:260.

Holotype male: USA, South Carolina, Clemson College, 30.IV.1932, D. Duncan (MCZ, specimen not found).

DISTRIBUTION.—USA: AL<sup>1</sup>, AZ<sup>1</sup>, DC<sup>1</sup>, FL<sup>1</sup>, IL<sup>1</sup>, IN<sup>2</sup>, KS<sup>1</sup>, LA<sup>3</sup>, MI<sup>1</sup>, MO<sup>1</sup>, MS<sup>2</sup>, NC<sup>1</sup>, NY<sup>1</sup>, SC<sup>1</sup>, VA<sup>1</sup>, (Carpenter, 1940; <sup>2</sup>Lawson & McCafferty, 1984:130; <sup>3</sup>Penny, NEW STATE RECORD).

**Lomamyia flavigornis** (Walker), 1853:278 [*Hermerobius*].

Lectotype (without abdomen): North America (BMNH).

= *Isoscelipteron pennsylvanicum* Brauer, 1864:898. Type(s) sex(es) unknown: USA, Pennsylvania (Vienna).

= *Lomamyia nearctica* Navás, 1913b:19. Holotype male: USA, New York, Japhank [Long Island], 23.IX.1911, De la Torre Bueno (MZB).

TAXONOMY.—To *Micromus* (see Hagen, 1861:198); to *Lomamyia* (see Banks, 1905a:27); (*L. nearctica* = *L. flavigornis*; *I. pennsylvanicum* = *L. flavigornis*) (see Navás, 1929a:26); lectotype designated (see Carpenter, 1940:260).

DISTRIBUTION.—USA: AL<sup>8</sup>, DC<sup>2</sup>, FL<sup>8</sup>, GA<sup>1</sup>, IN<sup>9</sup>, KS<sup>4</sup>, KY<sup>8</sup>, LA<sup>8</sup>, MS<sup>10</sup>, NC<sup>7</sup>, NJ<sup>3</sup>, NY<sup>5</sup>, PA<sup>8</sup>, VA<sup>6</sup> (Walker, 1853:278; <sup>2</sup>Hagen, 1861:199; <sup>3</sup>Smith, 1900:56; <sup>4</sup>Smith, 1925:167; <sup>5</sup>Leonard, 1928:40; <sup>6</sup>Navás, 1929:26; <sup>7</sup>Brimley, 1938:30; <sup>8</sup>Carpenter, 1940:260; <sup>9</sup>Lawson & McCafferty, 1984:130; <sup>10</sup>Penny, NEW STATE RECORD).

**Lomamyia fulva** Carpenter, 1940:264.

Holotype female: USA, California, Riverside Co., San Jacinto Mts., 21.VII.1929, R. H. Beamer (MCZ, specimen not found).

DISTRIBUTION.—USA: CA (Carpenter, 1940:264).

**Lomamyia hamata** (Walker), 1853:278 [*Hermerobius*].

Holotype male: North America (BMNH).

= *Lomamyia hubbardi* Banks, 1924:430. Type(s) sex(es) unknown: USA, Florida, Cedar Keys, 5.IV.1878, Hubbard and Schwarz (MCZ).

**TAXONOMY.**—To *Micromus* (see Hagen, 1861:199); to *Lomamyia* (see Banks, 1905a:48); (*H. hamatus* = *L. flavigornis*) (see Navás, 1929a:26); (*L. hamatus* not equal to *L. flavigornis*, *L. hubbardi* = *L. hamata*) (see Carpenter, 1940:265).

**DISTRIBUTION.**—**USA:** FL<sup>1</sup>, SC<sup>2</sup> (<sup>1</sup>Carpenter, 1940:265; <sup>2</sup>Brushwein, 1987a:671).

**Lomamyia latipennis** Carpenter, 1940:262.

Holotype male: USA, California, Marin Co., Phoenix Lake, 4.VII.1927, H. H. Kiefer (CAS).

**DISTRIBUTION.**—AZ, CA (Carpenter, 1940:262).

**Lomamyia longicollis** (Walker), 1853:281 [*Hermetobius*].

Two syntype males: USA, Georgia, Abbot (BMNH).

**TAXONOMY.**—To *Lomamyia* (see Carpenter, 1940:267).

**DISTRIBUTION.**—GA<sup>1</sup>, MS<sup>2</sup>, NC<sup>2</sup>, SC<sup>3</sup> (<sup>1</sup>Walker, 1853:281; <sup>2</sup>Carpenter, 1940:267; <sup>3</sup>Brushwein, 1987b:151).

**Lomamyia occidentalis** (Banks), 1905a:89 [*Berotha*].

Lectotype female: USA, Nevada, Ormsby Co., 6.VII, Baker (MCZ).

**TAXONOMY.**—To *Lomamyia* (see Banks, 1905c:48); considered a synonym of *L. texana* (see Banks, 1907c:23); considered a valid species (see Carpenter, 1940:263); lectotype designated (see Carpenter, 1940:263).

**DISTRIBUTION.**—**CAN:** BC<sup>2</sup>, **USA:** AZ<sup>1</sup>, CA<sup>2</sup>, NV<sup>1</sup> (<sup>1</sup>Banks, 1905a:89; <sup>2</sup>Carpenter, 1940:263).

**Lomamyia squamosa** Carpenter, 1940:266.

Holotype male: USA, Texas, Brownsville, 1116.VI.1933, P. J. Darlington, Jr. (MCZ).

**DISTRIBUTION.**—**USA:** TX (Carpenter, 1940:266).

**Lomamyia tenuis** Carpenter, 1940:261.

Holotype male: USA, New Mexico, Torrance Co., VII.1925, C. H. Martin (MCZ).

**DISTRIBUTION.**—**USA:** NM (Carpenter, 1940:262).

**Lomamyia texana** (Banks), 1897:24 [*Berotha*].

Lectotype male: USA, Texas (MCZ).

**TAXONOMY.**—To *Lomamyia* (see Banks, 1905a:27); lectotype designated (see Carpenter, 1940:265).

**DISTRIBUTION.**—**USA:** AZ<sup>2</sup>, TX<sup>1</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1897:24; <sup>2</sup>Carpenter, 1940:265).

## CHrysopidae

Most chrysopid adults are green with golden eyes, but members of the genera *Nothochrysa*, *Pimachrysa*, *Eremochrysa*, and *Yumachrysa* are usually predominately brown or black. Most species have characteristic red or dark markings, which often fade within a few months after death. Eggs are laid on long stalks, usually singly. Larvae are voracious predators, those of some species carrying packets of trash or corpses of their prey on their backs (Smith 1922). Most species will feed upon a wide variety of slow-moving insects, although some have more specific food requirements. Larvae of Belonopterygini (*Abachrysa* and *Nacarina* in our region) live in ant nests, presumably feeding upon the ants. Adult chrysopids are very conservative morphologically, with many of the most taxonomically significant characteristics found in the male terminalic structures. Thus, to determine the generic placement of an unfamiliar male specimen, the abdomen must be cleared (in 10% KOH), and subsequently may be stained with chlorazol black-E. A key to subfamilies and genera of Nothochrysinae was provided by Adams (1967). A worldwide generic revision of the family contains helpful descriptions and illustrations of wings and genitalic structures, as well as a key to genera (Brooks and Barnard 1990). The species of Chrysopidae of Canada were included in a key by Garland (1985). All other literature is too old to be of much help with identification. There are 14 genera of Chrysopidae known from North America; several new species are in collections, awaiting description.

## NOTHOCHRYSINAЕ

**Nothochrysa** McLachlan, 1868:195

**Nothochrysa californica** Banks, 1892:373.

Holotype male: USA, California, Los Angeles (MCZ).

**DISTRIBUTION.**—**CAN:** BC<sup>2</sup>, **USA:** CA<sup>1</sup>, OR<sup>3</sup>, WA<sup>3</sup> (<sup>1</sup>Banks, 1892:373; <sup>2</sup>Smith, 1932:582; <sup>3</sup>Adams, 1967:224).

**Pimachrysa** Adams, 1957a:67

**Pimachrysa albicostales** Adams, 1967:226.

Holotype male: Mexico, Baja California, 22 miles north of Punta Prieta, 9.XII.1958, H. B. Leech (CAS).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup> (<sup>1</sup>Adams, 1967:227; <sup>2</sup>Penny, NEW STATE RECORD).

**Pimachrysa fusca** Adams, 1967:227.

Holotype male: USA, California, Riverside Co., L. Covington Flat, Joshua Tree National Monument, 19.III.1961, E. L. Sleeper (CAS).

DISTRIBUTION.—USA: CA (Adams, 1967:227).

**Pimachrysa grata** Adams, 1957a:68.

Holotype female: USA, Arizona, Santa Rita Mountains, Madera Canyon, 26.VIII.1949, P. A. Adams (MCZ).

DISTRIBUTION.—USA: AZ (Adams, 1957a:67).

**Pimachrysa intermedia** Adams, 1967:228.

Holotype female: USA, California, Riverside Co., Whitewater, Snow Creek, 1500 ft., 8.III.1955, W. R. M. Mason (CNC).

DISTRIBUTION.—USA: CA (Adams, 1967:228).

**Pimachrysa nigra** Adams, 1967:228.

Holotype male: USA, California, Riverside Co., Gavilan, 19.III.1936, P. Timberlake (CAS).

DISTRIBUTION.—USA: CA (Adams, 1967:228).

CHRYSOPINAE  
BELONOPTERYGINI

**Abachrysa** Banks, 1938a:75

**Abachrysa eureka** (Banks), 1931:174  
[*Chrysopa*].

Holotype (sex unknown): USA, Arkansas, Hope, Knobel (MCZ).

TAXONOMY.—To *Abachrysa* (see Banks, 1938a:75).

DISTRIBUTION.—USA: AR<sup>1</sup>, FL<sup>5</sup>, GA<sup>3</sup>, MS<sup>2</sup>, TX<sup>4</sup>, (<sup>1</sup>Banks, 1931:174; <sup>2</sup>Banks, 1938a:75; <sup>3</sup>Bickley & MacLeod, 1956:197; <sup>4</sup>Agnew et al., 1981:4; <sup>5</sup>Brooks & Barnard, 1990:165).

**Nacarina** Navás, 1915c:133

**Nacarina robusta** (Banks), 1905b:5  
[*Chrysopa*].

Type(s) (sex unknown): USA, North Carolina, Tyron (MCZ).

TAXONOMY.—To *Nacarina* by Adams (NEW COMBINATION).

DISTRIBUTION.—USA: FL<sup>2</sup>, NC<sup>1</sup> (<sup>1</sup>Banks 1905b:5; <sup>2</sup>Stange, NEW STATE RECORD).

LEUCOCHRYGINI

**Leucochrysa** (*Leucochrysa*) McLachlan, 1868:208

**Leucochrysa** (*Leucochrysa*) *ampla* (Walker), 1853:268 [*Chrysopa*].

Lectotype female: no locality data (BMNH).

TAXONOMY.—Listed as possible synonym of *C. lineaticornis* (see Banks 1907c:27); to *Leucochrysa* (*Leucochrysa*) as valid species (see Brooks & Barnard, 1990:276); lectotype designated (see Kimmings, 1940:444).

DISTRIBUTION.—USA: GA (Walker, 1853:268).

**Leucochrysa** (*Leucochrysa*) *arizonica* (Banks), 1906a:98 [*Allochrysa*].

Holotype male: USA, Arizona, Palmerlee, VII (MCZ).

TAXONOMY.—To *Leucochrysa* (see Adams, 1977:95).

DISTRIBUTION.—USA: AZ (Banks, 1906a:98).

**Leucochrysa** (*Leucochrysa*) *insularis* (Walker), 1853:269 [*Chrysopa*].

Holotype male: Jamaica (BMNH).

= *Chrysopa virginica* Fitch, 1855:795. Holotype (sex unknown): USA, Virginian Cartersville, T. A. Culbertson (type probably lost).

= *Nothochrysa phantasma* MacGillivray, 1894:170. Lectotype male: USA, Massachusetts, West Chop, 8.VIII.1893 (MCZ).

= *Leucochrysa cerverai* Navás, 1924a:325.

Type(s) (sex unknown): Cuba, Habana, Santiago de las Vegas, 18.IX.1923, F. Z. Cervera (MCZ).

= *Leucochrysa joannisi* Navás, 1925d:13. No type specimen designated: Cuba, Habana, Santiago de las Vegas, 17.VII and 20.VIII.1924, F. Cervera (no types located).

= *Allochrysa virginica ocala* Banks, 1938b:122. Holotype female: USA, Florida, Jefferson Co., Lloyd Sink, 9.VIII.1935, G. Fairchild (MCZ).

**TAXONOMY.** — To *Protochrysopa* (see Kolbe, 1888:74); to *Leucochrysa* (see Adams, 1977:97); *N. phantasma* = *C. virginica* (see Banks, 1895b:315); *L. joannisi* = *L. insularis* (see Alayo, 1968:57); *L. virginica* = *L. insularis*, *L. cerverai* = *L. insularis*, *L. virginica ocala* = *L. insularis* (see Adams, 1977:97). It appears that no formal description of *L. joannisi* was ever published; referred to by Alayo, 1968:57.

**DISTRIBUTION.** — USA: AL<sup>8</sup>, AR<sup>8</sup>, DC<sup>7</sup>, FL<sup>4</sup>, IA<sup>9</sup>, MA<sup>2</sup>, MO<sup>8</sup>, MS<sup>8</sup>, NC<sup>5</sup>, NJ<sup>3</sup>, TN<sup>6</sup>, VA<sup>1</sup>, WV<sup>8</sup> (<sup>1</sup>Fitch, 1855:795, <sup>2</sup>MacGillivry, 1894:170, as *Nothochrysa phantasma*; <sup>3</sup>Smith, 1900:55; <sup>4</sup>Banks, 1938b:122; <sup>5</sup>Brimley, 1938:30; <sup>6</sup>Bickley, 1941:187; <sup>7</sup>Bickley & MacLeod, 1956:184; <sup>8</sup>Adams, 1977:98; <sup>9</sup>Penny, **NEW STATE RECORD**).

**Leucochrysa (Nodita)** Navás, 1916a:21

**Leucochrysa (Nodita) americana** Banks, 1897b:175.

Holotype female: USA, Alabama, Auburn, 1938 (MCZ).

**TAXONOMY.** — To *Nodita* (see Navás, 1917b:280); to *Leucochrysa (Nodita)* (see Brooks & Barnard, 1990:277).

**DISTRIBUTION.** — USA: AL<sup>1</sup>, KS<sup>3</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1897b:175; <sup>2</sup>Banks, 1907c:26; <sup>3</sup>Bickley & MacLeod, 1956:188).

**Leucochrysa (Nodita) antennata** Banks, 1905b:5.

Type (sex unknown): Mexico, Tuxpan (MCZ).

**TAXONOMY.** — To *Nodita* (see Banks, 1939:2); to *Leucochrysa (Nodita)* (see Brooks & Barnard, 1990:277).

**NOTE.** — A male specimen in MCZ from Texas, Brownsville, 13.VIII.1909, H. S. Barber is erroneously labelled as type.

**DISTRIBUTION.** — USA: TX (Banks, 1907c:26).

**Leucochrysa (Nodita) callota** Banks, 1915b:626.

Holotype female: USA, Texas, Austin, 3.V.1901 (MCZ).

**TAXONOMY.** — To *Nodita* (see Banks, 1939:2); returned to *Leucochrysa (Leucochrysa)* (see Brooks & Barnard, 1990:276); to *Leucochrysa (Nodita)* by Adams, **NEW COMBINATION**.

**DISTRIBUTION.** — USA: CA<sup>2</sup>, FL<sup>2</sup>, LA<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Banks, 1915b:626; <sup>2</sup>Adams, **NEW STATE RECORDS**).

**Leucochrysa (Nodita) explorata** (Hagen), 1861:217 [*Chrysopa*].

Type(s) (sex unknown): Mexico, Cordova, Saussure (MCZ, specimen not found).

**TAXONOMY.** — To *Leucochrysa (Nodita)* (see Brooks and Barnard, 1990:277).

**DISTRIBUTION.** — USA: NM (Hagen, 1875:921).

**Leucochrysa (Nodita) floridana** Banks, 1897c:184.

Holotype female: USA, Florida, Lake Worth (MCZ).

**TAXONOMY.** — To *Nodita* (see Navás, 1917b:280); to *Leucochrysa (Nodita)* (see Brooks & Barnard, 1990:277).

**DISTRIBUTION.** — USA: FL<sup>1</sup>, MS<sup>2</sup>, NC<sup>3</sup>, TX<sup>4</sup> (<sup>1</sup>Banks, 1897c:184; <sup>2</sup>Banks, 1903a:144; <sup>3</sup>Brimley, 1938:30; <sup>4</sup>Agnew et al., 1981:14).

**Leucochrysa (Nodita) nigrinervis** (Banks), 1939:1 [*Nodita*].

Type(s) (sex unknown): USA, New Mexico, McKinley Co., Satan Pass (MCZ, specimen not found).

**TAXONOMY.** — To *Leucochrysa (Nodita)* (see Brooks & Barnard, 1990:277).

**DISTRIBUTION.** — USA: AZ<sup>3</sup>, CA<sup>3</sup>, CO<sup>3</sup>, NM<sup>1</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1939:1; <sup>2</sup>Bickley & MacLeod, 1956:188; <sup>3</sup>Adams, **NEW STATE RECORDS**).

**Leucochrysa (Nodita) pavida** (Hagen), 1861:216 [*Chrysopa*].

Syntype series (sex unknown): Mexico, Cordova, Saussure, Deppe (NHB); USA, South Carolina, Zimmerman (female) (MCZ).

**TAXONOMY.** — To *Nodita* (see Banks, 1939:2); to *Leucochrysa (Nodita)* (see Brooks & Barnard, 1990:277).

**DISTRIBUTION.** — USA: FL<sup>2</sup>, IN<sup>4</sup>, NC<sup>2</sup>, SC<sup>1</sup>, TX<sup>3</sup> (<sup>1</sup>Hagen, 1861:216; <sup>2</sup>Bickley & MacLeod, 1956:188; <sup>3</sup>Agnew et al., 1981:14; <sup>4</sup>Lawson & McCafferty, 1984:130).

**Leucochrysa (Nodita) rufina** (Banks), 1950:54 [*Eremochrysa*].

Holotype male: USA, Arizona, Coconino Co., Grand Canyon, 24.VII.1934 (MCZ).

**TAXONOMY.** — To *Leucochrysa (Nodita)* by Adams, **NEW COMBINATION**.

**DISTRIBUTION.** — USA: AZ (Banks, 1950:55).

**Leucochrysa (Nodita) texana** (Banks), 1939:3  
[*Nodita*].

Holotype male: USA, Texas, Travis Co. (MCZ).

TAXONOMY.—To *Leucochrysa (Nodita)* (see Brooks & Barnard, 1990:278).

DISTRIBUTION.—USA: TX (Banks, 1939:3).

#### CHrysopini

**Ceraeochrysa** Adams, 1982a:70

**Ceraeochrysa cincta** (Schneider), 1851:86  
[*Chrysopa*].

Syntype series (sex unknown): Brazil, RJ, Rio de Janeiro (HMB and IRSNB).

= *Chrysopa bilineata* Navás, 1914e:91. Holotype female: Guatemala, Vulcan de Attilan, 2500–3500 ft., Champion (BMNH).

= *Chrysopa lafonei* Navás, 1914a:222. Holotype male: Argentina (MLP - Bruch collection).

= *Chrysopa bruchi* Navás, 1914a:222. Lectotype male: Argentina, Huasan, 25.II.1912 (MACN) (Adams, NEW SYNONYMY).

= *Chrysopa incalis* Banks, 1915b:627. Syntypes (sex unknown): Peru, Matucana, 7780 ft., 14.VI, Parish (male); Chosica, 2800 ft., 10.VI, Parish (MCZ).

= *Chrysopa bicarnea* Banks, 1920:338. Holotype male: USA, Florida, Miami, G. F. Mozzette (MCZ).

= *Chrysopa advena* Navás, 1922a:51. Holotype male: Coast of Uruguay, 1920, Renato Martin (MNHN).

= *Chrysopa habana* Navás, 1922a:52. Holotype (sex unknown): Cuba, Havana, Fermín Z. Cervera (MZB).

= *Chrysopa bessona* Navás, 1924d:361. Holotype female: Argentina, La Plata, 20.XI.1920, Bruch (MLP).

= *Chrysopa mestiza* Navás, 1924a:330. Lectotype female: Cuba, Santiago de las Vegas, IX.1923; Havana, Rio Almendares, 19.VII.1923, Fermín Z. Cervera (MCZ). NEW LECTOTYPE DESIGNATION.

= *Chrysopa villosoala* Navás, 1924b:337. Holotype (missing abdomen): Cuba, Havana, Santiago de las Vegas, 4.XII.1923 (MCZ).

= *Cintameva bina* Navás, 1924a:331. Holotype male: Cuba, Havana, Rio Almendares, 9.VIII.1923 (MCZ).

= *Chrysopa cornuta* Navás, 1925a:65. Holotype female: Brazil, RJ, Niterói, 21.XI.1924 (MNHN).

= *Chrysopa alternans* Navás, 1933a:306. Holotype male: Peru, Lima, 4.X.1933 (HMB).

= *Chrysopa wollebaeki* Esben-Petersen, 1934:291. Seven syntypes (six males, one female): Ecuador, Galapagos Islands, Santa Maria, Post Office Bay, 1–10.X.1925, one male; 29.X.1925, four males, one female; 48.XI.1925, one male, all collected by Wollebaek (ZMUO).

= *Chrysopa iona* Banks, 1944:12. Holotype male: Surinam, Paramaribo, 7.XII.1938, Geijskes (MCZ).

= *Chrysopodes sallei* Banks, 1946:171. Holotype male: Mexico, Salle (MCZ).

TAXONOMY.—*C. mestiza* = *C. habana* (see Alayo, 1968:30); *C. advena*, *C. alternans*, *C. bessona*, *C. bicarnea*, *C. bilineata*, *C. bina*, *C. caligata*, *C. cornuta*, *C. habana*, *C. incalis*, *C. iona*, *C. lafonei*, *C. mestiza*, *Chrysopodes sallei*, *C. villosoala*, *C. wollebaeki* = *C. cincta* (Adams, 1982:72); *C. caligata* not = *C. cincta* (Adams & Penny, 1987:442).

DISTRIBUTION.—USA: FL (Banks, 1920:338).

**Ceraeochrysa cubana** (Hagen), 1861:215  
[*Chrysopa*].

Neotype male: Cuba, Soledad, near Cienfuegos, 6–20.VIII, N. Banks (MCZ).

= *Chrysopa tolteca* Banks, 1901a:364. Holotype (female): Mexico, Oaxaca, Tomellin, VI (MCZ).

= *Chrysopa venularis* Navás, 1913b:20. Holotype female: Jamaica (ZSBS).

= *Chrysopa albatala* Banks, 1913a:139. Holotype (abdomen missing): Guyana, Bartica, XII, Parish (MCZ).

= *Chrysopa imbecilla* Navás, 1914e:107. Holotype male: Barbados (BMNH).

= *Chrysopa epheba* Navás, 1924a:328. Holotype (sex unknown): Cuba, Calabazas, 28.VIII.1923 (MZB).

= *Chrysopa seminole* Banks, 1924:432. Holotype female: USA, Florida, Marco, 24.VII, Mott (MCZ).

= *Chrysopa freemani* Smith, 1931:811. Holotype female: Haiti, Port-au-Prince, 500 ft., 16.XI.1929 (MCZ).

= *Chrysopa fischerina* Navás, 1933:196. Holotype female: Brazil, Central. Mang., 13.X.1913, R. Fischer (DEIC) (Adams, NEW SYNONYMY).

= *Chrysopa damiensis* var. *jamaicensis* Banks, 1941a:394.

Syntype series (male, female): Jamaica, Kingston, 27–29.VIII, P. J. Darlington; Trelawny, Perkins (MCZ).

**TAXONOMY.** — To *Ceraeochrysa* (see Adams, 1982:72); *C. epheba* = *C. cubana*, *C. jacobaea* = *C. cubana* (see Alayo, 1968:28); Adams, 1982:72 (*C. jacobaea* = *C. gundlachi*, *C. tolteca* = *C. cubana*, *C. albatala* = *C. cubana*, *C. venularis* = *C. cubana*, *C. imbecilla* = *C. cubana*, *C. epheba* = *C. cubana*, *C. seminole* = *C. cubana*, *C. freemani* = *C. cubana*, *C. damiensis jamaicensis* = *C. cubana*, *C. scapularis* = *C. cubana* (see Adams, 1982:72); *C. scapularis* not = *C. cubana* (see Adams & Penny, 1987:448); neotype designated (see Adams & Penny, 1987:446).

**DISTRIBUTION.** — USA: FL<sup>2</sup>, NC<sup>3</sup>, VA<sup>1</sup> (<sup>1</sup>Hagen, 1861:215; <sup>2</sup>Banks, 1924:432, as *C. seminole*; <sup>3</sup>Banks, 1938b:121).

***Ceraeochrysa lineaticornis*** (Fitch), 1855:795  
[*Chrysopa*].

Holotype (sex unknown): USA, New York (MCZ).

= *Chrysopa puncticornis* Fitch, 1855:796. Type(s) (sex unknown): USA, New York (MCZ, specimen not found).

= *Allochrysa parvula* Banks, 1903a:143. Holotype (male): USA, Florida, Runnymede (MCZ).

= *Chrysopa columbiana* Banks, 1903a:150. Holotype male: USA, D. C., Washington (MCZ).

= *Chrysopa stichoptera* Navás, 1914c:61. Holotype (sex unknown): USA, New York, Long Island, Yaphank, 10.VII.1913, J. R. de la Torre-Bueno (MZB).

**TAXONOMY.** — To *Ceraeochrysa* (see Adams, 1982:73) *C. puncticornis* = *C. lineaticornis* (see Banks, 1907c:27); *C. stichoptera* = *C. lineaticornis* (see Bickley & MacLeod, 1956:190); *C. columbiana* = *C. lineaticornis* (see Bram & Bickley, 1963:16); *A. parvula* = *C. lineaticornis* (see Adams, 1982:73).

**DISTRIBUTION.** — CAN: BC<sup>3</sup>, PQ<sup>3</sup>, USA: DC<sup>2</sup>, FL<sup>2</sup>, GA<sup>10</sup>, IN<sup>9</sup>, KS<sup>10</sup>, MA<sup>2</sup>, MD<sup>2</sup>, ME<sup>10</sup>, MI<sup>2</sup>, NC<sup>4</sup>, NH<sup>2</sup>, NY<sup>1</sup>, PA<sup>10</sup>, TN<sup>5</sup>, TX<sup>8</sup>, VA<sup>6</sup>, WS<sup>7</sup> (<sup>1</sup>Fitch, 1855:795; <sup>2</sup>Banks, 1903a:143, 150, 151 as *A. parvula*, *C. lineaticornis* and *C. colum-*

*biana*; <sup>3</sup>Smith, 1932:585, as *C. columbiana* and *C. lineaticornis*; <sup>4</sup>Brimley, 1938:30; <sup>5</sup>Bickley, 1941:197; <sup>6</sup>Bickley & MacLeod, 1956:191; <sup>7</sup>Throne, 1971a:74; <sup>8</sup>Agnew et al., 1981:12; <sup>9</sup>Lawson & McCafferty, 1984:130; <sup>10</sup>Adams, NEW STATE RECORDS).

***Ceraeochrysa placita*** (Banks), 1908a:259  
[*Chrysopa*].

Male syntypes: USA, Colorado, Chimney Gulch, 20.I.1907; Clear Creek (MCZ).

= *Chrysopa intacta* Navás, 1912a:199. Holotype (sex unknown): Canada, Quebec, Toronto, 1.VI.1908 (MZB).

= *Chrysopa forreri* Navás, 1914e:97. Type(s) (sex unknown): Mexico, Mexico City, 8100 ft., Forrer (BMNH).

**TAXONOMY.** — To *Ceraeochrysa* (see Adams, 1982:73); *C. forreri* = *C. placita* (see Adams, 1982:73); *C. intacta* = *C. placita* (see Brooks & Barnard, 1990:269).

**DISTRIBUTION.** — CAN: PQ<sup>2</sup>, USA: CA<sup>4</sup>, CO<sup>1</sup>, MO<sup>3</sup>, NH<sup>4</sup>, QR<sup>4</sup> (<sup>1</sup>Banks, 1908a:259; <sup>2</sup>Navás, 1912a:200; <sup>3</sup>Froeschner, 1947:134, as *C. intacta*; <sup>4</sup>Adams, NEW STATE RECORDS).

***Ceraeochrysa smithi*** (Navás), 1914e:105  
[*Chrysopa*].

Two male syntypes: Grenadines, Union Island, W. J. H. H. Smith; West Indies, Mustiques (BMNH).

= *Chrysopa poeyi* Navás, 1924a:330. Holotype (male): Cuba, Havana, Rio Almendares, 20.VII.1923, Cervera (MCZ).

= *Chrysopa neotropica* Navás, 1929b:317. Syntype series (sex unknown): Honduras, Amapala, 9.VII.1908, R. Paessler; El Salvador, La Unión, 3.XII.1907, R. Paessler; Mexico, Jalisco, La Garita, 1912, W. Fritsche; Colombia, Behn verid., 3.XII.1900 (Hamburg, specimen lost).

**TAXONOMY.** — To *Ceraeochrysa* (see Adams, 1982:73); *C. neotropica* = *C. smithi*, *C. poeyi* = *C. smithi* (see Adams, 1982:73).

**DISTRIBUTION.** — USA: FL (Adams, NEW NATIONAL RECORD).

**REMARKS.** — This species has been mentioned before from Florida by Banks under the names *C. claveri* and *C. cubana sanchezi*. Several specimens have been identified by other chrysopid specialists, but this distribution has not been reported in the taxonomic literature.

**Ceraeochrysa valida** (Banks), 1895:517  
[*Chrysopa*].

Two syntypes (sex unknown): Mexico, Baja California Sur, Serra El Taste, IX; San José del Cabo, IX (MCZ).

= *Chrysopa bimaculata* McClendon, 1901:215. Lectotype female: USA, Texas, Laredo, VIII.1900 (MCZ) **NEW LECTOTYPE DESIGNATION**.

= *Chrysopa limitata* Navás, 1913c:84. Holotype (sex unknown): Brazil, Rio de Janeiro, Nova Friburgo, XII.1911, Ramos (MZB).

= *Chrysopa longicella* Navás, 1914e:97. Holotype male: Guatemala, Zapote, G. C. Champion (BMNH).

= *Chrysopa breviata* Banks, 1915b:628. One male syntype, one syntype of unknown sex. Ecuador, Guayaquil, 8.VI, Parish; Quevedo (MCZ).

= *Chrysopa lioni* Navás, 1927a:54. Holotype female: Haiti, Port-au-Prince, 1910, G. Lion (MNHN).

= *Chrysopa damiensis* Smith, 1931:803. Holotype male: Haiti, Hatte Lathan, 18.I.1929, R. C. Smith (MCZ).

= *Chrysopa wolcotti* Smith, 1931:810. Holotype male: Haiti, Damien (MCZ).

**TAXONOMY.** — To *Ceraeochrysa* (see Adams, 1982:73); *C. bimaculata* = *C. tolteca*, *C. longicella* = *C. tolteca*, *C. tolteca* (?) = *C. valida* (see Banks, 1946:148); *C. bimaculata* = *C. valida*, *C. limitata* = *C. valida*, *C. longicella* = *C. valida*, *C. breviata* = *C. valida*, *C. lioni* = *C. valida*, *C. damiensis* = *C. valida*, *C. wolcotti* = *C. valida*, *C. camposana* = *C. valida*, *C. seminole* = *C. valida*, *C. tolteca* = *C. cubana* (see Adams, 1982:73); *C. seminole* = *C. cubana*; *C. camposana* = *Chrysopa incertae sedis* (see Brooks & Barnard, 1990:268, 279).

**DISTRIBUTION.** — **USA:** FL<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>McClendon, 1901:215, as *C. bimaculata*; <sup>2</sup>Banks, 1903a:153).

**Chrysopa** Leach in Brewster, 1815:138**Chrysopa chi** Fitch, 1855:791.

Holotype female: USA, New York (MCZ). = *Chrysopa epsilon* Fitch, 1855:791. Holotype female: no label data (MCZ).

= *Chrysopa epsilon* var. *haematica* Navás, 1918a:354. (as *Chrysopa hypsilon* [sic]) Holo-

type (sex unknown): USA, New York, White Plains, 31.V.1915, de la Torre Bueno (MZB).

**TAXONOMY.** — *C. epsilon* = *C. chi* (see Smith, 1922:1352); *C. hypsilon* var. *haematica* = *C. chi* (see Smith, 1932:592).

**DISTRIBUTION.** — **CAN:** AB<sup>5</sup>, BC<sup>5</sup>, MB<sup>5</sup>, NS<sup>5</sup>, ON<sup>5</sup>, PQ<sup>5</sup>, SK<sup>5</sup>, **USA:** DC<sup>2</sup>, MD<sup>9</sup>, ME<sup>7</sup>, MN<sup>8</sup>, NJ<sup>3</sup>, NH<sup>4</sup>, NY<sup>1</sup>, TN<sup>6</sup>, WA<sup>11</sup>, WS<sup>10</sup> (<sup>1</sup>Fitch, 1855:791; <sup>2</sup>Hagen, 1861:214; <sup>3</sup>Smith, 1900:55; <sup>4</sup>Banks, 1903a:148; <sup>5</sup>Smith, 1932:592; <sup>6</sup>Bickley, 1941:188; <sup>7</sup>Procter, 1946:43; <sup>8</sup>Parfin, 1952:424; <sup>9</sup>Bram & Bickley, 1963:8; <sup>10</sup>Throne, 1971a:70; <sup>11</sup>Penny, **NEW STATE RECORD**). **Chrysopa coloradensis** Banks, 1895b:314.

Holotype male: USA, Colorado, Fort Collins (MCZ).

**DISTRIBUTION.** — **CAN:** BC<sup>3</sup>, **USA:** AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>1</sup>, ID<sup>3</sup>, NM<sup>2</sup>, OR<sup>2</sup>, UT<sup>4</sup>, WA<sup>2</sup> (<sup>1</sup>Banks, 1895b:315; <sup>2</sup>Banks, 1903a:151; <sup>3</sup>Smith, 1932:585; <sup>4</sup>Bickley & MacLeod, 1956:190).

**Chrysopa excepta** Banks, 1911:340.

Holotype (sex unknown): USA, New Mexico, Fort Wingate, 25.VII, Woodgate (MCZ).

**DISTRIBUTION.** — **CAN:** AB<sup>2</sup>, BC<sup>2</sup>, **USA:** NM<sup>1</sup>, UT<sup>3</sup>, WY<sup>3</sup> (<sup>1</sup>Banks, 1911:340; <sup>2</sup>Smith, 1932:585; <sup>3</sup>Bickley & MacLeod, 1956:190).

**Chrysopa incompleta** Banks, 1911:340.

Syntypes (sex unknown): USA: North Carolina, Beaufort, 15.V.1906, Woglum (male); Raleigh, 15.VII.1902, Sherman (MCZ).

**DISTRIBUTION.** — **USA:** FL<sup>6</sup>, GA<sup>2</sup>, IN<sup>5</sup>, MA<sup>2</sup>, MD<sup>2</sup>, NC<sup>1</sup>, NJ<sup>2</sup>, TX<sup>4</sup>, VA<sup>2</sup>, WS<sup>3</sup> (<sup>1</sup>Banks, 1911:341; <sup>2</sup>Bram & Bickley, 1963:7; <sup>3</sup>Throne, 1971a:72; <sup>4</sup>Agnew et al., 1981:12; <sup>5</sup>Lawson & McCafferty, 1984:130; <sup>6</sup>Adams, **NEW STATE RECORD**).

**Chrysopa nigricornis** Burmeister, 1839:980.

Holotype male: North America (Berlin).

= *Chrysopa colon* Fitch, 1855:792. Type (sex unknown): (MCZ).

= *Chrysopa erythrocephala* Banks, 1898:201 not *C. erythrocephala* (Rambur) 1842:428 [Hemerobius]. Holotype female: USA, California, San Francisco, VII.1897, Morse (MCZ).

= *Chrysopa majuscula* Banks, 1906a:98. Replacement name for *Chrysopa erythrocephala* Banks.

= *Chrysopa vegeta* Navás, 1917:6. Type(s) (sex unknown): USA, New Mexico, Jemez Springs, 21.VI.1916, John Woodgate (MZB).

= *Chrysopa crotchi* Banks, 1938a:76. Holotype (female): Canada, British Columbia, Vancouver Island, Victoria, VII, Crotch (MCZ).

**TAXONOMY.** — *C. colon* = *C. nigricornis* (see Hagen, 1875:920); *C. majuscula*, new name for *C. erythrocephala* Banks (see Banks, 1906a:98); *C. majuscula* = *C. nigricornis* (see Bram & Bickley, 1963:10); *C. crotchi* = *C. nigricornis*; *C. vegeta* = *C. nigricornis* (Adams, **NEW SYNONYMY**).

**DISTRIBUTION.** — CAN: AB<sup>5</sup>, BC<sup>5</sup>, ON<sup>5</sup>, PQ<sup>4</sup>. USA: CA<sup>3</sup>, CO<sup>2</sup>, DC<sup>4</sup>, IN<sup>6</sup>, KS<sup>4</sup>, MA<sup>4</sup>, MD<sup>12</sup>, ME<sup>9</sup>, MN<sup>11</sup>, MO<sup>10</sup>, NC<sup>7</sup>, NM<sup>2</sup>, NY<sup>1</sup>, OH<sup>4</sup>, RI<sup>4</sup>, TN<sup>8</sup>, TX<sup>14</sup>, VA<sup>4</sup>, WS<sup>13</sup> (Fitch, 1855:792; <sup>2</sup>Hagen, 1875:920; <sup>3</sup>Banks, 1898:201; <sup>4</sup>Banks, 1903a:150; <sup>5</sup>Smith, 1932:584; <sup>6</sup>Montgomery & Trippel, 1933:260; <sup>7</sup>Brimley, 1938:30; <sup>8</sup>Bickley, 1941:188; <sup>9</sup>Procter, 1946:43; <sup>10</sup>Froeschner, 1947:133; <sup>11</sup>Parfin, 1952:424; <sup>12</sup>Bram & Bickley, 1963:9; <sup>13</sup>Throne, 1971a:69; <sup>14</sup>Agnew et al., 1981:12).

### *Chrysopa oculata* Say, 1839:45.

Type(s) (sex unknown): USA (depository unknown).

= *Chrysopa chlorophana* Burmeister, 1839:979. Type(s) (sex unknown): North America (depository unknown).

= *Chrysopa euryptera* Burmeister, 1839:980. Type(s) (sex unknown): North America (?) (Halle).

= *Chrysopa latipennis* Schneider, 1851:118. Holotype (sex unknown): North America (HMB or Frankfurt).

= *Chrysopa albicornis* Fitch, 1855:788. Holotype (sex unknown): USA, Mississippi (MCZ).

= *Chrysopa illepeda* Fitch, 1855:788. Holotype (female): USA, New York; Illinois (MCZ).

= *Chrysopa omikron* Fitch, 1855:789. Holotype (female): USA, New York (MCZ).

= *Chrysopa xanthocephala* Fitch, 1855:789. Holotype (sex unknown): USA, New York; Michigan (MCZ).

= *Chrysopa fulvibucca* Fitch, 1855:790. Type(s) (sex unknown): USA, New York (Type not found).

= *Chrysopa mississippiensis* Fitch, 1855:790. Holotype female: USA, Mississippi, Jackson, IV, Fitch (MCZ).

= *Chrysopa bipunctata* Fitch, 1855:791. Holotype female: USA, New York (MCZ).

= *Chrysopa transmarina* Hagen, 1861:213. Syntypes (sex unknown): Canada, Ontario, La

Chine, near Montreal, Barnston; Trenton Falls, New York; Newfoundland, St. John; Nova Scotia (Types should be in BMNH).

= *Nothochrysa annulata* MacGillivray, 1894:169. Holotype female: USA, Massachusetts, Wellesley, 22.VI.1892 (CORN).

= *Chrysopa assimilis* Banks, 1898:202. One male syntype, one syntype of unknown sex: USA, Oregon, Ashland (male) and Hood River (MCZ).

= *Chrysopa mexicana* Banks, 1901:364. Six syntypes (sex unknown): Mexico, Veracruz, Thalpan, VII (MCZ) Adams, **NEW SYNONYMY**.

= *Chrysopa separata* Banks, 1911:341. Syntypes (sex unknown): USA, Colorado, Golden, Chimney Gulch, Oslar; New Mexico, Pecos, 23.VI, Cockerell (MCZ).

= *Chrysopa rubicunda* Navás, 1913b:20. Type(s) (sex unknown): USA, New York, Long Island, Yapbank, 23.IX.1911, Torre Bueno (MZB).

= *Cintameva conspersa* Navás, 1929:318. Adams, **NEW SYNONYMY**. Holotype male: Colombia (?), Behn vend., 3.XII.1900 (Hamburg, specimen destroyed).

**TAXONOMY.** — *C. euryptera* = *C. oculata*, *C. illepeda* = *C. oculata*, *C. omikron* = *C. oculata*, *C. fulvibucca* = *C. oculata*, *C. mississippiensis* = *C. oculata*, *C. latipennis* = *C. chlorophana*, *C. xanthocephala* = *C. chlorophana*, *C. bipunctata* = *C. chlorophana*, *C. transmarina* = *C. chlorophana* (see Banks, 1903a:161); *C. chlorophana* and earlier synonymies = *C. oculata*, *C. albicornis* = *C. oculata*, *Nothochrysa annulata* (?) = *C. oculata*, *C. separata* = *C. oculata*, *C. rubicunda* = *C. oculata* (see Bickley & MacLeod, 1956:191); *C. assimilis* = *C. oculata*, *C. separata* is valid species, *C. chlorophana* is valid species with associated earlier synonymies (see Brooks & Barnard, 1990:270). We consider that the synonymy of *C. chlorophana* and *C. oculata* by Bickley and MacLeod is a correct decision.

**DISTRIBUTION.** — CAN: AB<sup>8</sup>, BC<sup>8</sup>, MB<sup>8</sup>, NF<sup>6</sup>, NS<sup>6</sup>, ON<sup>8</sup>, PQ<sup>6</sup>, SK<sup>8</sup>, USA: AZ<sup>6</sup>, CO<sup>6</sup>, DC<sup>2</sup>, GA<sup>2</sup>, IA<sup>16</sup>, IL<sup>2</sup>, IN<sup>9</sup>, KS<sup>7</sup>, LA<sup>2</sup>, MD<sup>2</sup>, ME<sup>11</sup>, MI<sup>3</sup>, MN<sup>13</sup>, MO<sup>12</sup>, MS<sup>2</sup>, NC<sup>10</sup>, NJ<sup>5</sup>, NM<sup>6</sup>, NY<sup>1</sup>, OH<sup>6</sup>, OR<sup>4</sup>, PA<sup>2</sup>, TN<sup>2</sup>, TX<sup>15</sup>, VA<sup>2</sup>, WS<sup>14</sup> (Fitch, 1855:788; <sup>2</sup>Hagen, 1861:212; <sup>3</sup>Banks, 1892:358; <sup>4</sup>Banks, 1898:202, as *C. chlorophana*; <sup>5</sup>Smith, 1900:55; <sup>6</sup>Banks, 1903a:148, as *C. chlorophana*; <sup>7</sup>Smith, 1925:168; <sup>8</sup>Smith,

1932:587; <sup>9</sup>Montgomery & Trippel, 1933:260;  
<sup>10</sup>Brimley, 1938:30; <sup>11</sup>Procter, 1946:43;  
<sup>12</sup>Froeschner, 1947:133; <sup>13</sup>Parfin, 1952:424;  
<sup>14</sup>Throne, 1971a:69; <sup>15</sup>Agnew et al., 1981:12;  
<sup>16</sup>Penny, **NEW STATE RECORD**).

REMARKS.—Carpenter (1992:538) has indicated that Burmeister's *Handbuch der Entomologie* was apparently published in two parts. The first 400 pages were published in 1839, while the remainder, including all Neuroptera, were published in 1838. If this is found to be correct, it could lead to serious nomenclatorial instability. The synonymy of *Chrysopa oculata* and *Chrysopa chlorophana* would be reversed. Almost certainly the International Commission on Zoological Nomenclature would be petitioned to declare, for nomenclatorial purposes, that the entire book was published in 1839.

***Chrysopa pleuralis* Banks, 1911:341.**

One male syntype, one syntype of unknown sex: USA, Colorado, Boulder Co., North Boulder Creek, 21.VIII.1907, S. A. Rohwer; Steamboat Springs, 27.V. Cockerell (male) (MCZ).

DISTRIBUTION.—CAN: AB<sup>1</sup>, BC<sup>2</sup>, USA: CO<sup>1</sup>, MT<sup>3</sup>, UT<sup>3</sup>, WY<sup>3</sup> (<sup>1</sup>Banks, 1911:342; <sup>2</sup>Smith, 1932:591; <sup>3</sup>Adams, **NEW STATE RECORDS**).

***Chrysopa quadripunctata* Burmeister, 1839:980.**

Type(s) (sex unknown): North America (Halle).

= *Chrysopa sulphurea* Fitch, 1855:793. Type(s) (sex unknown): USA, New Jersey, IX (MCZ, specimen not found).

= *Chrysopa sichelii* Fitch, 1855:793. Holotype male: USA, New York (MCZ).

TAXONOMY.—*C. sulphurea* = *C. quadripunctata*, *C. sichelii* = *C. quadripunctata* (see Banks, 1903a:162).

DISTRIBUTION.—CAN: BC<sup>6</sup>, USA: DC<sup>2</sup>, IL<sup>6</sup>, IN<sup>7</sup>, KS<sup>5</sup>, MD<sup>12</sup>, MN<sup>11</sup>, MO<sup>10</sup>, NC<sup>8</sup>, NJ<sup>3</sup>, NY<sup>1</sup>, PA<sup>2</sup>, SC<sup>2</sup>, TN<sup>9</sup>, TX<sup>4</sup>, VA<sup>4</sup>, WS<sup>13</sup> (<sup>1</sup>Fitch, 1855:793; <sup>2</sup>Banks, 1892:359; <sup>3</sup>Smith, 1900:55; <sup>4</sup>Banks, 1903a:153; <sup>5</sup>Smith, 1925:168; <sup>6</sup>Smith, 1932:597; <sup>7</sup>Montgomery & Trippel, 1933:260; <sup>8</sup>Brimley, 1938:30; <sup>9</sup>Bickley, 1941:189; <sup>10</sup>Froeschner, 1947:134; <sup>11</sup>Parfin, 1952:425; <sup>12</sup>Bram & Bickley, 1963:8; <sup>13</sup>Throne, 1971a:70).

***Chrysopa slossonae* Banks, 1924:432.**

Syntypes females: USA, North Carolina, Hendersonville, F. Sherman; New Hampshire, Fran-

conia, Mrs. A. T. Slosson; Virginia, Great Falls (MCZ).

DISTRIBUTION.—USA: NC<sup>1</sup>, NH<sup>1</sup>, NY<sup>2</sup> VA<sup>1</sup> (<sup>1</sup>Banks, 1924:432; <sup>2</sup>Eisner et al., 1978:790).

***Chrysoperla* Steinmann, 1964:260**

***Chrysoperla adamsi* Henry, Wells and Pupedis, 1993:9.**

Holotype male: USA, Oregon, Benton Co., Philomath (Mary's Peak), 28.IX.1989, C. S. Henry & M. M. Wells (YPM).

DISTRIBUTION.—USA: CA, ID, OR, WA (Henry, Wells & Pupedis, 1993:9).

***Chrysoperla comanche* (Banks), 1938b:119 [*Chrysopa*].**

Holotype female: USA, Texas, Laredo (MCZ).

= *Chrysopa sperryae* Banks, 1943:74. Lectotype female: USA, California, Riverside, 7.IX.1940, Sperry (MCZ).

TAXONOMY.—To *Chrysoperla* (see Brooks & Barnard, 1990:271); *C. sperryae* = *C. comanche* (see Adams, 1958:45); lectotype designated for *C. sperryae* (see Adams, 1958:45).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: AZ<sup>1</sup>, CA<sup>1</sup>, CO<sup>2</sup>, NM<sup>1</sup>, TX<sup>1</sup> (<sup>1</sup>Banks, 1938b:119; <sup>2</sup>Bickley & MacLeod, 1956:194, as *C. sperryae*).

***Chrysoperla downesi* (Smith), 1932:594 [*Chrysopa*].**

Holotype female: Canada, British Columbia, Kelowna, W. Downes (CNC).

= *Chrysopa mohave* Banks, 1938b:120. Syntypes (sex unknown): USA, California, Claremont; Stanford University campus; Arizona, Chiricahua Mountains (MCZ).

TAXONOMY.—To *Chrysoperla* (see Henry, 1985:965); *C. mohave* synonymized with *C. carnea* (see Tauber & Tauber, 1973:1153); *C. downesi* synonymized with *C. carnea* (see Garland, 1985:753); *C. downesi* and *C. mohave* considered synonyms of *C. carnea* (see Brooks & Barnard, 1990:271); *C. downesi* considered valid species with *C. mohave* as synonym (see Henry, 1993:23).

DISTRIBUTION.—CAN: BC<sup>1</sup>, SK<sup>1</sup>, USA: AZ<sup>2</sup>, CA<sup>2</sup>, ID<sup>3</sup>, MA<sup>3</sup>, MT<sup>3</sup>, NH<sup>3</sup>, NY<sup>3</sup>, VT<sup>3</sup> (<sup>1</sup>Smith, 1932:595; <sup>2</sup>Banks, 1938b:120, as *C. mohave*; <sup>3</sup>Henry, 1985:967).

***Chrysoperla externa* (Hagen), 1861:221 [*Chrysopa*].**

Neotype female: USA, Winter Park, Florida, 3.IX, E. M. Davis (MCZ).

= *Chrysopa lanata* Banks, 1910:154. Holotype male: Argentina, Mendoza, Jensen-Haarup Expedition (MCZ).

= *Chrysopa graciana* Navás, 1919b:301. Holotype (sex unknown): Argentina, Alta Gracia, II.1918, C. Bruch (MLP).

**TAXONOMY.** — *C. lanata* = *C. externa* (Adams, 1963:222); *C. graciana* = *C. externa* (Adams & Penny, 1987:421).

**DISTRIBUTION.** — USA: FL<sup>1</sup>, SC<sup>1</sup>, TX<sup>2</sup> (Adams, 1963:222; Agnew et al., 1981:9).

**NOTE.** — The specimens listed by Hagen, 1875:921 and Smith, 1932:596 are misidentifications.

***Chrysoperla harrisii* (Fitch), 1855:794** [*Chrysopa*].

Holotype male: USA, New York (MCZ).

= *Chrysopa stenostigma* Navás, 1914c:61. Holotype (abdomen missing): USA, New York, Long Island, Yaphank, 25.VII.1913 (MZB).

**TAXONOMY.** — *C. harrisii* is a replacement name for the *C. perla* of Harris, 1841:197. To *Chrysoperla* (see Garland, 1985:755); *C. stenostigma* = *C. harrisii* (Bickley & MacLeod, 1956:193).

**DISTRIBUTION.** — CAN: BC<sup>5</sup>, ON<sup>5</sup> USA: AK<sup>11</sup>, DC<sup>3</sup>, IN<sup>6</sup>, KS<sup>4</sup>, MD<sup>12</sup>, MI<sup>3</sup>, MN<sup>10</sup>, MO<sup>9</sup>, NC<sup>7</sup>, NH<sup>3</sup>, NJ<sup>2</sup>, NY<sup>1</sup>, TN<sup>8</sup>, TX<sup>14</sup>, WS<sup>13</sup> (Fitch, 1855:794; Smith, 1900:55; Banks, 1903a:155; Smith, 1925:169; Smith, 1932:595; Montgomery & Trippel, 1933:260; Brimley, 1938:30; Bickley, 1941:188; Froeschner, 1947:134; Parfin, 1952:425; Bickley & MacLeod, 1956:193; Bram & Bickley, 1963:7; Throne, 1971a:73; Agnew et al., 1981:9).

***Chrysoperla johnsoni* Henry, Wells and Pupedis, 1993:10.**

Holotype male: USA, Oregon, Benton Co., Philomath (Mary's Peak), 28.IX.1989, C. S. Henry & M. M. Wells (YPM).

**DISTRIBUTION.** — USA: AZ, CA, ID, OR, WA (Henry et al., 1993:10).

***Chrysoperla plorabunda* (Fitch), 1855:792** [*Chrysopa*].

One male syntype, one syntype of unknown sex: USA, Illinois (no head or abdomen) (USNM); female (no label data, except #8592) (MCZ).

= *Chrysopa robertsonii* Fitch, 1855:792. Holotype female: USA, Arkansas, V. W. S. Robertson (MCZ).

= *Chrysopa pseudographa* Fitch, 1855:793. Lectotype male: USA, Illinois, X (MCZ) **NEW LECTOTYPE DESIGNATION.**

= *Chrysopa illinoiensis* Shimer, 1865:208. Type(s) (sex unknown): USA, Illinois (depository unknown).

= *Chrysopa californica* Coquillett, 1890:288. Holotype female: USA, California, Los Angeles, Coquillett (USNM).

= *Chrysopa simplex* Navás, 1908:402. Type(s) (sex unknown): USA, Rocky Mountains, R. P. Joannis (MZB).

**TAXONOMY.** — To *Chrysoperla* (see Brooks & Barnard, 1990:271); *C. robertsonii* = *C. plorabunda*, *C. pseudographa* = *C. plorabunda*, *C. illinoiensis* = *C. plorabunda* (see Banks, 1903a:162); *C. californica* = *C. plorabunda* (see Smith, 1932:594); *C. californica* = subspecies of *C. plorabunda* (see Adams, 1956:45); *C. plorabunda* = *C. carnea* (see Tjeder, 1960:148); *C. plorabunda* not = *C. carnea* (see Henry, 1983:293); *C. simplex* = *C. plorabunda* (Adams, 1956:193). **NEW SYNONYMY.**

**DISTRIBUTION.** — CAN: AB<sup>5</sup>, BC<sup>5</sup>, MB<sup>5</sup>, SK<sup>5</sup>, USA: AK<sup>10</sup>, AZ<sup>14</sup>, CO<sup>3</sup>, CT<sup>13</sup>, DC<sup>14</sup>, FL<sup>14</sup>, IA<sup>3</sup>, ID<sup>13</sup>, IL<sup>1</sup>, IN<sup>6</sup>, KS<sup>4</sup>, MD<sup>11</sup>, ME<sup>14</sup>, MI<sup>3</sup>, MN<sup>3</sup>, MO<sup>9</sup>, MS<sup>3</sup>, MT<sup>13</sup>, NC<sup>7</sup>, NJ<sup>2</sup>, NM<sup>14</sup>, NY<sup>1</sup>, OH<sup>3</sup>, OK<sup>1</sup>, OR<sup>14</sup>, TN<sup>8</sup>, TX<sup>3</sup>, WA<sup>14</sup>, WS<sup>12</sup> (Fitch, 1855:792; Smith, 1900:55; Banks, 1903a:155; Smith, 1925:168; Smith, 1932:593; Montgomery & Trippel, 1933:260; Brimley, 1938:30; Bickley, 1941:188; Froeschner, 1947:134; Parfin, 1952:425; Bickley & MacLeod, 1956:193; Bram & Bickley, 1963:4, as *C. carnea*; Throne, 1971a:72, as *C. carnea*; Henry, 1983:293; Agnew et al., 1981:9).

**REMARKS.** — Henry (1983:293) indicated that the holotype of *C. plorabunda* is at MCZ. In fact, this is one of two syntypes, and no lectotype has yet been designated. Henry, Wells and Pupedis (1993:9) reported that *C. plorabunda* possibly does not occur in California, so that previous records from this state probably pertain to *C. adamsi* or *C. johnsoni*, which are morphologically indistinguishable from *C. plorabunda*.

***Chrysoperla rufilabris* (Burmeister), 1839:979** [*Chrysopa*].

Syntypes (sex unknown): USA and Mexico (ZMH, specimen destroyed).

= *Chrysopa interrupta* Schneider, 1851:76. Holotype male: USA, Pennsylvania (HMB).

= *Chrysopa attenuata* Walker, 1853:242. Lectotype male: USA, east Florida, St. John's Bluff, E. Doubleday (BMNH).

= *Chrysopa repleta* Walker, 1853:244. Holotype male: USA, Georgia, Abbot (BMNH).

= *Chrysopa novaeboracensis* Fitch, 1855:794. Holotype female: USA, New York (MCZ).

= *Chrysopa tabida* Fitch, 1855:796. Type(s) (sex unknown): USA (MCZ, specimen not found).

= *Chrysopa citri* Ashmead, 1880:12. Holotype male: USA, Florida, W. H. Ashmead & S. Henshaw (MCZ).

= *Chrysopa medialis* Banks, 1903a:154. Holotype female: USA, Maryland, High Island, IX (MCZ).

TAXONOMY.—To *Chrysoperla* (see Garland, 1985:756); Hagen, 1866:395, 398 (*C. novaeboracensis* = *C. rufilabris*, *C. tabida* = *C. rufilabris*) (see Hagen, 1866:395, 398); *C. repleta* = *C. rufilabris*, *C. attenuata* = *C. rufilabris* (see Banks, 1903a:161); *C. citri* = *C. rufilabris* (see Bickley & MacLeod, 1956:194); *C. medialis* = *C. rufilabris*, *C. interrupta* = *C. rufilabris* (see Bram & Bickley, 1963:6); lectotype designated for *C. attenuata* (see Kimmings, 1940:447).

DISTRIBUTION.—CAN: NB<sup>4</sup>, ON<sup>6</sup> USA: DC<sup>4</sup>, FL<sup>2</sup>, GA<sup>2</sup>, IN<sup>7</sup>, KS<sup>5</sup>, LA<sup>4</sup>, MD<sup>12</sup>, MI<sup>4</sup>, MN<sup>11</sup>, MO<sup>10</sup>, NC<sup>8</sup>, NJ<sup>3</sup>, NY<sup>3</sup>, OH<sup>4</sup>, PA<sup>1</sup>, TN<sup>9</sup>, TX<sup>14</sup>, VA<sup>4</sup>, WS<sup>13</sup> (Schneider, 1851:79; <sup>2</sup>Walker, 1853:242, 244; <sup>3</sup>Smith, 1900:55; <sup>4</sup>Banks, 1903a:152; <sup>5</sup>Smith, 1925:168; <sup>6</sup>Smith, 1932:597; <sup>7</sup>Montgomery & Trippel, 1933:260; <sup>8</sup>Brimley, 1938:30; <sup>9</sup>Bickley, 1941:189; <sup>10</sup>Froeschner, 1947:134; <sup>11</sup>Parfin, 1952:425; <sup>12</sup>Bram & Bickley, 1963:5; <sup>13</sup>Throne, 1971a:74; <sup>14</sup>Agnew et al., 1981:5).

### **Chrysopodes (Neosuarius) Adams and Penny, 1987:435**

#### **Chrysopodes (Neosuarius) collaris** (Schneider, 1851:80 [*Chrysopa*]).

Holotype (abdomen missing): Puerto Rico (HMB).

= *Chrysopa thoracica* Walker, 1853:243. Holotype (abdomen missing): Dominican Republic, Santo Domingo, M. A. Pierret (BMNH).

= *Chrysopa krugii* Kolbe, 1888:173. Type(s) (sex unknown): Puerto Rico (HMB).

= *Chrysopa signatalis* Banks, 1911:342. Syntype females: USA, Texas, Brownsville, VI, Schaeffer (MCZ, USNM).

= *Chrysopa rufolinea* Banks, 1914a:24. Holotype male: Colombia, Cali, 1000m, Fassl (MCZ).

= *Chrysopa acolhua* Banks, 1948:156. Holotype male: Mexico, Guerrero, Pungarabato, 1260 m, 22.VIII.1930, José Paner (MCZ).

TAXONOMY.—*C. thoracica* = *C. collaris*, *C. krugii* = *C. collaris*, *C. rufolinea* = *C. collaris*, *C. signatalis* = *C. collaris*, *C. acolhua* = *C. collaris* (see Adams & Penny, 1987:436).

DISTRIBUTION.—USA: FL<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Banks, 1911:342; <sup>2</sup>Banks, 1938b:121, as *C. thoracica*).

### **Eremochrysa (Chrysopiella) Banks, 1911:344**

#### **Eremochrysa (Chrysopiella) brevisetosa** (Adams & Garland), 1981:1 [*Chrysopiella*].

Holotype male: USA, Utah, Uintah Co., 22.V.1954, M. Cazier (AMNH).

TAXONOMY.—To *Eremochrysa (Chrysopiella)* (see Brooks & Barnard, 1990:272).

DISTRIBUTION.—CAN: AB, USA: ID, NV, OR, UT, WY (Adams & Garland, 1981:3).

#### **Eremochrysa (Chrysopiella) minora** (Banks), 1935:55 [*Chrysopiella*].

Holotype (sex unknown): USA, Oregon, Umatilla, 24.VI.1882, S. Henshaw (MCZ).

TAXONOMY.—To *Eremochrysa (Chrysopiella)* (see Brooks & Barnard, 1990:272).

DISTRIBUTION.—USA: CA<sup>2</sup>, CO<sup>2</sup>, OR<sup>1</sup> (<sup>1</sup>Banks, 1935:55; <sup>2</sup>Adams, NEW STATE RECORDS).

#### **Eremochrysa (Chrysopiella) pallida** (Banks), 1911:345 [*Chrysopiella*].

Holotype male: USA, New Mexico, Rincon, 31.VIII, Cockerell (MCZ).

TAXONOMY.—To *Eremochrysa (Chrysopiella)* (see Brooks & Barnard, 1990:272).

DISTRIBUTION.—AZ<sup>2</sup>, CA<sup>3</sup>, NM<sup>1</sup>, OR<sup>2</sup>, UT<sup>2</sup>, (<sup>1</sup>Banks, 1911:345; <sup>2</sup>Bickley & MacLeod, 1956:198; <sup>3</sup>Adams, NEW STATE RECORD).

#### **Eremochrysa (Chrysopiella) sabulosa** (Banks), 1897b:174 [*Chrysopa*].

Holotype female: USA, Colorado, Fort Collins, 1916 (MCZ).

TAXONOMY.—To *Chrysopiella* (see Banks, 1911:344); to *Eremochrysa (Chrysopiella)* (see Brooks & Barnard, 1990:272).

DISTRIBUTION.—CO<sup>1</sup>, KS<sup>3</sup>, NM<sup>2</sup>, UT<sup>4</sup>, WY<sup>4</sup> (<sup>1</sup>Banks, 1897b:174; <sup>2</sup>Banks, 1903a:151;

<sup>3</sup>Smith, 1925:169; <sup>4</sup>Bickley & MacLeod, 1956:198).

REMARKS.—Bickley and MacLeod (1956:198) erroneously reported that Banks (1948) listed specimens from Texas.

**Eremochrysa (Eremochrysa)** Banks,  
1903a:158

**Eremochrysa (Eremochrysa) altilis** Banks, 1950:56.

Syntype male and females: USA, Arizona, Graham Co., Pinaleño Mtns., Stockton Pass, 5440 ft., 4.VIII.1948, W. Nutting and F. Werner (MCZ).

DISTRIBUTION.—AZ (Banks, 1950:57).

**Eremochrysa (Eremochrysa) californica** Banks, 1905b:6.

Holotype male: USA, California, Santa Clara Co. (MCZ).

DISTRIBUTION.—AZ<sup>2</sup>, CA<sup>1</sup> (<sup>1</sup>Banks, 1905b:6; <sup>2</sup>Banks, 1950:63).

**Eremochrysa (Eremochrysa) canadensis** (Banks), 1911:339 [*Chrysopa*].

Holotype female: Canada, Ontario, Lake Huron, Go Home Bay, 12.VII.1907 (MCZ).

TAXONOMY.—To *Eremochrysa* (see Banks, 1950:64).

DISTRIBUTION.—CAN: ON<sup>1</sup>, USA: MA<sup>2</sup>, ME<sup>2</sup>, NH<sup>2</sup>, WS<sup>3</sup> (<sup>1</sup>Banks, 1911:339; <sup>2</sup>Banks, 1950:66; <sup>3</sup>Throne, 1971a:76).

**Eremochrysa (Eremochrysa) fraterna** (Banks), 1897b:174 [*Chrysopa*].

Holotype female: USA, Colorado (MCZ).

TAXONOMY.—To *Eremochrysa* (see Banks, 1903a:159).

DISTRIBUTION.—CAN: BC<sup>4</sup>, USA: CO<sup>1</sup>, KS<sup>3</sup>, NV<sup>2</sup> (<sup>1</sup>Banks, 1897b:174; <sup>2</sup>Banks, 1903a:159; <sup>3</sup>Smith, 1925:169; <sup>4</sup>Smith, 1932:582).

**Eremochrysa (Eremochrysa) hageni** (Banks), 1903a:158 [*Chrysopa*].

Holotype female: USA, Texas, Austin, 20.V.1900 (MCZ).

DISTRIBUTION.—USA: AZ<sup>2</sup>, NM<sup>2</sup>, TX<sup>1</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1903a:159; <sup>2</sup>Banks, 1950:60). Procter (1946:44) recorded a specimen of *E. ha-geni* from Maine, but *E. canadensis* is the only species of *Eremochrysa* which is found in the northeastern U. S.

**Eremochrysa (Eremochrysa) pima** Banks, 1950:61.

Two syntypes (one female, one missing abdomen): USA, Arizona, White Mtns., South Fork Camp, 22.VI.1947, G. H. and J. L. Sperry (missing abdomen); New Mexico, Santa Fe, Cockerell (female) (MCZ). Note that MCZ records only indicate presence of paratype.

DISTRIBUTION.—USA: AZ, NM (Banks, 1950:62).

**Eremochrysa (Eremochrysa) pumilis** Banks, 1950:58.

Holotype male: USA, Colorado, Garland, VII, Yarrow (MCZ).

DISTRIBUTION.—USA: CA<sup>2</sup>, CO<sup>1</sup>, TX<sup>1</sup>, UT<sup>1</sup> (<sup>1</sup>Banks, 1950:58; <sup>2</sup>Bickley & MacLeod, 1956:200).

**Eremochrysa (Eremochrysa) punctinervis** (McLachlan), 1869:24 [*Chrysopa*].

Holotype female: USA, Texas, Bosque Co. (BMNH).

TAXONOMY.—To *Eremochrysa* (see Banks, 1903a:159).

DISTRIBUTION.—CAN: BC<sup>4</sup>, USA: AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>2</sup>, FL<sup>5</sup>, KS<sup>3</sup>, NM<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>McLachlan, 1869:24; <sup>2</sup>Banks, 1903a:159; <sup>3</sup>Smith, 1925:169; <sup>4</sup>Smith, 1932:583; <sup>5</sup>Banks, 1950:53).

**Eremochrysa (Eremochrysa) rufifrons** Banks, 1950:57.

Syntype series (both sexes): USA, Arizona, Globe, 9.VIII.1933, Parker; Arizona, Globe, 19.VII, F. Werner and W. Nutting (MCZ).

DISTRIBUTION.—USA: AZ (Banks, 1950:58).

**Eremochrysa (Eremochrysa) spilota** Banks, 1950:61.

Five syntype females: USA, California, Fort Yuma, 6–13.VI.1948, Andahl; California, Brawley, 8.IV, A. L. Melander (MCZ). Note that MCZ records only indicate presence of paratype.

DISTRIBUTION.—USA: CA (Banks, 1950:61).

**Eremochrysa (Eremochrysa) tibialis** Banks, 1950:55.

Syntype male: USA, Arizona, Florence Junction, 18.IV.1935, F. H. Parker; Utah, Wasont 22.VII, F. M. Carpenter; California, Vidal, 9.IV, Sperry (MCZ).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>1</sup>, NV<sup>2</sup>, UT<sup>1</sup> (<sup>1</sup>Banks, 1950:56; <sup>2</sup>Bickley & MacLeod, 1956:199).

**Eremochrysa (Eremochrysa) yosemite** Banks, 1950:63.

Holotype male: USA, California, Yosemite, 12.VI.1931, Essig (MCZ).

DISTRIBUTION.—USA: CA (Banks, 1950:64).

**Meleoma** Fitch, 1855:786

**Meleoma arizonensis** (Banks), 1903a:155 [*Chrysopa*].

Holotype female: USA, Arizona, Prescott (MCZ).

TAXONOMY.—To *Meleoma* (see Tauber, 1969:18).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NM<sup>3</sup>, TX<sup>3</sup>, UT<sup>3</sup> (<sup>1</sup>Banks, 1903a:155; <sup>2</sup>Bickley & MacLeod, 1956:195; <sup>3</sup>Tauber, 1969:18).

**Meleoma beardi** Tauber, 1969:20.

Holotype male: USA, Arizona, Chiricahua Mountains, Rustler's Park, 8,400 ft., 11.VI.1956, H. & A. Howden (CNC).

DISTRIBUTION.—USA: AZ (Tauber, 1969:21).

**Meleoma dolicharthra** (Navás), 1914e:96 [*Chrysopa*].

Holotype male: Guatemala: Capetillo, XI.1914, G. C. Champion (BMNH).

= *Meleoma cavifrons* Banks, 1950:46. Holotype male: USA, California, Tuolumne Co., Pinecrest, 10.VII.1948, P. H. Arnaud, Jr. (MCZ).

TAXONOMY.—To *Meleoma* (see Kimmings, 1940:448); *M. cavifrons* = *M. dolicharthra* (see Adams, 1969:6).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: AZ<sup>2</sup>, CA<sup>1</sup>, CO<sup>2</sup>, OR<sup>2</sup>, UT<sup>2</sup>, WA<sup>2</sup> (<sup>1</sup>Banks, 1950:46; <sup>2</sup>Tauber, 1969:36).

**Meleoma emuncta** (Fitch), 1855:792 [*Chrysopa*].

Holotype (sex unknown): USA, New York (MCZ).

= *Meleoma slossonae* Banks, 1896:95. Holotype male: USA, New Hampshire, Mount Washington (MCZ).

= *Meleoma verticalis* Banks, 1908a:259. Holotype male: USA, Colorado, Golden, Chimney Gulch, 3.VIII.1907, Oslar (MCZ).

= *Meleoma comata* Banks, 1950:45. Holotype male: USA, California, San Bernardino Co., Upper Santa Ana River, 15.VII.1948, G. H. & J. L. Sperry (MCZ).

TAXONOMY.—To *Meleoma* (see Banks, 1924:432); *M. slossonae* = *M. emuncta* (see Banks, 1924:432); *M. verticalis* = *M. emuncta*, *M. comata* = *M. emuncta* (see Tauber, 1969:23).

DISTRIBUTION.—CAN: BC<sup>5</sup>, MB<sup>8</sup>, NB<sup>8</sup>, NF<sup>8</sup>, NS<sup>8</sup>, ON<sup>5</sup>, PQ<sup>5</sup>, USA: AZ<sup>7</sup>, CA<sup>4</sup>, CO<sup>3</sup>, ID<sup>8</sup>, MA<sup>8</sup>, ME<sup>6</sup>, MI<sup>8</sup>, NC<sup>8</sup>, NH<sup>2</sup>, NM<sup>7</sup>, NV<sup>8</sup>, NY<sup>1</sup>, OR<sup>8</sup>, UT<sup>8</sup>, VA<sup>7</sup>, VT<sup>8</sup>, WA<sup>8</sup>, WS<sup>1</sup> (Fitch, 1855:792; <sup>2</sup>Banks, 1896:95; <sup>3</sup>Banks, 1908a:259; <sup>4</sup>Banks, 1950:45; <sup>5</sup>Smith, 1932:583; <sup>6</sup>Procter, 1946:44; <sup>7</sup>Bickley & MacLeod, 1956:185, 186; <sup>8</sup>Tauber, 1969:23).

**Meleoma furcata** (Banks), 1911:342 [*Chrysopa*].

Holotype female: USA, New Mexico, Fort Wingate, 19.VII (MCZ).

= *Meleoma delicata* Banks, 1950:48. Holotype male: USA, New Mexico, Fort Wingate, 14.VIII.1908, J. Woodgate (MCZ).

TAXONOMY.—To *Meleoma* (see Tauber, 1969:40); *M. delicata* = *M. furcata* (see Tauber, 1969:40).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>2</sup>, NM<sup>1</sup>, UT<sup>3</sup> (<sup>1</sup>Banks, 1911:342; <sup>2</sup>Bickley & MacLeod, 1956:186, 196; <sup>3</sup>Tauber, 1969:41).

**Meleoma hageni** Banks, 1948:170.

Holotype male: Mexico, Distrito Federal, Mexico City, Lomas de Chapultepec, 25.VII.1939, A. Dampf (MCZ).

DISTRIBUTION.—USA: AZ, NM (Tauber, 1969:38).

**Meleoma kennethi** Tauber, 1969:42.

Holotype male: USA, Nevada, Clark Co., 38 miles northwest of Las Vegas, Lee Canyon, 6,900 ft., 29.VII.1966, F., P., and M. Rindge (AMNH).

DISTRIBUTION.—USA: CA, NV (Tauber, 1969:42).

**Meleoma pallida** Banks, 1908a:260.

Holotype male: USA, Arizona, Huachuca Mountains, 8,000 ft., 20.VIII.1905, Oslar (MCZ).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup> (<sup>1</sup>Banks, 1908a:260; <sup>2</sup>Bickley & MacLeod, 1956:186).

**Meleoma pinaleana** (Banks), 1950:49 [*Chrysopa*].

Holotype (abdomen missing): USA, Arizona, Pinale-Globe, 18.VII.1948, F. Werner & W. Nutting (MCZ).

TAXONOMY.—To *Meleoma* (see Tauber, 1969:13).

DISTRIBUTION.—USA: AZ<sup>1</sup>, NM<sup>2</sup> (<sup>1</sup>Banks, 1950:49; <sup>2</sup>Tauber, 1969:13).

**Meleoma schwarzi** (Banks), 1903a:146 [*Chrysopa*].

Holotype (abdomen missing): USA, New Mexico, Las Vegas Hot Springs, 5.VIII, Barber & Schwarz (USNM).

TAXONOMY.—To *Meleoma* (see Adams, 1962:178).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>2</sup>, NM<sup>1</sup>, NV<sup>2</sup>, OR<sup>2</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1903a:147; <sup>2</sup>Tauber, 1969:14).

**Meleoma signoretii** Fitch, 1855:786.

Type(s) (sex unknown): USA, Vermont, Rupert (depository unknown)

DISTRIBUTION.—CAN: BC<sup>3</sup>, MB<sup>9</sup>, NS<sup>9</sup>, ON<sup>3</sup>, PQ<sup>3</sup>, USA: DC<sup>8</sup>, IL<sup>9</sup>, IN<sup>10</sup>, MA<sup>9</sup>, MD<sup>8</sup>, ME<sup>6</sup>, MI<sup>9</sup>, MN<sup>7</sup>, NC<sup>4</sup>, NH<sup>8</sup>, NJ<sup>9</sup>, NY<sup>2</sup>, PA<sup>9</sup>, TN<sup>5</sup>, VA<sup>8</sup>, VT<sup>1</sup>, WS<sup>9</sup>, WV<sup>9</sup>, (<sup>1</sup>Fitch, 1855:786; <sup>2</sup>Leonard, 1928:40; <sup>3</sup>Smith, 1932:584; <sup>4</sup>Brimley, 1938:31; <sup>5</sup>Bickley, 1941:187; <sup>6</sup>Procter, 1946:44; <sup>7</sup>Parfin, 1952:424; <sup>8</sup>Bickley & MacLeod, 1956:185; <sup>9</sup>Tauber, 1969:33; <sup>10</sup>Lawson & McCafferty, 1984:130).

**Meleoma tezcucana** (Banks), 1948:157 [Chrysopa].

Holotype female: Mexico, Distrito Federal, Lomas de Chapaltepec, Mexico City, 8.VIII.1940, A. Dampf (MCZ).

TAXONOMY.—To *Meleoma* (see Adams, 1969:8).

DISTRIBUTION.—USA: AZ (Adams, 1969:8).

**Nineta** Navás, 1912b:98

**Nineta gravida** (Banks), 1911:343 [Chrysopa].

Type male: USA, California, Yosemite (MCZ).

TAXONOMY.—to *Nineta* (see Brooks & Barnard, 1990:275).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: CA<sup>1</sup> (<sup>1</sup>Banks, 1911:343; <sup>2</sup>Smith, 1932:597).

**Nineta nanina** (Banks), 1911:344 [Chrysopa].

Type(s) (sex unknown): USA, Arizona, Palmerlee (MCZ).

TAXONOMY.—To *Nineta* (see Brooks & Barnard, 1990:275).

DISTRIBUTION.—USA: AZ<sup>1</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1911:340; <sup>2</sup>Adams, NEW STATE RECORD).

**Plesiochrysa** Adams, 1982b:28

**Plesiochrysa brasiliensis** (Schneider), 1851: 83 [Chrysopa].

Holotype (female): Brazil (HMB).

= *Chrysopa cubana* Navás, 1921:120. A junior primary homonym of *Chrysopa cubana* Hagen, 1861:215.

= *Chrysopa antillana* Navás, 1922c:392. A replacement name for *Chrysopa cubana* Navás. Lectotype male: Cuba, Havana, F. Cervera (MCZ).

= *Chrysopa bouvieri* Navás, 1924c:112. Holotype female: Costa Rica, Paul Serre (MNHN).

= *Chrysopa rata* Lacroix, 1926:68. Syntype series (one male, one female, one without abdomen): Brazil, Minas Gerais, Uberaba (MNHN).

= *Chrysopa uribei* Navás, 1927b:5. Holotype male: Colombia, Cundinamarca, San Antonio de Tena, San Pedro Claver (MZB, specimen not found).

= *Chrysopa scalaris* Navás, 1929c:26. Holotype male: Colombia, Behn vend., 3.XII.1900 (Hamburg, specimen destroyed).

= *Chrysopa yucatanensis* Navás, 1929c:28. Holotype male: Mexico, Yucatan, Progreso, 5.V.1905, W. Schwinghamer (Hamburg, specimen destroyed).

= *Chrysopa divergens* Navás, 1931:82. Holotype (sex unknown): Costa Rica, San José, 8 km wsw San José, Farm La Caja, V.1925 (Hamburg, specimen destroyed).

TAXONOMY.—To *Plesiochrysa*; *C. bouvieri* = *C. brasiliensis*, *C. antillana* = *C. brasiliensis*, *C. rata* = *C. brasiliensis*, *C. uribei* = *C. brasiliensis*, *C. scalaris* = *C. brasiliensis*, *C. yucatanensis* = *C. brasiliensis*, *C. divergens* = *C. brasiliensis* (see Adams, 1982:29).

DISTRIBUTION.—USA: FL (Adams, NEW NATIONAL RECORD).

**Pseudomallada** Tsukaguchi, 1995:67

**Pseudomallada luctuosus** (Banks), 1911:343 [Chrysopa].

Holotype male: USA, New Mexico, Fort Winnebago, 26.VI (MCZ).

TAXONOMY.—To *Mallada* (see Adams, 1975:172); to *Pseudomallada* by Adams, NEW COMBINATION.

DISTRIBUTION.—USA: AZ<sup>2</sup>, CO<sup>2</sup>, IN<sup>3</sup>, NE<sup>2</sup>, NM<sup>1</sup> (<sup>1</sup>Banks, 1911:343; <sup>2</sup>Adams & Garland, 1982:246; <sup>3</sup>Lawson & McCafferty, 1984:130).

**Pseudomallada macleodi** (Adams & Garland), 1982:240 [*Mallada*].

Holotype male: USA, Texas, Erath Co., Stephenville, 20.IV.1981, C. W. Agnew (MCZ).

TAXONOMY.—To *Pseudomallada* by Adams, **NEW COMBINATION**.

DISTRIBUTION.—CAN: ON, USA: AZ, KS, MD, TX (Adams & Garland, 1982:242).

**Pseudomallada perfectus** (Banks), 1895a:516  
[*Chrysopa*].

Holotype female: Mexico, Baja California Sur, Serra El Taste (MCZ).

= *Chrysopa cockerelli* Banks, 1903a:154.

Holotype female: USA, New Mexico, East Las Vegas (MCZ).

= *Chrysopa injusta* Banks, 1906a:98. Holotype female: USA, California, mountains near Claremont, Baker (MCZ).

= *Chrysopa marginalis* Banks, 1905b:5 not Navás 1905:122. Holotype (sex unknown): USA, California, mountains near Claremont, Baker (MCZ, specimen not found).

TAXONOMY.—To *Mallada* (see Adams, 1975:172); *C. cockerelli* = *M. perfectus*, *C. injusta* = *M. perfectus* (see Adams & Garland, 1982:244); to *Pseudomallada* by Adams, **NEW COMBINATION**.

DISTRIBUTION.—CAN: BC<sup>4</sup>, USA: AZ<sup>5</sup>, CA<sup>2</sup>, CO<sup>5</sup>, KS<sup>3</sup>, NM<sup>1</sup>, OR<sup>5</sup>, UT<sup>5</sup>, WA<sup>5</sup>, WY<sup>5</sup> (<sup>1</sup>Banks, 1903a:154; <sup>2</sup>Banks, 1906a:98; <sup>3</sup>Smith, 1925:168, as *C. cockerelli*; <sup>4</sup>Smith, 1932:597, as *C. cockerelli*; <sup>5</sup>Adams & Garland, 1982:244).

REMARKS.—The Navás name of *Chrysopa marginalis* was published in June, 1905 while the *Chrysopa marginalis* of Banks was published in November, 1905.

**Pseudomallada sierra** (Banks), 1924:431  
[*Chrysopa*].

Holotype female: USA, California, Los Angeles Co., San Gabriel Mtns., Sister Elsie Peak, 10.VI, F. Grinnell (MCZ).

TAXONOMY.—To *Mallada* (see Adams & Garland, 1982:245); to *Pseudomallada* by Adams, **NEW COMBINATION**.

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, OR<sup>2</sup>, WA<sup>2</sup> (<sup>1</sup>Banks, 1924:431; <sup>2</sup>Adams & Garland, 1982:245).

### **Yumachrysa** Banks, 1950:51

**Yumachrysa apache** (Banks), 1938b:120  
[*Chrysopa*].

Holotype female: USA, Arizona, Globe, 27.VI.1933, Parker (MCZ).

TAXONOMY.—To *Chrysopa* (*Yumachrysa*) (see Bickley & MacLeod, 1956:189); to *Suarius* (*Prochrysopa*) (see Tauber, 1975:696), to *Yumachrysa* (see Brooks & Barnard, 1990:276).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>3</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1938b:121; <sup>2</sup>Bickley & MacLeod, 1956:189; <sup>3</sup>Tauber, 1975:697).

**Yumachrysa clarivena** (Banks), 1950:50  
[*Chrysopa* (*Yumachrysa*)].

Holotype female: USA, Arizona, Yuma Co., Ehrenberg, 11.VII.1948, W. Nutting and F. Werner (MCZ).

TAXONOMY.—*Yumachrysa* elevated to generic status (see Brooks & Barnard, 1990:241).

DISTRIBUTION.—USA: AZ (Banks, 1950:50).

**Yumachrysa yuma** (Banks), 1950:49  
[*Chrysopa* (*Yumachrysa*)].

Two syntypes (male and female): USA, California, Fort Yuma, 6.VI and 13.VI, Andahl (MCZ).

TAXONOMY.—To *Suarius* (*Prochrysopa*) (see Tauber, 1975:697); to *Yumachrysa* (see Brooks & Barnard, 1990:241).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup> (<sup>1</sup>Banks, 1950:50; <sup>2</sup>Tauber, 1975:698).

NOTE.—**Leucochrysa californica** Navás, 1928a:35 was described from California, and the type specimen is present in the Stockholm Museum. Adams, 1977:94 has subsequently shown that this species is a synonym of *Leucochrysa colombia* Banks, 1910:150, known only from Colombia. The locality label associated with the type specimen of *L. californica* dates from the last century and may have originally been noted as "Cali" or "Col."

### CONIOPTERYGIDAE

Coniopterygidae, or dusky-wings, comprise the smallest and some of the most highly modified lacewings. Adults are generally only 2–3 mm long, with reduced wing venation and a white powder covering the wings and body. Because of this white powder, they often are mistaken for white-flies (Aleyrodidae) in flight. They are most active near dawn and dusk, fluttering slowly between plants, where they lay eggs and feed on soft-bodied insects. Eggs are usually laid singly on bark or leaves. Withycombe (1923) indicated that a larval dusky-wing will consume between 150 and 300 aphid eggs.

or young nymphs before metamorphosing to a pupa. Larvae and adults can feed on a wide variety of sluggish, soft-bodied insects, like aphids, white-flies, mealy-bugs, scale insects, etc. This makes them one of our more important biological control agents, although because of their size, they are often overlooked in the field. Although a wide variety of insects are consumed by the family in general, many species are closely associated with a specific species of tree or bush, indicating a close association with a specific type of host insect. The taxonomy of Coniopterygidae has been covered in a series of monographs and papers by Meinander (1972, 1974, 1975, 1990) and Johnson (1976, 1981a, 1981b).

## ALEUROPTERYGINAE

**Aleuropteryx** Löw, 1885:79**Aleuropteryx arceuthobii** Meinander, 1975:28.

Holotype male: USA, Colorado (USNM).

DISTRIBUTION.—USA: CO<sup>1</sup>, TX<sup>2</sup> (<sup>1</sup>Meinander, 1975:28; <sup>2</sup>Johnson, 1981a:272).

**Aleuropteryx arizonica** Johnson, 1981a:272.

Holotype male: USA, Arizona, Pinal Co., near Superior, Boyce Thompson Southwestern Arboretum, 11.VII.1949, B. W. Benson (INHS).

DISTRIBUTION.—USA: AZ (Johnson, 1981a:274).

**Aleuropteryx cupressi** Meinander, 1974:218.

Holotype male: USA, California, Marin Co., emerged ex *Cupressus goveniana*, 2.V.1961, J. Powell (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1974:218).

**Aleuropteryx dragoonica** Johnson, 1981a:275.

Holotype male: USA, Arizona, Cochise Stronghold, Dragoon Mountains, 15–18.VIII.1969, R. J. Shaw (USNM).

DISTRIBUTION.—USA: AZ (Johnson, 1981a:275).

**Aleuropteryx juniperi** Ohm, 1968:14.

Holotype male: Germany, Treisbach (Ohm Collection).

DISTRIBUTION.—USA: PA<sup>1</sup>, VA<sup>2</sup> (<sup>1</sup>Henry, 1974:643; <sup>2</sup>Flint, 1974:703).

**Aleuropteryx knowltoni** Johnson, 1981a:278.

Holotype male: USA, Utah, Logan, 29.VI.1950, G. F. Knowlton (USNM).

DISTRIBUTION.—USA: UT (Johnson, 1981a:279).

**Aleuropteryx longipennis** Meinander, 1974:218.

Holotype male: Mexico, Baja California Norte, 1 mile north of Meling Ranch, 2100 ft., 17.III.1972, J. Doyen & J. Powell (CAS).

DISTRIBUTION.—USA: CA<sup>1</sup>, NM<sup>2</sup>, UT<sup>2</sup> (<sup>1</sup>Meinander, 1974:219; <sup>2</sup>Johnson, 1981a:281).

**Aleuropteryx maculipennis** Meinander, 1972:45.

Holotype male: USA, California, Kern Co., 19 miles north northeast of Mojave, 14.VI.1962, C. MacNeill, D. Rentz & R. Brown (CAS).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, NM<sup>3</sup>, TX<sup>2</sup> (<sup>1</sup>Meinander, 1972:45; <sup>2</sup>Johnson, 1981a:282; <sup>3</sup>Meinander, 1990:7).

**Aleuropteryx megacornis** Johnson, 1981a:284.

Holotype male: USA, Arizona, Cochise Stronghold, Dragoon Mountains, 9–12.V.1970, R. J. Shaw (USNM).

DISTRIBUTION.—USA: AZ, NM, TX (Johnson, 1981a:284).

**Aleuropteryx punctata** Meinander, 1974:220.

Holotype male: Mexico, Sinaloa, 20 mi W Rosario, 30.I.1964, E. Schlinger (CAS).

DISTRIBUTION.—AZ<sup>1</sup>, NM<sup>2</sup> (<sup>1</sup>Meinander, 1974:220; <sup>2</sup>Johnson, 1981a:287).

**Aleuropteryx simillima** Meinander, 1972:46.

Holotype male: Mexico, Baja California, San Felipe, 7.III.1963, P. H. Arnaud (CAS).

DISTRIBUTION.—USA: AZ<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Wheeler, 1980:51; <sup>2</sup>Johnson, 1981a:289).

**Aleuropteryx unicolor** Meinander, 1972:48.

Holotype male: USA, California, Riverside Co., Palm Desert, 4 miles south of Boyd Desert Research Center, 11.IV.1963, W. A. Steffan (CAS).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup> (<sup>1</sup>Meinander, 1972:49; <sup>2</sup>Johnson, 1981a:290).

**Aleuropteryx vulgaris** Meinander, 1972:51.

Holotype male: USA, Texas, Kerrville, VII.1954, L. J. Bottimer (USNM).

DISTRIBUTION.—USA: AZ, CA, NM, TX, UT (Meinander, 1972:51).

**Aleuropteryx wernerii** Johnson, 1981a:294.

Holotype male: USA, New Mexico, Las Cruces, 28.VII–3.VIII.1975, W. P. Morrison (USNM).

DISTRIBUTION.—USA: NM (Johnson, 1981a:294).

**Bidesmida** Johnson, 1976:192

**Bidesmida morrisoni** Johnson, 1976:193.

Holotype male: USA, New Mexico, Las Cruces, 30.XI.1975, W. P. Morrison (USNM).

DISTRIBUTION.—USA: NM (Johnson, 1976:294).

**Helicoconis** Enderlein, 1905b:226

**Helicoconis (Helicoconis) californica** Meinander, 1972:119.

Holotype male: USA, California, Mono Co., 1 mile southwest of Tom's Place, 10.VIII.1963, M. J. Tauber & C. A. Toschi (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1972:119).

**Helicoconis (Helicoconis) lutea** (Wallengren), 1871:55 [*Coniopteryx*].

Type lost: Sweden, Gotland (NRS).

TAXONOMY.—To *Aleuropteryx* (see Klapálek, 1894:121); to *Helicoconis* (see Enderlein, 1905b:226).

DISTRIBUTION.—USA: PA (Meinander, 1972:124).

**Helicoconis (Helicoconis) similis** Meinander, 1972:124.

Holotype male: Canada, British Columbia, Top of Moyle Mountains., 6868 ft., East Kootenaya, H. B. Leech (CAS).

DISTRIBUTION.—CAN: BC (Meinander, 1972:124).

**Helicoconis (Helicoconis) walshi** (Banks), 1906b:83 [*Aleuropteryx*].

Holotype male: USA, Michigan, Michigan Agricultural College, 3.VII.1896 (MCZ).

TAXONOMY.—To *Helicoconis* (see Enderlein, 1908:16); to *Cryptoscenea* (see Enderlein, 1930:113); returned to *Helicoconis* (see Meinander, 1972:125).

DISTRIBUTION.—USA: CA<sup>3</sup>, IN<sup>4</sup>, ME<sup>2</sup>, MI<sup>1</sup>, WS<sup>2</sup> (Banks, 1906:83; Meinander, 1972:125; Meinander, 1974:223; <sup>4</sup>Lawson & McCafferty, 1984:130).

**Neoconis** Enderlein, 1930:112

**Neoconis bifurcata** Meinander, 1974:223.

Holotype male: USA, Arizona, Cochise Co., Stewart Camp, 1 mile south of Portal, 12–15.VIII.1971, J. Doyen (CAS).

DISTRIBUTION.—USA: AZ (Meinander, 1974:223).

**Neoconis inexpectata** Meinander, 1972:155.

Holotype male: USA, Arizona, Cochise Co., Southwest Research Station, 5400 ft., Chiracahua Mts., 58.VII.1964, D. R. Davis (MZB).

DISTRIBUTION.—USA: AZ (Meinander, 1972:155).

**Neoconis marginata** Meinander, 1972:156.

Holotype male: USA, Texas, Kerrville, 13.VIII.1954, L. J. Bottimer (USNM).

DISTRIBUTION.—USA: AZ, CA, TX (Meinander, 1972:157).

#### CONIOPTERYGINAE

**Coniopteryx (Coniopteryx)** Curtis, 1834:

unpaginated letterpress to plate 528

**Coniopteryx (Coniopteryx) californica** Meinander, 1974:225.

Holotype male: USA, California, San Luis Obispo Co., Oro Haco Lake, 5.VIII.1962, E. Schlinger (CAS).

DISTRIBUTION.—USA: CA<sup>1</sup>, TX<sup>2</sup> (<sup>1</sup>Meinander, 1974:226; <sup>2</sup>Meinander, 1990:61).

**Coniopteryx (Coniopteryx) dorsicornis** Johnson, 1981b:183.

Holotype male: USA, Florida, Highlands Co., Archbold Biological Station, 30.III–5.IV.1967, S. W. Frost (PSU).

DISTRIBUTION.—USA: FL (Johnson, 1981b:183).

**Coniopteryx (Coniopteryx) fitchi** Banks, 1895b:315.

Type lost: USA, Colorado (MCZ).

= *Malacomyza farinosa* Banks, 1906b:85.

Type lost: USA, California, San Mateo Co. (MCZ).

TAXONOMY.—To *Malacomyza* (see Banks, 1906b:84); to *Semidalis* (see Enderlein, 1907:12); returned to *Coniopteryx* (see Meinander, 1972:256); *M. farinosa* = *C. fitchi* (see Meinander, 1972:256).

DISTRIBUTION.—USA: AZ, CA, NV, TX (Meinander, 1972:257).

**Coniopteryx (Coniopteryx) forcipata** Johnson, 1981b:185.

Holotype male: USA, Arizona, Cochise Stronghold, Dragoon Mountains, 20–24.VI.1970, R. J. Shaw (USNM).

DISTRIBUTION.—USA: AZ (Johnson, 1981b:185).

**Coniopteryx (Coniopteryx) latipalpis** Meinander, 1972:257.

Holotype male: USA, Colorado, Junction Creek Road, 10,000 ft., 12–17.VII.1968, E. C. Becker (CNC).

DISTRIBUTION.—USA: AZ, CA, CO (Meinander, 1972:259).

**Coniopteryx (Coniopteryx) mexicana** Meinander, 1974:226.

Holotype male: Mexico, Sinaloa, 20 miles east of Villa Union, 31.I.1964, E. Schlänger & M. Irwin (CAS).

DISTRIBUTION.—USA: TX (Meinander, 1990:62).

**Coniopteryx (Coniopteryx) minuta** Meinander, 1972:259.

Holotype male: USA, California, San Luis Obispo Co., Morro Bay 8 miles, 2.V.1962, C. A. Toschi (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1972:260).

**Coniopteryx (Coniopteryx) palpalis** Meinander, 1972:260.

Holotype male: Mexico, Mexico, Route 15, km 152 west of Bosencheve, 20.VII.1966, Flint & Ortiz (USNM).

DISTRIBUTION.—USA: CA (Meinander, 1990:62).

**Coniopteryx (Coniopteryx) quadricephala** Johnson, 1981b:188.

Holotype male: USA, Utah, Cache Co., Green Canyon, 25–30.VII.1968, W. J. Hanson (USNM)

DISTRIBUTION.—USA: UT (Johnson, 1981b:188).

**Coniopteryx (Coniopteryx) simplex** Meinander, 1974:228.

Holotype male: USA, California, El Dorado Co., Pino Grande, southwest of Lake Edson, 8.VII.1967, W. J. Turner (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1974:228).

**Coniopteryx (Coniopteryx) simplicior** Meinander, 1972:261.

Holotype male: USA, Texas, Kerr Co., Kerrville, VII.1953, L. J. Bottimer (USNM).

DISTRIBUTION.—USA: AR<sup>2</sup>, CA<sup>2</sup>, FL<sup>1</sup>, GA<sup>1</sup>, IN<sup>3</sup>, MD<sup>1</sup>, NY<sup>1</sup>, TN<sup>1</sup>, TX<sup>1</sup>, VA<sup>1</sup> (Meinander, 1972:262; Meinander, 1974:64; Lawson & McCafferty, 1984:130).

**Coniopteryx (Coniopteryx) tineiformis** Curtis, 1834: text to plate 528.

Lectotype male: England (NMV).

= *Aleyrodes dubia* Stephens, 1829:367(nomen nudum).

= *Malacomyza lactea* Wesmael, 1836:166.

Type (sex unknown): (locality unknown) (IRSNB).

= *Sciodus lacteus* Zetterstedt, 1840:1051.

Type (sex unknown): Sweden, Bohuslän (NRS).

= *Sciodus fuscus* Zetterstedt, 1840:1051. Type (sex unknown): Sweden, Gotland (NRS).

= *Coniopteryx tineiformis* var. *xaveriana* Navás, 1918:20. Holotype female: Spain, Javier (MZB).

TAXONOMY.—Originally described from a mixed syntype series of seven specimens, with the text describing one species, and associated plate describing another. Meinander, 1972:254 has corrected this problem by designating a lectotype. *Aleyrodes dubia* is a nomen nudum, with no description and no type designation. *A. dubia* = *C. tineiformis*, *M. lactea* = *C. tineiformis* (see Enderlein, 1905b:197); *S. lacteus* = *C. tineiformis*, *S. fuscus* = *C. tineiformis*, *S. albus* = *C. tineiformis*, *C. t. xaverina* = *C. tineiformis* (see Meinander, 1972:256).

DISTRIBUTION.—CAN: BC<sup>3</sup>, MB<sup>4</sup>, NF<sup>1</sup>, PQ<sup>1</sup>, YU<sup>3</sup>, USA: AK<sup>1</sup>, CA<sup>1</sup>, ID<sup>1</sup>, IN<sup>2</sup>, MI<sup>1</sup>, TN<sup>1</sup>, VA<sup>1</sup> (<sup>1</sup>Meinander, 1972:254; <sup>2</sup>Lawson & McCafferty, 1984:130; <sup>3</sup>Meinander, 1990:61; <sup>4</sup>Penny, NEW STATE RECORD).

**Coniopteryx (Coniopteryx) westwoodii** (Fitch), 1855:802 [*Aleuronia*].

Type lost: USA, New York

= *Malacomyza ventralis* Navás, 1912a:198.

Type (very deteriorated): USA, New York, White Plains (MZB).

= (?) *Coniopteryx gidae* Procter, 1938:44 (nomen nudum).

TAXONOMY.—No description or type were designated for *C. gidae*. *M. ventralis* = *C. westwoodii* (see Meinander, 1972:263).

DISTRIBUTION.—CAN: MB<sup>8</sup>, USA: FL<sup>6</sup>, IN<sup>7</sup>, ME<sup>3</sup>, MI<sup>6</sup>, MN<sup>5</sup>, MO<sup>4</sup>, NJ<sup>2</sup>, NY<sup>1</sup>, TN<sup>6</sup>, TX<sup>6</sup>, VA<sup>6</sup>, WS<sup>6</sup> (<sup>1</sup>Fitch, 1855:98; <sup>2</sup>Smith, 1900:54; <sup>3</sup>Procter, 1946:44, as *C. gidae*; <sup>4</sup>Froeschner, 1947:135; <sup>5</sup>Parfin, 1952:426; <sup>6</sup>Meinander, 1972:263; <sup>7</sup>Lawson & McCafferty, 1984:130; <sup>8</sup>Penny, NEW STATE RECORD).

**Coniopteryx (Xeroconiopteryx)** Meinander, 1972:203

**Coniopteryx (Xeroconiopteryx) canadensis** Meinander, 1972:211.

Holotype male: Canada, Saskatchewan, Elbow, 10.VI.1960, A. R. Brooks (CNC).

DISTRIBUTION.—CAN: SA<sup>1</sup>, USA: AK<sup>2</sup>, WS<sup>1</sup> (<sup>1</sup>Meinander, 1972:212; <sup>2</sup>Meinander, 1990:40).

**Coniopteryx (Xeroconiopteryx) diversicornis** Meinander, 1972:213.

Holotype male: USA, Texas, Kerr Co., Kerrville, 13.X.1953, L. J. Bottimer (USNM).

DISTRIBUTION.—USA: AZ, CA, FL, TX (Meinander, 1972:214).

**Coniopteryx (Xeroconiopteryx) meinanderi** Johnson, 1981b:186.

Holotype male: USA, Arizona, Cochise Stronghold, Dragoon Mountains, R. J. Shaw (USNM).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup> (<sup>1</sup>Johnson, 1981b:186; <sup>2</sup>Meinander, 1990:40).

**Coniopteryx (Xeroconiopteryx) texana** Meinander, 1972:214.

Holotype male: USA, Texas, Kerr Co., Kerrville, 30.VI.1953, L. J. Bottimer (USNM).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Meinander, 1972:215; <sup>2</sup>Meinander, 1990:40).

**Conwentzia** Enderlein, 1905a:10

**Conwentzia barretti** (Banks), 1898:202  
[*Coniopteryx*].

Holotype female: Mexico, Amecameca (MCZ).

= *Parasemidalis flaviceps* Banks, 1906b:81. Lectotype female: USA, California, Los Angeles (MCZ).

TAXONOMY.—To *Semidalis* (see Enderlein, 1907:12); to *Conwentzia* (see Meinander, 1972:295); *P. flaviceps* = *C. barretti* (see Meinander, 1972:295).

DISTRIBUTION.—USA: CA (Fleschner & Ricker, 1953:458, as *Parasemidalis flaviceps*).

**Conwentzia californica** Meinander, 1972:297.

Holotype male: USA, California, Mono Co., Lee Vining Campground, west of Mono Lake, VII.1961, D. C. Rentz (CAS).

DISTRIBUTION.—CAN: BC<sup>2</sup> USA: AZ<sup>1</sup>, CA<sup>1</sup>, CO<sup>1</sup>, NM<sup>3</sup>, OR<sup>3</sup>, UT<sup>1</sup>, WA<sup>1</sup>, WY<sup>1</sup> (<sup>1</sup>Meinander, 1972:298; <sup>2</sup>Meinander, 1990:72; <sup>3</sup>Penny, NEW STATE RECORDS).

nander, 1972:298; <sup>2</sup>Meinander, 1990:72; <sup>3</sup>Penny, NEW STATE RECORDS).

**Conwentzia pineticola** Enderlein, 1905a:40.

Lectotype male: Berlin, Germany (HMB).

= *Conwentzia pineticola* var. *furcilla* Enderlein, 1906a:194. Two syntype males: Berlin, Germany (probably lost).

= *Conwentzia pineticola* var. *tetensi* Enderlein, 1906a:195. Lectotype female: Finland, Jakobstad (MZB).

= *Conwentzia hageni* Banks, 1906b:82. Lectotype female: USA, Virginia (MCZ).

= *Coniopteryx reticulata* Tullgren, 1906:14. Syntypes male and female: Sweden, Stockholm (depository unknown).

= *Conwentzia angulata* Navás, 1914b:16. Holotype female: USA, New York (MZB).

= *Conwentzia axillata* Navás, 1914b:17. Holotype male: USA, New York (MZB).

= *Conwentzia cryptoneurus* Bagnall, 1915:192. Syntype male, 2 females: England, Newcastle-upon-Tyne (depository unknown).

TAXONOMY.—*C. p.* var. *furcilla* = *C. pineticola*, *C. p.* var. *tetensi* = *C. pineticola*, *C. hageni* = *C. pineticola*, *C. angulata* = *C. pineticola*, *C. axillata* = *C. pineticola*; lectotypes designated for *C. pineticola* and *C. hageni* (see Meinander, 1972:301).

DISTRIBUTION.—CAN: MB<sup>9</sup>, NF<sup>6</sup>, NS<sup>6</sup>, ON<sup>6</sup>, USA: AR<sup>7</sup>, DC<sup>6</sup>, FL<sup>6</sup>, IA<sup>9</sup>, IN<sup>8</sup>, MA<sup>6</sup>, MD<sup>6</sup>, ME<sup>3</sup>, MI<sup>6</sup>, MN<sup>5</sup>, MO<sup>4</sup>, NC<sup>6</sup>, NH<sup>6</sup>, NJ<sup>6</sup>, NY<sup>2</sup>, OH<sup>6</sup>, PA<sup>6</sup>, VA<sup>1</sup>, WS<sup>6</sup> (<sup>1</sup>Banks, 1906b:82; <sup>2</sup>Navás, 1914b:16, as *C. angulata*; <sup>3</sup>Procter, 1946:44, as *C. hageni*; <sup>4</sup>Froeschner, 1947:135, as *C. hageni*; <sup>5</sup>Parfin, 1952:426, as *C. hageni*; <sup>6</sup>Meinander, 1972:303; Throne, 1972:126; <sup>8</sup>Lawson & McCafferty, 1984:130; <sup>9</sup>Penny, NEW STATE RECORDS).

**Conwentzia psociformis** (Curtis), 1834: plate 528 [*Coniopteryx*].

Type deposition unknown: England, specimen presumed lost.

= *Coniopteryx aphidiformis* Rambur, 1842:316. Type depository unknown.

TAXONOMY.—To *Coniortes* (see Walker, 1853:299); to *Conwentzia* (see Enderlein, 1905:10); *C. aphidiformis* = *C. psociformis* (see Meinander, 1972:305).

DISTRIBUTION.—CAN: BC, USA: NJ, OH (Meinander, 1972:306).

**Parasemidalis** Enderlein, 1905c:197

**Parasemidalis fuscipennis** (Reuter), 1894:13  
[*Coniopteryx*].

Holotype male: Finland (MZB).

= *Parasemidalis annae* Enderlein, 1905c:198.

Holotype female: Germany, Berlin (HMB).

TAXONOMY.—To *Parasemidalis* (see Enderlein, 1906:219); *P. annae* = *P. fuscipennis* (see Meinander, 1972:281).

DISTRIBUTION.—USA: AZ, CA, FL, MI (Meinander, 1972:283).

**Semidalis** Enderlein, 1905c:197

**Semidalis angusta** (Banks), 1906b:81 [*Coniopteryx*].

Lectotype male: USA, Arizona, Williams, 29.V. Barber & Schwarz (USNM).

TAXONOMY.—To *Semidalis* (see Enderlein, 1908:11); lectotype designated (see Meinander, 1972:327).

DISTRIBUTION.—USA: AR, AZ, CA, MT, TX (Meinander, 1972:327).

**Semidalis arnaudi** Meinander, 1972:328.

Holotype male: Mexico, Sonora, Alamos, 27.II.1963, P. H. Arnaud (CAS).

DISTRIBUTION.—USA: AZ (Meinander, 1990:78).

**Semidalis bituberculata** Meinander, 1990:79

Holotype male: USA, California, Riverside Co., hills on west side of Menifee Valley, 27.VII.1980, J. D. Pinto (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1990:79).

**Semidalis deserta** Meinander, 1974:230.

Holotype male: USA, California, Riverside Co., Boyd Desert Research Center, 3 miles south of Palm Desert, 24–26.VI.1969, S. Frommer & B. Worley (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1974:231).

**Semidalis flinti** Meinander, 1972:335.

Holotype male: USA, Texas, Kerr Co., Kerrville, 17.VIII.1954, L. J. Bottimer (USNM).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, TX<sup>1</sup> (Meinander, 1972:336; Meinander, 1974:231).

**Semidalis frommeri** Meinander, 1974:231.

Holotype male: USA, California, Riverside Co., Deep Canyon, 22.VII.1964, M. Irwin (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1974:232).

**Semidalis inconspicua** Meinander, 1972:336.

Holotype male: USA, Virginia, Falls Church, Holmes Run, 6.VI.1961, W. W. Wirth (USNM).

DISTRIBUTION.—USA: AR<sup>2</sup>, AZ<sup>1</sup>, CA<sup>1</sup>, IN<sup>3</sup>, MD<sup>1</sup>, OK<sup>1</sup>, TX<sup>1</sup>, VA<sup>1</sup>, WS<sup>1</sup> (Meinander, 1972:336; Meinander, 1974:232; Lawson & McCafferty, 1984:130).

**Semidalis tricornis** Johnson, 1981b:190.

Holotype male: USA, Arizona, Cochise Stronghold, Dragoon Mountains, R. J. Shaw (USNM).

DISTRIBUTION.—USA: AZ (Johnson, 1981b:191).

**Semidalis vicina** (Hagen), 1861:197 [*Coniopteryx*].

Holotype female: USA, Washington, D. C. (MCZ).

= *Parasemidalis nigriceps* Navás, 1918a:355.

Holotype female: USA, New York, White Plains, 18.VII.1915 (MZB).

= *Semidalis ribesi* Ohm, 1973:237. Holotype male: France, Pyrénées orientales, above the Bains du Boulou, 27.V.1963 (Ohm Collection).

TAXONOMY.—To *Semidalis* (see Enderlein, 1906:215); to *Niphetia* (see Enderlein, 1930:106); returned to *Semidalis* (see Procter, 1938:44); *P. nigriceps* = *S. vicina*, *S. ribesi* = *S. vicina* (Meinander, 1972:330).

DISTRIBUTION.—CAN: AB<sup>8</sup>, PQ<sup>8</sup>, USA: CT<sup>8</sup>, DC<sup>1</sup>, FL<sup>8</sup>, GA<sup>8</sup>, IN<sup>9</sup>, KS<sup>3</sup>, ME<sup>5</sup>, MO<sup>6</sup>, MN<sup>7</sup>, MA<sup>8</sup>, ME<sup>8</sup>, MI<sup>8</sup>, MS<sup>8</sup>, NC<sup>4</sup>, NH<sup>8</sup>, NY<sup>2</sup>, OH<sup>8</sup>, PA<sup>8</sup>, SC<sup>8</sup>, TN<sup>8</sup>, VA<sup>8</sup>, WS<sup>8</sup> (Hagen, 1861:197; Navás, 1918:19, as *P. nigriceps*; Smith, 1925:170; Brimley, 1938:32; Procter, 1946:44; Froeschner, 1947:135; Parfin, 1952:425; Meinander, 1972:331; Lawson & McCafferty, 1984:130).

**Semidalis wallacei** Meinander, 1972:337.

Holotype male: USA, Pennsylvania, Pittsburgh, 15.VII.1932, H. Kahl (CMNH).

DISTRIBUTION.—USA: PA (Meinander, 1972:338).

**Semidalis xerophila** Meinander, 1990:84.

Holotype male: USA, California, Riverside Co., south of Palm Desert, Boyd Desert Research Center, 19–21.VI.1978 (CAS).

DISTRIBUTION.—USA: CA (Meinander, 1990:84).

## DILARIDAE

The dilarids, or pleasing lacewings, comprise a small family of tropical, moth-like lacewings which are rarely seen in North America. This family has broad, darkly-banded wings with considerable long pilosity on the wings, giving them a "hairy" appearance. Males have distinctive pectinate antennae. However, unlike most moths, the eyes are very large in proportion to the rest of the head. Females bear ovipositors which are recurved over the abdomen. Larvae are very elongate, and live under the bark of dead trees, where they are presumed to feed on beetle adults or larvae (Gurney, 1947). Tropical American species appear to be most abundant in the forest canopy as adults, and seem to emerge most frequently during the driest part of the year (Penny and Arias, 1981). Only two species are known from the U. S., both in the genus *Nallachius* Navás, 1909a, and although one species is rather widely distributed over eastern North America, it is seldom collected. A key to New World species can be found in Adams (1970).

***Nallachius*** Navás, 1909a:666***Nallachius americanus*** (McLachlan), 1881:55  
[*Dilar*].

Holotype female: USA, Kentucky, Bee Spring, VI.1874, Sanborn (MCZ).

TAXONOMY.—To *Nallachius* (see Navás, 1909a:669).

DISTRIBUTION.—USA: GA<sup>5</sup>, IN<sup>4</sup>, KY<sup>1</sup>, MD<sup>2</sup>, SC<sup>5</sup>, TX<sup>3</sup>, VA<sup>2</sup> (<sup>1</sup>McLachlan, 1881:55; <sup>2</sup>Carpenter, 1940:274; <sup>3</sup>Stange, 1961b:144; <sup>4</sup>Lawson & McCafferty, 1984:130; <sup>5</sup>Hoffman, 1990:155).

***Nallachius pulchellus*** (Banks), 1938c:289 [*Dilar*].

Male and female syntypes: Cuba, Soledad (Cienfuegos), 4.V.1930, P. J. Darlington, Jr. (MCZ).

TAXONOMY.—To *Nallachius* (see Carpenter, 1940:274).

DISTRIBUTION.—USA: AZ (Carpenter, 1940:275).

## HEMEROBIIDAE

Hemerobiidae, or brown lacewings, comprise one of the more commonly encountered families of Neuroptera. Adults are predominantly brown

or black in coloration. They can be characterized as having the Rs and MA fused with R in the forewing to give the appearance of having multiple radial sectors. The number of these apparent radial sectors is important in recognizing individual genera. They are rather small insects, and individuals of the genus *Symppherobius* can have forewing length 2–3 mm. Adults are most abundant on trees and bushes, although some groups, most notably *Micromus*, can be found on grasses. Both adults and larvae feed on soft-bodied insects. The genus *Symppherobius* was revised by Oswald (1988), and he has dealt with the family level classification (Oswald, 1993). The Canadian species have been redescribed by Kevan and Klimaszewski (1986, 1987) and Klimaszewski and Kevan (1985, 1987a, 1987b, 1988a, 1988b, 1989, 1990a, 1990b).

## HEMEROBIINAE

***Hemerobius*** Linnaeus, 1758:549***Hemerobius alpestris*** Banks, 1908a:261.

Holotype male: USA, Colorado, Sugar Loaf Mountain, 13.V, S. A. Rohwer (MCZ).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CO<sup>1</sup> (<sup>1</sup>Banks, 1908a:261; <sup>2</sup>Carpenter, 1940:208).

***Hemerobius bistrigatus*** Currie, 1904a:79.

Holotype female: USA, California, Humboldt Co., Little River, 31.V.1903 (USNM).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: CA<sup>1</sup>, ID<sup>2</sup>, OR<sup>2</sup> (<sup>1</sup>Currie, 1904a:79; <sup>2</sup>Carpenter, 1940:208).

***Hemerobius conjunctus*** Fitch, 1855:798.

Holotype male: Kelson's Cabin, 7.VII (MCZ).

= *Hemerobius citrinus* Hagen, 1861:204.

Holotype (male): America Septentrionalis, Knob (HMB).

= *Hemerobius venustus* Banks, 1897a:25.

Holotype female: USA, New Hampshire, Mount Washington, N. Banks (MCZ).

= *Hemerobius cockerelli* Banks, 1901b:286.

Holotype male: USA, New Mexico, Las Vegas Range, 11000 ft., Cockerell (MCZ).

= *Hemerobius caudelli* Currie, 1904b:87.

Holotype female: Canada, British Columbia, London Hill Mine, Bear Lake, 7000 ft., 29.VII.1903 (USNM).

= *Hemerobius glacialis* Currie, 1904b:88.

Holotype female: Canada, British Columbia,

Kokanee Mountain, 9000 ft., 10.VIII.1903 (USNM).

**TAXONOMY.** — *H. venustus* = *H. conjunctus*, *H. caudelli* = *H. cockerelli* (see Banks, 1905c:49); *H. citrinus* = *H. conjunctus*, *H. glacialis* = *H. conjunctus*, *H. cockerelli* = *H. conjunctus* (see Carpenter, 1940:209). Carpenter, 1940:209 considered the single Fitch specimen of *H. conjunctus* at MCZ to be the holotype. Klimaszewski and Kevan (1985:40) rejected this point of view and designated this specimen as lectotype. As there is no evidence of additional specimens in the original type series, we agree with Carpenter and herein list it as holotype.

**DISTRIBUTION.** — CAN: AB<sup>6</sup>, BC<sup>3</sup>, LB<sup>6</sup>, MB<sup>10</sup>, NF<sup>6</sup>, NS<sup>6</sup>, ON<sup>6</sup>, PE<sup>6</sup>, PQ<sup>6</sup>, SK<sup>6</sup>, YT<sup>10</sup>, USA: AK<sup>6</sup>, CO<sup>6</sup>, IN<sup>9</sup>, ME<sup>6</sup>, MN<sup>7</sup>, MT<sup>6</sup>, NC<sup>5</sup>, NH<sup>1</sup>, NM<sup>2</sup>, NY<sup>4</sup>, TN<sup>6</sup>, VT<sup>6</sup>, WS<sup>8</sup>, WY<sup>6</sup> (<sup>1</sup>Banks, 1897a:25, as *H. venustus*; <sup>2</sup>Banks, 1901b:286, as *H. cockerelli*; <sup>3</sup>Currie, 1904b:87, as *H. caudelli*; <sup>4</sup>Leonard, 1928:39; <sup>5</sup>Brimley, 1938:29; <sup>6</sup>Carpenter, 1940:209; <sup>7</sup>Parfin, 1952:423; <sup>8</sup>Throne, 1971b:80; <sup>9</sup>Lawson & McCafferty, 1984:130; <sup>10</sup>Klimaszewski & Kevan, 1985:39).

**Hemerobius costalis** Carpenter, 1940:213.

Holotype male: USA, Maine, Eastport, 14.VII.1909, C. W. Johnson (MCZ).

**DISTRIBUTION.** — CAN: AB<sup>1</sup>, BC<sup>1</sup>, NS<sup>1</sup>, NT<sup>1</sup>, ON<sup>1</sup>, PQ<sup>1</sup>, SK<sup>3</sup>, YT<sup>2</sup>, USA: AK<sup>1</sup>, CO<sup>1</sup>, MA<sup>1</sup>, ME<sup>1</sup>, NH<sup>1</sup>, VT<sup>1</sup> (<sup>1</sup>Carpenter, 1940:213; <sup>2</sup>Klimaszewski & Kevan, 1985:69; <sup>3</sup>Klimaszewski & Kevan, 1989:205).

**Hemerobius discretus** Navás, 1917a:5.

Lectotype female: USA, New Mexico, Jemez Springs, 16.VI.1916 (MZB).

= *Hemerobius neglectus* Hagen, 1861:206.

Type(s) unknown: Mexico, Ehrenberg (ZMB).

= *Hemerobius mexicanus* Navás, 1921:123.

= *Hemerobius pallidulus* Kimmins, 1928:366.

= *Hemerobius neadelphus* Gurney, 1948:214.

Holotype male: Canada, British Columbia, Kaslo, 4.VII.1903, R. P. Currie (USNM).

**TAXONOMY.** — *H. discretus* = *H. pacificus* (see Carpenter, 1940:204). *H. discretus* not equal *H. pacificus*; *H. mexicanus* = *H. discretus*, *H. neglectus* = *H. discretus*, *H. pallidulus* = *H. discretus* (see Monserrat, 1996:416–417). *H. neadelphus* = *H. discretus* (Penny, NEW SYNONYMY).

**DISTRIBUTION.** — CAN: BC<sup>1</sup>, USA: AZ<sup>3</sup>, CA<sup>2</sup>, CO<sup>3</sup>, LA<sup>3</sup>, NM<sup>3</sup>, OK<sup>3</sup>, OR<sup>1</sup>, TX<sup>3</sup>, UT<sup>2</sup>

(<sup>1</sup>Gurney, 1948:214; <sup>2</sup>Klimaszewski & Kevan, 1985:25; <sup>3</sup>Monserrat, 1996:421).

**Hemerobius dorsatus** Banks, 1904a:61.

Lectotype male: USA, Colorado, Veta Pass, 1.VII (MCZ).

**TAXONOMY.** — Lectotype designated (see Carpenter, 1940:211).

**DISTRIBUTION.** — CAN: AB<sup>2</sup>, BC<sup>2</sup>, NT<sup>2</sup>, ON<sup>3</sup>, PQ<sup>3</sup>, SA<sup>2</sup>, YT<sup>3</sup>, USA: AK<sup>2</sup>, CO<sup>1</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1904a:61; <sup>2</sup>Carpenter, 1940:212; <sup>3</sup>Klimaszewski & Kevan, 1985:66).

**Hemerobius humulinus** Linnaeus, 1758:550.

Type lost.

= *Hemerobius crispus* Stephens, 1836:112. Type lost: England.

= *Hemerobius maculatus* Wesmael, 1841:215. Type(s) (unknown): Belgium, Brussels (IRSCB).

= *Hemerobius oblitteratus* Walker, 1853:289. Holotype male: USA, Georgia (BMNH).

= *Hemerobius castaneae* Fitch, 1855:798. Holotype female: USA (MCZ).

= *Hemerobius tutatrix* Fitch, 1855:798. Holotype male: USA (MCZ).

= *Hemerobius gossypii* Ashmead, 1895:27. Lectotype female: USA, Georgia (BMNH).

= *Hemerobius algonquinus* Banks, 1924:429. Lectotype female: USA, New Hampshire, Hampton, 6.III.1904, S. A. Shaw (MCZ).

= *Hemerobius obtusus* Nakahara, 1954:42. Holotype male: Japan, Mejiro, Tokyo (Nakahara Collection).

= *Hemerobius shikotanus* Kuwayama, 1956:77. Holotype female: Kurile Islands, Shikotan Island, Shakotan, 23–27.VI.1935, Y. Sugihara (Kuwayama Collection).

**TAXONOMY.** — *H. humuli* = *H. humulinus*, *H. maculatus* = *H. humulinus*, *H. castaneae* = *H. humulinus*, *H. tutatrix* = *H. humulinus*, *H. gossypii* = *H. humulinus*, *H. algonquinus* = *H. humulinus*, *H. oblitteratus* = *H. humulinus* (see Carpenter, 1940:201); *H. shikotanus* = *H. humulinus* (see Makarkin, 1985:105).

**DISTRIBUTION.** — CAN: AB<sup>12</sup>, BC<sup>6</sup>, MB<sup>6</sup>, NF<sup>6</sup>, NS<sup>6</sup>, ON<sup>6</sup>, PQ<sup>6</sup>, SK<sup>6</sup>, YT<sup>12</sup>, USA: AK<sup>10</sup>, CO<sup>6</sup>, DC<sup>3</sup>, FL<sup>6</sup>, GA<sup>1</sup>, IN<sup>4</sup>, KS<sup>3</sup>, MD<sup>13</sup>, ME<sup>7</sup>, MI<sup>3</sup>, MN<sup>9</sup>, MO<sup>8</sup>, NC<sup>5</sup>, ND<sup>6</sup>, NH<sup>3</sup>, NJ<sup>2</sup>, NY<sup>2</sup>, PA<sup>3</sup>, SD<sup>6</sup>, TN<sup>6</sup>, TX<sup>6</sup>, VA<sup>2</sup>, WA<sup>10</sup>, WS<sup>11</sup> (<sup>1</sup>Walker, 1853:289; <sup>2</sup>Smith, 1900:56, as *H. castanea* and *H. tutatrix*; <sup>3</sup>Banks, 1905c:32; <sup>4</sup>Montgomery & Trippel, 1933:259; <sup>5</sup>Brimley, 1938:29, as *H. humuli*; <sup>6</sup>Carpenter, 1940:202;

<sup>7</sup>Procter, 1946:43; <sup>8</sup>Froeschner, 1947:131; <sup>9</sup>Parfin, 1952:423; <sup>10</sup>Nakahara, 1966:212; <sup>11</sup>Throne, 1971b:80; <sup>12</sup>Klimaszewski & Kevan, 1985:56; <sup>13</sup>Monserrat, 1996:429).

**Hemerobius kokaneeanus** Currie, 1904b:86.

Holotype female: Canada, British Columbia, Kokanee Mountain, 9000 ft., 10.VIII.1903 (USNM).

= *Hemerobius hesperus* Banks, 1924:429.

Lectotype female (abdomen missing): USA, California, Tahoe, Angora Pk., 10.VII.1915, E. P. Van Duzee (MCZ).

TAXONOMY.—*H. hesperus* = *H. kokaneeanus* (see Carpenter, 1940:206); lectotype designated for *H. hesperus* (see Klimaszewski & Kevan, 1985:30).

DISTRIBUTION.—CAN: BC<sup>1</sup>, MB<sup>4</sup>, PQ<sup>2</sup>, YT<sup>3</sup>; USA: AK<sup>5</sup>, CA<sup>2</sup>, CO<sup>2</sup>, MT<sup>2</sup>, NV<sup>5</sup>, WA<sup>2</sup> (<sup>1</sup>Currie, 1904b:85; <sup>2</sup>Carpenter, 1940:206; <sup>3</sup>Klimaszewski & Kevan, 1985:32; <sup>4</sup>Klimaszewski & Kevan, 1989:205; <sup>5</sup>Penny, NEW STATE RECORDS).

**Hemerobius nigrans** Carpenter, 1940:207.

Holotype male: Canada, British Columbia, Kamloops, Mount Lobo, 2.VI.1938, J. K. Jacob (CNC).

DISTRIBUTION.—CAN: BC<sup>1</sup>, YT<sup>2</sup>; USA: CA<sup>1</sup>, CO<sup>1</sup>, WY<sup>1</sup> (<sup>1</sup>Carpenter, 1940:207; <sup>2</sup>Klimaszewski & Kevan, 1985:36).

**Hemerobius ovalis** Carpenter, 1940:205.

Holotype male: USA, California, Claremont, C. F. Baker (MCZ).

DISTRIBUTION.—CAN: AB<sup>1</sup>, BC<sup>1</sup>, NF<sup>2</sup>, PQ<sup>2</sup>, YT<sup>2</sup>; USA: AK<sup>1</sup>, CA<sup>1</sup>, OR<sup>1</sup>, UT<sup>1</sup>, WA<sup>1</sup>, WY<sup>1</sup>, (<sup>1</sup>Carpenter, 1940:205; <sup>2</sup>Klimaszewski & Kevan, 1985:28).

**Hemerobius pacificus** Banks, 1897a:24.

Lectotype male: USA, Washington, Olympia, T. Kincaid (MCZ).

= *Hemerobius pallescens* Currie, 1904a:80. Holotype female: USA, California, Humboldt Co., Fieldbrook, 30.V.1903 (USNM).

TAXONOMY.—*H. pallescens* = *H. pacificus* (see Banks, 1905c:49). Lectotype designated for *H. pacificus* (see Carpenter, 1940:204).

DISTRIBUTION.—CAN: AB<sup>4</sup>, BC<sup>4</sup>, MB<sup>5</sup>, SK<sup>4</sup>; USA: AK<sup>6</sup>, AZ<sup>4</sup>, CA<sup>2</sup>, CO<sup>4</sup>, ID<sup>4</sup>, IN<sup>5</sup>, MT<sup>8</sup>, NM<sup>3</sup>, NV<sup>7</sup>, OR<sup>4</sup>, TX<sup>4</sup>, UT<sup>8</sup>, WA<sup>1</sup> (<sup>1</sup>Banks, 1897a:24; <sup>2</sup>Currie, 1904a:80; <sup>3</sup>Navás, 1917a:5; <sup>4</sup>Carpenter, 1940:204; <sup>5</sup>Lawson & McCafferty, 1984:130; <sup>6</sup>Klimaszewski & Kevan, 1985:23; <sup>7</sup>

Monserrat, 1996:415; <sup>8</sup>Penny, NEW STATE RECORDS).

REMARKS.—The Indiana record is probably in error. However, one of us (NDP) has seen a series of specimens of the closely related *H. neadelphus* labelled from northwestern Minnesota.

**Hemerobius pinidumus** Fitch, 1855:799.

Holotype male: no locality data (MCZ).

= *Hemerobius hyalinatus* Fitch, 1855:799.

Holotype male: no locality data (MCZ).

= *Hemerobius canadensis* Banks, 1897a:26.

Holotype female: Canada, Quebec, Sherbrooke, October, L'Abbe, P. A. Bégin (MCZ).

= *Hemerobius kootenayensis* Currie, 1904b:88.

Holotype female: Canada, British Columbia, Kaslo, 17.VI.1903 (USNM).

TAXONOMY.—*H. canadensis* = *H. hyalinatus* (see Banks, 1905c:34); *H. kootenayensis* = *H. conjunctus* (see Carpenter, 1940:209); *H. hyalinatus* = *H. conjunctus* var. *pinidumus* (see Carpenter, 1940:211); *H. kootenayensis* = *H. pinidumus* (see Klimaszewski & Kevan, 1985:41); *H. pinidumus* variety of *H. conjunctus* (see Carpenter, 1940:209); *H. pinidumus* is valid species (see Throne, 1971b:81).

DISTRIBUTION.—CAN: AB<sup>6</sup>, BC<sup>3</sup>, PQ<sup>1</sup>; USA: CO<sup>4</sup>, IN<sup>5</sup>, MA<sup>4</sup>, ME<sup>4</sup>, MN<sup>4</sup>, NH<sup>4</sup>, NJ<sup>2</sup>, NY<sup>2</sup>, WS<sup>4</sup> (<sup>1</sup>Banks, 1897a:26, as *H. canadensis*;

<sup>2</sup>Smith, 1900:56, as *H. pinidumus* and *H. hyalinatus*; <sup>3</sup>Currie, 1904b:88, as *H. kootenayensis*;

<sup>4</sup>Carpenter, 1940:211; <sup>5</sup>Lawson & McCafferty, 1984:130; <sup>6</sup>Klimaszewski & Kevan, 1985:42).

**Hemerobius simulans** Walker, 1853:285.

Holotype male: Canada, Hudson's Bay, St. Martin's Falls, Albany River (BMNH).

= *Hemerobius orotopus* Wallengren, 1870:155. Syntypes (number unknown): Sweden, regions of Dalarne, Norbotten and Västbotten; one syntype in BMNH, others may be in ZIL and NRS.

= *Hemerobius nevadensis* Banks, 1904a:61.

Holotype female: USA, Nevada, Ormsby Co., 3.VII.1903, Baker (MCZ).

= *Hemerobius placidus* Banks, 1908a:260.

Lectotype male: USA, New York, Lake Placid, 12.VIII.1904, E. C. Van Duzee (MCZ).

= *Hemerobius piceus* Navás, 1925e:2. Three male, four female syntypes: Russia, Kamchatka Peninsula (NRS).

**TAXONOMY.** — *H. orotypus* = *H. simulans* (see Kimmins, 1932:88); *H. placidus* = *H. simulans*, *H. nevadensis* = *H. simulans* (see Carpenter, 1940:212); *H. piceus* = *H. simulans* (see Tjeder, 1960:148); lectotype designated for *H. placidus* (see Klimaszewski & Kevan, 1985:60).

**DISTRIBUTION.** — CAN: AB<sup>3</sup>, BC<sup>3</sup>, MB<sup>5</sup>, NB<sup>3</sup>, NF<sup>3</sup>, NS<sup>5</sup>, ON<sup>3</sup>, PQ<sup>3</sup>, SK<sup>5</sup>, YT<sup>5</sup>, USA: AK<sup>3</sup>, CO<sup>4</sup>, MA<sup>3</sup>, ME<sup>3</sup>, MI<sup>3</sup>, NH<sup>3</sup>, NV<sup>1</sup>, NY<sup>2</sup> (<sup>1</sup>Banks, 1904a:61; <sup>2</sup>Banks, 1908a:260; <sup>3</sup>Carpenter, 1940:212; <sup>4</sup>Nakahara, 1966:213; <sup>5</sup>Klimaszewski & Kevan, 1985:63).

**Hemerobius stigma** Stephens, 1836:112.

Lectotype (without abdomen): England, near Ripley (BMNH).

= *Hemerobius irroratus* Stephens, 1836:111. Lectotype (without abdomen): England, London metropolitan district (BMNH).

= *Hemerobius strigosus* Zetterstedt, 1840:1049. Type(s) (unknown): Lapponia-Upland-Ostrogoth-cania (ZIL).

= *Hemerobius limbatellus* Zetterstedt, 1840:1050. Type(s) (unknown): Lapponia meridional (ZIL).

= *Hemerobius limbatus* Wesmael, 1841:215. Type(s) (unknown): Belgium, Brussels (IRSNB).

= *Hemerobius crispus* Stephens, 1836:112. Single remaining syntype male: North America (BMNH).

= *Hemerobius stigmaterus* Fitch, 1855:797. Lectotype male: no locality data (MCZ).

= *Hemerobius moestus* Banks, 1897a:25. Holotype female: USA, Washington, Olympia, T. Kincaid (MCZ).

= *Hemerobius dyari* Currie, 1904b:85. Holotype female: Canada, British Columbia, Kaslo, 17.VII.1903 (USNM).

= *Hemerobius simplex* Banks, 1905c:32. Lectotype male: USA, Arizona, Prescott, 18.V.1902, Oslar (MCZ).

= *Hemerobius buyssoni* Navás, 1909b:217. Type(s) (unknown): France, La Bouboule (Puy-de-Dôme) (MNHN).

= *Hemerobius periphericus* Navás, 1913d:81. Type(s) (unknown): Spain, Escorial (depository of type unknown).

**TAXONOMY.** — *H. crispus* = *H. stigmaterus*, *H. moestus* = *H. stigmaterus*, *H. dyari* = *H. stigmaterus*, *H. simplex* = *H. stigmaterus*) (see Carpenter, 1940:202); *H. irroratus* = *H. stigma*, *H. strigosus* = *H. stigma*, *H. limbatellus* = *H.*

*stigma*, *H. limbatus* = *H. stigma*, *H. stigmaterus* = *H. stigma*, *H. buyssoni* = *H. stigma*, *H. periphericus* = *H. stigma* (see Klimaszewski & Kevan, 1985:44); lectotype designated for *H. stigmaterus* (see Carpenter, 1940:202); lectotype designated for *H. simplex* (see Klimaszewski & Kevan, 1985:46).

**DISTRIBUTION.** — CAN: AB<sup>11</sup>, BC<sup>3</sup>, LB<sup>6</sup>, MB<sup>11</sup>, NF<sup>11</sup>, NS<sup>11</sup>, ON<sup>11</sup>, PQ<sup>11</sup>, SK<sup>11</sup>, YT<sup>11</sup>, USA: AK<sup>12</sup>, AZ<sup>3</sup>, CA<sup>3</sup>, CO<sup>3</sup>, DC<sup>3</sup>, FL<sup>10</sup>, IA<sup>3</sup>, ID<sup>3</sup>, IN<sup>4</sup>, KS<sup>3</sup>, MD<sup>6</sup>, ME<sup>7</sup>, MI<sup>3</sup>, MN<sup>3</sup>, MO<sup>3</sup>, NC<sup>5</sup>, NJ<sup>3</sup>, NM<sup>3</sup>, NV<sup>3</sup>, NY<sup>2</sup>, TX<sup>9</sup>, WA<sup>1</sup>, WS<sup>8</sup> (<sup>1</sup>Banks, 1897a:25; <sup>2</sup>Smith, 1900:56, as *H. stigmaterus*; <sup>3</sup>Banks, 1905c:49; <sup>4</sup>Montgomery & Trippel, 1933:259; <sup>5</sup>Brimley, 1938:29, as *H. stigmaterus*; <sup>6</sup>Carpenter, 1940:203; <sup>7</sup>Procter, 1946:43; <sup>8</sup>Throne, 1971b:79; <sup>9</sup>Agnew et al., 1981:16; <sup>10</sup>MacLeod & Stange, 1981:1; <sup>11</sup>Klimaszewski & Kevan, 1985:49; <sup>12</sup>Klimaszewski & Kevan, 1989:205).

**REMARKS.** — *Hemerobius californicus* Banks, 1905a:90 (nomen nudum).

### Wesmaelius Krüger, 1922:170

**Wesmaelius brunneus** (Banks), 1920:333 [*Boriomyia*].

Holotype female: USA, Montana, Midvale, C. E. Brown (MCZ).

**TAXONOMY.** — To *Kimminisia* (see Carpenter, 1940:221); to *Wesmaelius* (see Kevan & Klimaszewski, 1986:19).

**DISTRIBUTION.** — CAN: AB<sup>2</sup>, BC<sup>4</sup>, YT<sup>4</sup>, USA: AK<sup>3</sup>, CA<sup>2</sup>, CO<sup>2</sup>, MT<sup>1</sup>, UT<sup>3</sup>, WA<sup>2</sup>, WY<sup>2</sup> (<sup>1</sup>Banks, 1920:333; <sup>2</sup>Carpenter, 1940:222; <sup>3</sup>Parfin, 1956:208; <sup>4</sup>Klimaszewski & Kevan, 1987b:256).

**Wesmaelius coloradensis** (Banks), 1897a:26 [*Hemerobius*].

Lectotype male: USA, Colorado, Fort Collins, F. C. Baker (MCZ).

**TAXONOMY.** — To *Boriomyia* (see Banks, 1905c:38); to *Kimminisia* (see Carpenter, 1940:217); to *Wesmaelius* (see Kevan & Klimaszewski, 1986:19); lectotype designated (see Carpenter, 1940:217).

**DISTRIBUTION.** — CAN: AB<sup>2</sup>, BC<sup>2</sup>, YT<sup>4</sup>, USA: CA<sup>2</sup>, CO<sup>1</sup>, NV<sup>3</sup>, OR<sup>2</sup>, UT<sup>2</sup>, WA<sup>3</sup>, WY<sup>2</sup> (<sup>1</sup>Banks, 1897a:26; <sup>2</sup>Carpenter, 1940:217; <sup>3</sup>Parfin, 1956:208; <sup>4</sup>Klimaszewski & Kevan, 1987b:252).

**Wesmaelius constrictus** (Parfin), 1956:203  
[*Kimminsia*].

Holotype male: USA, Alaska, 10–15 miles below Gulkana Lake along Gulkana River, 27.VI–20.VII.1955, G. O. Schumann (USNM).

= *Kimminsia olympica* Nakahara, 1966:216. Holotype male: USA, Washington, Deer Park, 5,400 ft., 17.VII.1948, C. P. Alexander (USNM).

TAXONOMY.—To *Wesmaelius*; *K. olympica* = *W. constrictus* (see Kevan & Klimaszewski, 1986:19).

DISTRIBUTION.—USA: AK<sup>1</sup>, WA<sup>2</sup> (<sup>1</sup>Parfin, 1956:203; <sup>2</sup>Nakahara, 1966:216, as *K. olympica*).

**Wesmaelius fumatus** (Carpenter), 1940:225  
[*Kimminsia*].

Holotype male: USA, Colorado, Boulder Co., Long's Peak Inn, 9000 ft., 13.VII.1926, E. C. Van Dyke (CAS).

TAXONOMY.—To *Wesmaelius* (see Kevan & Klimaszewski, 1986:19).

DISTRIBUTION.—CAN: AB<sup>2</sup>, USA: CO<sup>1</sup> (<sup>1</sup>Carpenter, 1940:225; <sup>2</sup>Klimaszewski & Kevan, 1987b:258).

**Wesmaelius furcatus** (Banks), 1935:55  
[*Boriomyia*].

Holotype male: USA, Colorado, Argentine Pass, T. D. A. Cockerell (MCZ).

TAXONOMY.—To *Kimminsia* (see Carpenter, 1940:219); to *Wesmaelius* (see Kevan & Klimaszewski, 1986:19).

DISTRIBUTION.—CAN: AB<sup>2</sup>, BC<sup>3</sup>, MB<sup>3</sup>, NT<sup>3</sup>, YT<sup>3</sup>, USA: AK<sup>3</sup>, CA<sup>3</sup>, CO<sup>1</sup> (<sup>1</sup>Banks, 1935:55; <sup>2</sup>Carpenter, 1940:219; <sup>3</sup>Klimaszewski & Kevan, 1987b:254).

**Wesmaelius involutus** (Carpenter), 1940:21  
[*Kimminsia*].

Holotype male: Canada, Alberta, Banff, Rundle Mountain, 5000–7000 ft., 25.VI.1925, Owen Bryant (MCZ).

TAXONOMY.—To *Wesmaelius* (see Kevan & Klimaszewski, 1986:19).

DISTRIBUTION.—CAN: AB<sup>1</sup>, BC<sup>1</sup>, NF<sup>3</sup>, NS<sup>3</sup>, NT<sup>3</sup>, PQ<sup>3</sup>, YU<sup>1</sup>, USA: AK<sup>2</sup>, CO<sup>1</sup>, ID<sup>1</sup>, UT<sup>2</sup>, WY<sup>1</sup> (<sup>1</sup>Carpenter, 1940:220; <sup>2</sup>Parfin, 1956:209; <sup>3</sup>Klimaszewski & Kevan, 1987a:175).

**Wesmaelius longifrons** (Walker), 1853:291  
[*Hemerobius*].

Holotype female: Canada, Ontario, Albany River, St. Martin's Falls (BMNH).

= *Hemerobius alternatus* Fitch, 1855:797. Holotype female: USA, New York (MCZ).

= *Hemerobius transversus* Banks, 1904a:61. Holotype male: USA, Colorado, Denver (MCZ).

= *Allotomyia borealis* Banks, 1935:56. Holotype male: USA, New Hampshire, Hillsboro, 11.VI (MCZ).

TAXONOMY.—To *Boriomyia* (see Banks, 1905c:37); to *Wesmaelius* (see Carpenter, 1940:226).

DISTRIBUTION.—CAN: AB<sup>3</sup>, BC<sup>3</sup>, MB<sup>3</sup>, NB<sup>5</sup>, NF<sup>5</sup>, NS<sup>3</sup>, ON<sup>1</sup>, PQ<sup>3</sup>, SK<sup>3</sup>, YT<sup>5</sup>, USA: AK<sup>3</sup>, AZ<sup>3</sup>, CO<sup>3</sup>, ME<sup>3</sup>, MI<sup>3</sup>, MN<sup>4</sup>, NH<sup>3</sup>, NY<sup>2</sup>, OR<sup>3</sup>, UT<sup>3</sup>, VT<sup>3</sup>, WA<sup>3</sup>, WY<sup>3</sup> (<sup>1</sup>Walker, 1853:291; <sup>2</sup>Smith, 1900:55; <sup>3</sup>Carpenter, 1940:226; <sup>4</sup>Parfin, 1952:423; <sup>5</sup>Klimaszewski & Kevan, 1987b:266).

**Wesmaelius longipennis** (Banks), 1920:333  
[*Boriomyia*].

Holotype female: USA, California, Alameda Co., Berkeley, 1.IV.1915 (MCZ).

TAXONOMY.—To *Kimminsia* (see Carpenter, 1940:221); to *Wesmaelius* (see Klimaszewski & Kevan, 1987a:163).

DISTRIBUTION.—USA: CA (Banks, 1920:333).

**Wesmaelius nervosus** (Fabricius), 1793:85  
[*Hemerobius*].

Two syntypes of unknown sex: France (Kiel).

= *Hemerobius betulinus* Ström, 1788:387 (nomen dubium). Type(s) lost.

= *Hemerobius nebulosus* Stephens, 1836:107. Type(s) (sex unknown): England, presumably near London (depository unknown).

= *Hemerobius conspersus* Burmeister, 1839:974. Type(s) (sex unknown): Germany, Halle (Halle).

= *Mucropalpus distinctus* Rambur, 1842:421. Type(s) (sex unknown): Spain, Andalusia (depository unknown).

= *Hemerobius disjunctus* Banks, 1897:25. Lectotype female: USA, New Hampshire, Mount Washington (MCZ).

= *Hemerobius frostinus* Navás, 1933b:109. Type(s) (sex unknown): USA, Massachusetts, Framingham (depository unknown).

= *Kimminsia alexanderi* Nakahara, 1966:217. Holotype male: USA, Alaska, Haines Highway, M. P., 5.VII.1952, C. P. Alexander (USNM).

= *Kimminsia melaleuca* Nakahara, 1966:219. Holotype female: Canada, Yukon Territory,

Alaska Highway, M. P. 320, 2.VII.1952, C. P. Alexander (USNM).

**TAXONOMY.** — To *Boriomyia* (see Banks, 1905c:29); to *Kimminsia* (see Hale Carpenter, 1938:532); to *Wesmaelius* (see Tjeder, 1967:5); *H. conspersus* = *H. nervosus* (see Wallengren, 1871:37); *H. nebulosus* = *B. betulina*, *M. distinctus* = *B. betulina* (see Killington, 1937:5); *H. disjunctus* = *W. nervosus*, *H. frostinus* = *W. nervosus*, *K. alexanderi* = *W. nervosus*, *K. melaleuca* = *W. nervosus* (Kevan & Klimaszewski, 1986:11); lectotype designated for *H. disjunctus* (see Klimaszewski & Kevan, 1987a:169).

**DISTRIBUTION.** — CAN: BC<sup>5</sup>, MB<sup>4</sup>, NB<sup>8</sup>, NF<sup>8</sup>, NS<sup>4</sup>, NT<sup>8</sup>, ON<sup>4</sup>, PQ<sup>4</sup>, SK<sup>4</sup>. USA: AK<sup>4</sup>, CO<sup>4</sup>, ID<sup>4</sup>, IN<sup>7</sup>, MA<sup>3</sup>, ME<sup>4</sup>, MI<sup>5</sup>, MT<sup>4</sup>, NC<sup>4</sup>, NH<sup>1</sup>, NY<sup>2</sup>, OR<sup>5</sup>, RI<sup>5</sup>, UT<sup>5</sup>, WS<sup>6</sup>, WY<sup>4</sup> (<sup>1</sup>Banks, 1897:25; <sup>2</sup>Leonard, 1928:40; <sup>3</sup>Navás, 1933b:109; <sup>4</sup>Carpenter, 1940:216; <sup>5</sup>Parfin, 1956:208, as *K. disjuncta*; <sup>6</sup>Throne, 1971b:84, as *K. disjuncta*; <sup>7</sup>Lawson & McCafferty, 1984:130, as *K. disjuncta*; <sup>8</sup>Klimaszewski & Kevan, 1987a:172).

**Wesmaelius posticatus** (Banks), 1905c:39 [*Boriomyia*].

Holotype female: USA, south Utah, VII.1900 (MCZ).

**TAXONOMY.** — To *Kimminsia* (see Carpenter, 1940:218); to *Wesmaelius* (see Kevan & Klimaszewski, 1986:19).

**DISTRIBUTION.** — CAN: AB<sup>2</sup>, YT<sup>3</sup>, USA: AZ<sup>2</sup>, CA<sup>3</sup>, CO<sup>2</sup>, NM<sup>2</sup>, UT<sup>1</sup>, (<sup>1</sup>Banks, 1905c:39; <sup>2</sup>Carpenter, 1940:219; <sup>3</sup>Nakahara, 1966:217).

**Wesmaelius pretiosus** (Banks), 1908a:260 [*Boriomyia*].

Lectotype male: USA, Colorado, Golden, 22.VII.1907, Oslar (MCZ).

**TAXONOMY.** — To *Kimminsia* (see Carpenter, 1940:223); to *Wesmaelius* (see Kevan & Klimaszewski, 1986:19; lectotype designated (see Carpenter, 1940:223)).

**DISTRIBUTION.** — CAN: BC<sup>5</sup>, PQ<sup>5</sup>, USA: AZ<sup>2</sup>, CO<sup>1</sup>, NE<sup>2</sup>, NV<sup>6</sup>, OR<sup>2</sup>, UT<sup>3</sup>, WS<sup>4</sup> (<sup>1</sup>Banks, 1908a:260; <sup>2</sup>Carpenter, 1940:223; <sup>3</sup>Parfin, 1956:209; <sup>4</sup>Throne, 1971b:84; <sup>5</sup>Klimaszewski & Kevan, 1987b:259; <sup>6</sup>Penny, NEW STATE RECORD).

**Wesmaelius schwarzi** (Banks), 1903b:24 [*Hemerobius*].

Holotype female: USA, Arizona, Williams, 23.VII, H. S. Barber (USNM).

= *Boriomyia ultima* Banks, 1930a:223 Holotype female: USA, California, Angora Park, Tahoe, E. P. Van Duzee (MCZ).

**TAXONOMY.** — To *Boriomyia* (see Banks, 1905c:38); to *Kimminsia* (see Carpenter, 1940:223); to *Wesmaelius* (see Kevan & Klimaszewski, 1986:19); *B. ultima* = *K. schwarzi* (see Carpenter, 1940:223).

**DISTRIBUTION.** — USA: AZ<sup>1</sup>, CA<sup>2</sup>, CO<sup>2</sup>, NM<sup>2</sup> (<sup>1</sup>Banks, 1903b:241; Carpenter, 1940:224).

**Wesmaelius subnebulosus** (Stephens), 1836:107 [*Hemerobius*].

Type(s) (sex unknown): England, presumably near London (depository unknown).

= *Hemerobius fuscus* Stephens, 1836:107. Type(s) (sex unknown): England, presumably near London (depository unknown).

= *Boriomyia maorica* Tillyard, 1923:221. Holotype female: New Zealand, Dunedin, George Street (CIC).

**TAXONOMY.** — To *Boriomyia* (see Banks, 1905c:29); to *Kimminsia* (see Killington, 1937:255); to *Wesmaelius* (see Tjeder, 1967:5); *H. fuscus* = *H. subnebulosus*, *H. nebulosus* = *H. subnebulosus* (see Wallengren, 1871:36); *B. maorica* = *W. subnebulosus* (see Wise, 1973:181, 182).

**DISTRIBUTION.** — CAN: NS<sup>3</sup>, ON<sup>3</sup>, PQ<sup>3</sup>, USA: CT<sup>1</sup>, IN<sup>2</sup>, NY<sup>1</sup> (<sup>1</sup>Parfin, 1956:206; <sup>2</sup>Lawson & McCafferty, 1984:130; <sup>3</sup>Klimaszewski & Kevan, 1987b:248).

**Wesmaelius yukonensis** Klimaszewski & Kevan, 1987b:262.

Holotype male: Canada, Yukon Territory, km 1,683 Alaska Highway, 7.VI.1979 (ROM).

**DISTRIBUTION.** — CAN: YT (Klimaszewski & Kevan, 1987b:263).

## MEGALOMINAE

### **Megalomus** Rambur, 1842:418

**Megalomus angulatus** Carpenter, 1940:242.

Holotype male: USA, New York, Ithaca, 3.IX.1939, J. S. Franclemont (Cornell).

**DISTRIBUTION.** — CAN: ON, USA: AZ, ME, NH, NY (Carpenter, 1940:242).

**Megalomus carpenteri** Penny, Adams & Stange, NEW NAME.

= *Hemerobius speciosus* (Banks), 1904a:62. A junior primary homonym of *Hemerobius specio-*

*sus* Linnaeus, 1758:551 (now *Palpares speciosus*). Holotype female: USA, Maryland, Plummer's Island, 9.IX (MCZ).

TAXONOMY.—To *Boriomyia* (see Banks, 1904c:209); to *Allotomyia* (see Banks, 1930a:224); returned to *Boriomyia* (see Carpenter, 1940:245); to *Megalomus* (see Oswald, 1993:241).

DISTRIBUTION.—USA: FL<sup>2</sup>, MD<sup>1</sup>, VA<sup>2</sup> (<sup>1</sup>Banks, 1904a:62; <sup>2</sup>Carpenter, 1940:245).

**Megalomus fidelis** (Banks), 1897:27 [*Hemerobius*].

Holotype female: USA, New York, Gowanda, 25.VI, E. P. Van Duzee (MCZ).

TAXONOMY.—To *Boriomyia* (see Banks, 1904c:209); to *Allotomyia* (see Banks, 1930a:224); returned to *Boriomyia* (see Carpenter, 1940:244); to *Megalomus* (see Oswald, 1993:239).

DISTRIBUTION.—CAN: BC<sup>4</sup>, ON<sup>4</sup>, USA: DC<sup>4</sup>, FL<sup>6</sup>, IL<sup>4</sup>, IN<sup>7</sup>, MA<sup>4</sup>, MD<sup>4</sup>, MN<sup>5</sup>, MS<sup>4</sup>, NC<sup>3</sup>, NH<sup>4</sup>, NJ<sup>2</sup>, NY<sup>1</sup>, PA<sup>4</sup>, SC<sup>4</sup>, TX<sup>4</sup>, VA<sup>4</sup>, WV<sup>4</sup> (<sup>1</sup>Banks, 1897:27; <sup>2</sup>Smith, 1900:56; <sup>3</sup>Brimley, 1938:29; <sup>4</sup>Carpenter, 1940:244; <sup>5</sup>Parfin, 1952:423, as *Boriomyia*; <sup>6</sup>MacLeod & Stange, 1981:1, as *Boriomyia*; <sup>7</sup>Lawson & McCafferty, 1984:130).

**Megalomus minor** Banks, 1905a:90.

Lectotype male: USA, Nevada, Ormsby Co., G. F. Baker (MCZ).

TAXONOMY.—Lectotype designated by Carpenter, 1940:242.

DISTRIBUTION.—USA: AL<sup>2</sup>, CA<sup>2</sup>, MS<sup>2</sup>, NV<sup>1</sup> (<sup>1</sup>Banks, 1905a:90; <sup>2</sup>Penny, NEW STATE RECORDS).

**Megalomus moestus** Banks, 1895b:314.

Holotype (missing abdomen): USA, New Mexico, Santa Fé, VII, T. D. A. Cockerell (MCZ).

= *Megalomus latus* Banks, 1903b:240.

Holotype female: USA, Arizona, Williams (USNM).

TAXONOMY.—*M. latus* = *M. moestus* (see Carpenter, 1940:241).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>2</sup>, NM<sup>1</sup>, TX<sup>2</sup>, UT<sup>2</sup>, WY<sup>2</sup> (<sup>1</sup>Banks, 1895b:314; <sup>2</sup>Carpenter, 1940:241).

**Megalomus parvulus** Kimmins, 1935:614.

Holotype male: West Indies, St. Vincent Island, windward side. H. H. Smith (BMNH).

DISTRIBUTION.—USA: FL (Stange, 1995:7).

**Megalomus uniformis** Banks, 1935:56.

Holotype female: USA, Texas, Brownsville, VI, F. H. Snow (MCZ).

DISTRIBUTION.—USA: TX (Banks, 1935:56).

## MICROMINAE

**Micromus** Rambur, 1842:416

**Micromus angulatus** (Stephens), 1836:106 [*Hemerobius*].

Holotype female: no locality data (BMNH).

= *Hemerobius villosus* Zetterstedt, 1840:1050 (nomen nudum). Type(s) (sex unknown): (locality unknown) (Lund).

= *Hemerobius intricatus* Wesmael, 1841:214.

Type(s) (sex unknown): (locality unknown) (IRSNB).

= *Micromus tendinosus* Rambur, 1842:417.

Type(s) (sex unknown): Sardinia (MNHN).

= *Hemerobius lineatus* Göszy, 1852:345.

Type(s) (sex unknown): no locality data (probably Vienna).

= *Micromus jonas* Needham, 1905:15. Type(s) (sex unknown): USA, New York (depository unknown).

TAXONOMY.—To *Micromus*; *H. villosus* = *M. angulatus*, *H. intricatus* = *M. angulatus*, *M. tendinosus* = *M. angulatus*, *H. lineatus* = *M. angulatus*, *M. aphidivorus* = *M. angulatus* (see Hagen, 1889:280); *M. jonas* = *M. angulatus* (see Carpenter, 1940:247).

DISTRIBUTION.—CAN: AB<sup>2</sup>, BC<sup>2</sup>, MB<sup>2</sup>, NF<sup>3</sup>, NS<sup>3</sup>, NT<sup>3</sup>, ON<sup>2</sup>, PQ<sup>2</sup>, SK<sup>2</sup>, YT<sup>2</sup>, USA: CO<sup>2</sup>, ME<sup>2</sup>, MI<sup>2</sup>, MN<sup>2</sup>, NH<sup>2</sup>, NY<sup>1</sup>, SD<sup>2</sup>, VT<sup>2</sup>, WS<sup>2</sup> (<sup>1</sup>Leonard, 1928:40; <sup>2</sup>Carpenter, 1940:247; Klimaszewski & Kevan, 1990b:65).

**Micromus borealis** Klimaszewski & Kevan, 1988:49.

DISTRIBUTION.—CAN: AB, BC, YT, USA: AK (Klimaszewski & Kevan, 1990b:65).

**Micromus montanus** Hagen, 1886:279.

Lectotype male: USA, Massachusetts, Natick (MCZ).

DISTRIBUTION.—CAN: AB<sup>4</sup>, BC<sup>4</sup>, NS<sup>4</sup>, PQ<sup>4</sup>, SK<sup>7</sup>, USA: AZ<sup>4</sup>, CA<sup>4</sup>, CO<sup>4</sup>, CT<sup>4</sup>, ID<sup>4</sup>, MA<sup>1</sup>, ME<sup>5</sup>, NC<sup>3</sup>, ND<sup>4</sup>, NH<sup>7</sup>, NV<sup>4</sup>, NY<sup>2</sup>, TN<sup>4</sup>, UT<sup>4</sup>, VT<sup>7</sup>, WA<sup>4</sup>, WS<sup>6</sup> (<sup>1</sup>Hagen, 1889:279; <sup>2</sup>Leonard, 1928:40; <sup>3</sup>Brimley, 1938:30; <sup>4</sup>Car-

penter, 1940:249; <sup>5</sup>Procter, 1946:43; <sup>6</sup>Throne, 1971b:83; <sup>7</sup>Klimaszewski & Kevan, 1990b:64). **Micromus posticus** (Walker), 1853:283 [*Hemerobius*].

Male and female syntypes: USA, Georgia (BMNH).

= *Micromus insipidus* Hagen, 1861:199. Syntypes (sex unknown): USA, New York; Pennsylvania, Philadelphia (MCZ).

= *Micromus sobrius* Hagen, 1861:199. Holotype (sex unknown): USA, Illinois, Chicago, Osten Sacken (MCZ).

**TAXONOMY.** — McLachlan, 1867:271 (*M. insipidus* = *M. posticus*, *M. sobrius* = *M. posticus* (see McLachlan, 1867:271)).

**DISTRIBUTION.** — CAN: BC<sup>11</sup>, MB<sup>11</sup>, ON<sup>11</sup>, PQ<sup>11</sup>, YT<sup>11</sup>, USA: AZ<sup>7</sup>, CO<sup>7</sup>, CT<sup>11</sup>, FL<sup>7</sup>, GA<sup>1</sup>, IA<sup>12</sup>, IL<sup>2</sup>, IN<sup>6</sup>, KS<sup>5</sup>, MA<sup>3</sup>, ME<sup>8</sup>, MN<sup>7</sup>, MO<sup>9</sup>, NC<sup>3</sup>, ND<sup>7</sup>, NE<sup>7</sup>, NJ<sup>4</sup>, NY<sup>2</sup>, OH<sup>12</sup>, PA<sup>2</sup>, TX<sup>7</sup>, VA<sup>11</sup>, WA<sup>11</sup>, WS<sup>10</sup> (<sup>1</sup>Walker, 1853:283; <sup>2</sup>Hagen, 1861:199, 200; <sup>3</sup>Hagen, 1889:286, as *M. insipidus*; <sup>4</sup>Smith, 1900:56; <sup>5</sup>Smith, 1925:167; <sup>6</sup>Montgomery & Trippel, 1933:259; <sup>7</sup>Carpenter, 1940:248; <sup>8</sup>Procter, 1946:43; <sup>9</sup>Froeschner, 1947:131; <sup>10</sup>Throne, 1971b:82; <sup>11</sup>Klimaszewski & Kevan, 1990b:65; <sup>12</sup>Penny, NEW STATE RECORDS).

**Micromus remiformis** Oswald, 1987:1.

Holotype male: USA, California, Madera Co., 4.75 miles east southeast of Fish Camp, 6400 ft., 9.VIII.1971, H. B. Leech (CAS).

**DISTRIBUTION.** — USA: CA, ID, NV, OR, UT (Oswald, 1987:3).

**Micromus subanticus** (Walker), 1853:282 [*Hemerobius*].

Holotype male: USA, East Florida, St. John's Bluff (BMNH).

= *Micromus angustus* Hagen, 1889:287. Seven syntypes: USA, Florida and North Carolina (MCZ and CNC).

= *Micromus nesoticus* Navás, 1914b:16. Holotype (sex unknown): USA, New York, Long Island, Yaphank (MZB).

= *Micromus haitiensis* Smith, 1931:800. Holotype female: Haiti, Petionville near Port-au-Prince, 17.II.1930, H. L. Dozier (MCZ).

**TAXONOMY.** — To *Micromus* (see Banks, 1905c:46); Carpenter, 1940:250, 251 (*M. angustus* = *M. subanticus*, *M. nesoticus* = *M. subanticus* (see Carpenter, 1940:250, 251); *M. haitiensis* = *M. subanticus* (see Monserrat, 1993:511)).

**DISTRIBUTION.** — CAN: BC<sup>11</sup>, MB<sup>11</sup>, ON<sup>11</sup>, PQ<sup>11</sup>, USA: AZ<sup>5</sup>, CA<sup>5</sup>, FL<sup>1</sup>, IA<sup>11</sup>, IL<sup>11</sup>, IN<sup>10</sup>, KS<sup>11</sup>, LA<sup>2</sup>, MA<sup>5</sup>, MN<sup>7</sup>, MO<sup>6</sup>, NC<sup>4</sup>, NV<sup>5</sup>, NY<sup>3</sup>, OH<sup>12</sup>, SC<sup>12</sup>, TX<sup>9</sup>, VA<sup>2</sup>, WS<sup>8</sup> (<sup>1</sup>Walker, 1853:282; <sup>2</sup>Banks, 1905c:46; <sup>3</sup>Leonard, 1928:40; <sup>4</sup>Brimley, 1938:30; <sup>5</sup>Carpenter, 1940:250, 251; <sup>6</sup>Froeschner, 1947:131; <sup>7</sup>Parfin, 1952:423; <sup>8</sup>Throne, 1971b:82; <sup>9</sup>Agnew et al., 1981:16; <sup>10</sup>Lawson & McCafferty, 1984:130; <sup>11</sup>Klimaszewski & Kevan, 1990b:65; <sup>12</sup>Monserrat, 1993:511).

**Micromus variegatus** (Fabricius, 1793:85) [*Hemerobius*].

Type(s): location unknown.

**DISTRIBUTION.** — CAN: BC (Klimaszewski & Kevan, 1990b:66).

**NOTE.** — This species appears to be established on Galiano Island, off the west coast of British Columbia, Canada. Klimaszewski and Kevan suspect this to be an introduction from Japan.

**Micromus variolosus** Hagen, 1886:284.

Holotype female: USA, Colorado, Denver (MCZ).

**DISTRIBUTION.** — CAN: BC<sup>3</sup>, SK<sup>6</sup>, YT<sup>6</sup>, USA: AZ<sup>3</sup>, CA<sup>3</sup>, CO<sup>1</sup>, ID<sup>3</sup>, IN<sup>5</sup>, KS<sup>2</sup>, MN<sup>4</sup>, NM<sup>3</sup>, NV<sup>3</sup>, OR<sup>3</sup>, UT<sup>3</sup>, WA<sup>3</sup>, WY<sup>3</sup> (<sup>1</sup>Hagen, 1886:284; <sup>2</sup>Smith, 1925:167; <sup>3</sup>Carpenter, 1940:251; <sup>4</sup>Parfin, 1952:423; <sup>5</sup>Lawson & McCafferty, 1984:130; <sup>6</sup>Klimaszewski & Kevan, 1990b:65).

#### NOTIOBIELLINAE

**Psectra** Hagen, 1866:376, 458

**Psectra diptera** (Burmeister), 1839:973 [*Hemerobius*].

Holotype male: Germany, Saxony, near Leipzig (Halle).

= *Hemerobius delicatulus* Fitch, 1855:800. Two syntypes: USA, Illinois (MCZ).

= *Psectra buenoi* Navás, 1909b:218. Five syntypes (1 male, 4 females): USA, New Jersey, Westfield, 5.VIII.1904; New York, Putnam Co., 5.IX.1903 (MZB).

**TAXONOMY.** — To *Psectra* (see Hagen, 1866:376); *H. delicatulus* = *P. diptera*, *P. buenoi* = *P. diptera* (see Carpenter, 1940:252).

**DISTRIBUTION.** — CAN: NF<sup>7</sup>, ON<sup>3</sup>, USA: IA<sup>8</sup>, IL<sup>1</sup>, IN<sup>6</sup>, MI<sup>3</sup>, MN<sup>4</sup>, NY<sup>2</sup>, VA<sup>3</sup>, WS<sup>5</sup>, WV<sup>3</sup>

(<sup>1</sup>Fitch, 1855:800; <sup>2</sup>Leonard, 1928:39; <sup>3</sup>Carpenter, 1940:252; <sup>4</sup>Parfin, 1952:423; <sup>5</sup>Throne, 1971b:84; <sup>6</sup>Lawson & McCafferty, 1984:130; <sup>7</sup>Kevan and Klimaszewski, 1986:9; <sup>8</sup>Penny, NEW STATE RECORD).

## SYMPHEROBIINAE

**Sypherobius** Banks, 1904c:209

**Sypherobius amiculus** (Fitch), 1855:799 [*Hemerobius*].

Lectotype female: USA, New York, Washington Co., Salem, 20.VIII.1852, A. Fitch (MCZ).

= *Sypherobius buenoi* Navás, 1912a:198  
Holotype (sex unknown): USA, New York, White Plains (MZB).

TAXONOMY.—To *Sypherobius* (see Banks, 1904c:209); *S. buenoi* = *S. amiculus* (see Carpenter, 1940:230). See note (Oswald, 1988:451) for clarification of type status.

DISTRIBUTION.—CAN: NS<sup>4</sup>, ON<sup>4</sup>, PQ<sup>8</sup>, USA: AL<sup>8</sup>, AR<sup>4</sup>, AZ<sup>8</sup>, CO<sup>8</sup>, CT<sup>8</sup>, DC<sup>8</sup>, FL<sup>4</sup>, GA<sup>4</sup>, IA<sup>8</sup>, IL<sup>4</sup>, IN<sup>7</sup>, KS<sup>2</sup>, LA<sup>4</sup>, MA<sup>4</sup>, MD<sup>8</sup>, ME<sup>5</sup>, MI<sup>4</sup>, MN<sup>4</sup>, MO<sup>6</sup>, MS<sup>8</sup>, NC<sup>3</sup>, NE<sup>4</sup>, NH<sup>4</sup>, NJ<sup>8</sup>, NY<sup>1</sup>, OH<sup>4</sup>, OK<sup>4</sup>, PA<sup>8</sup>, SC<sup>8</sup>, TN<sup>8</sup>, TX<sup>4</sup>, VA<sup>4</sup>, VT<sup>4</sup>, WS<sup>4</sup>, WV<sup>4</sup> (<sup>1</sup>Fitch, 1855:799; <sup>2</sup>Smith, 1925:167; <sup>3</sup>Brimley, 1938:29; <sup>4</sup>Carpenter, 1940:229; <sup>5</sup>Procter, 1946:42; <sup>6</sup>Froeschner, 1947:130; <sup>7</sup>Lawson & McCafferty, 1984:130; <sup>8</sup>Oswald, 1988:433).

**Sypherobius angustus** (Banks), 1904b:102 [*Hemerobius*].

Lectotype female: USA, New Mexico, Mesilla, A. P. Morse (MCZ).

= *Sypherobius tristis* Navás, 1914b:15.  
Type(s) (sex unknown): USA, New Mexico, Pecos (MZB).

= *Sypherobius strangei* Nakahara, 1960:16.  
Type (sex unknown): USA, California, San Bernardino Co., Barton Flats (probably in Nakahara Collection, Japan).

= *Sypherobius brunneus* Nakahara, 1966:207. Holotype male: USA, California, Mariposa Co., Miami Ranger Station (CAS).

TAXONOMY.—To *Sypherobius* (see Banks, 1904c:41); *S. tristis* = *S. angustus* (see Carpenter, 1940:233); *S. strangei* = *S. angustus*, *S. brunneus* = *S. angustus* (see Oswald, 1988:418).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: AZ<sup>3</sup>, CA<sup>3</sup>, CO<sup>2</sup>, ID<sup>3</sup>, NM<sup>1</sup>, NV<sup>3</sup>, OR<sup>3</sup>, SD<sup>3</sup>, UT<sup>2</sup>

(<sup>1</sup>Banks, 1904b:102; <sup>2</sup>Carpenter, 1940:233; <sup>3</sup>Oswald, 1988:421).

**Sypherobius arizonicus** Banks, 1911:346.

Holotype female: USA, Arizona, Prescott, Oslar (MCZ).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup> (<sup>1</sup>Banks, 1911:346; <sup>2</sup>Oswald, 1988:408).

**Sypherobius barberi** (Banks), 1903b:241 [*Hemerobius*].

Two female syntypes: USA, Arizona, Williams, 20,21.VII, H. S. Barber and E. A. Schwarz (USNM).

TAXONOMY.—To *Sypherobius* (see Banks, 1905c:42).

DISTRIBUTION.—USA: AL<sup>8</sup>, AR<sup>3</sup>, AZ<sup>1</sup>, CA<sup>3</sup>, CO<sup>3</sup>, FL<sup>3</sup>, IA<sup>8</sup>, IL<sup>8</sup>, IN<sup>7</sup>, KS<sup>2</sup>, LA<sup>8</sup>, MD<sup>8</sup>, MN<sup>5</sup>, MO<sup>4</sup>, MS<sup>8</sup>, NC<sup>3</sup>, NJ<sup>8</sup>, NM<sup>3</sup>, OK<sup>3</sup>, OR<sup>3</sup>, PA<sup>3</sup>, SC<sup>3</sup>, TX<sup>3</sup>, UT<sup>3</sup>, VA<sup>3</sup>, WS<sup>6</sup> (<sup>1</sup>Banks, 1903b:241; <sup>2</sup>Smith, 1925:166; <sup>3</sup>Carpenter, 1940:236; <sup>4</sup>Froeschner, 1947:130; <sup>5</sup>Parfin, 1952:423; <sup>6</sup>Throne, 1971b:83; <sup>7</sup>Lawson & McCafferty, 1984:130; <sup>8</sup>Oswald, 1988:429).

**Sypherobius beameri** Gurney, 1948:220.

Holotype male: USA, California, Lake Tahoe, 11.VIII.1940, R. H. Beamer (SEM).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, UT<sup>2</sup> (<sup>1</sup>Gurney, 1948:220; <sup>2</sup>Oswald, 1988:406).

**Sypherobius bifasciatus** Banks, 1911:347.

Holotype male: USA, Colorado, G. F. Baker (MCZ).

DISTRIBUTION.—CAN: AB<sup>2</sup>, USA: CO<sup>1</sup>, AZ<sup>3</sup>, CA<sup>3</sup>, UT<sup>3</sup> (<sup>1</sup>Banks, 1911:347; <sup>2</sup>Carpenter, 1940:233; <sup>3</sup>Oswald, 1988:423).

**Sypherobius californicus** Banks, 1911:346.

Lectotype male: USA, California, Pasadena (MCZ).

TAXONOMY.—Lectotype designated by Carpenter, 1940:232.

DISTRIBUTION.—USA: CA<sup>1</sup>, WA<sup>2</sup> (<sup>1</sup>Carpenter, 1940:232; <sup>2</sup>Oswald, 1988:425).

**Sypherobius constrictus** Oswald, 1988:425.

Holotype male: USA, California, San Francisco, Lobos Creek, 16.II.1967, J. Powell (CAS).

DISTRIBUTION.—USA: CA (Oswald, 1988:427).

**Sypherobius distinctus** Carpenter, 1940:238.

Holotype male: USA, Colorado, Pingree Park, 18.VIII.1926, R. C. Smith (MCZ).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CO<sup>1</sup> (<sup>1</sup>Carpenter, 1940:238; <sup>2</sup>Oswald, 1988:418).

**Sypherobius killingtoni** Carpenter, 1940:238.

Holotype male: USA, Utah, Eureka, 13.VIII, T. Spaulding (MCZ).

= *Sypherobius texanus* Nakahara, 1966:209.

Holotype male: USA, Texas, Kerr Co., Kerrville, VI.1954, L. J. Bottimer (USNM).

TAXONOMY.—*S. texanus* = *S. killingtoni* (see Oswald, 1988:408).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>1</sup>, CO<sup>2</sup>, ID<sup>1</sup>, NM<sup>1</sup>, NV<sup>2</sup>, OR<sup>2</sup>, TX<sup>1</sup>, UT<sup>1</sup>, WA<sup>2</sup> (<sup>1</sup>Carpenter, 1940:239; <sup>2</sup>Oswald, 1988:410).

**Sypherobius limbus** Carpenter, 1940:236.

Holotype male: USA, Texas, Alpine, 11.VII.1928, R. H. Beamer (SEM).

DISTRIBUTION.—USA: AZ<sup>2</sup>, NM<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Carpenter, 1940:236; <sup>2</sup>Oswald, 1988:416).

**Sypherobius occidentalis** (Fitch), 1855:799 [*Hemerobius*].

Holotype (sex unknown): USA, Illinois, Henderson River, 2.X.1854, A. Fitch (lost).

TAXONOMY.—To *Sypherobius* (see Banks, 1905c:40).

DISTRIBUTION.—USA: AL<sup>10</sup>, AR<sup>4</sup>, DC<sup>4</sup>, FL<sup>9</sup>, GA<sup>10</sup>, IL<sup>1</sup>, IN<sup>8</sup>, KS<sup>2</sup>, LA<sup>10</sup>, MO<sup>5</sup>, NC<sup>3</sup>, NY<sup>10</sup>, OH<sup>10</sup>, PA<sup>10</sup>, TX<sup>7</sup>, VA<sup>10</sup>, WS<sup>6</sup> (<sup>1</sup>Fitch, 1855:799; <sup>2</sup>Banks, 1905c:40; <sup>3</sup>Brimley, 1938:29; <sup>4</sup>Carpenter, 1940:231; <sup>5</sup>Froeschner, 1947:130; <sup>6</sup>Throne, 1971b:83; <sup>7</sup>Agnew et al., 1981:16; <sup>8</sup>Lawson & McCafferty, 1984:130; <sup>9</sup>MacLeod & Stange, 1981:2; <sup>10</sup>Oswald, 1988:415).

**Sypherobius perparvus** (McLachlan), 1869:22 [*Hemerobius*].

Lectotype male: USA, Texas, Bosque, G. W. Belfrage (BMNH).

= *Sypherobius sparsus* Banks, 1911:346.

Holotype female: USA, Kansas (MCZ).

TAXONOMY.—To *Sypherobius* (see Banks, 1905c:41); *S. sparsus* = *S. perparvus* (see Carpenter, 1940:238); lectotype designated by Oswald, 1988:403.

DISTRIBUTION.—CAN: SK<sup>3</sup>, USA: AZ<sup>3</sup>, CA<sup>3</sup>, CO<sup>3</sup>, NM<sup>3</sup>, OK<sup>3</sup>, IA<sup>4</sup>, KS<sup>2</sup>, MN<sup>4</sup>, MT<sup>4</sup>, NE<sup>4</sup>, NV<sup>4</sup>, OR<sup>4</sup>, SD<sup>4</sup>, TX<sup>1</sup>, UT<sup>4</sup>, WY<sup>4</sup> (<sup>1</sup>McLachlan, 1869:22; <sup>2</sup>Banks, 1911:346; <sup>3</sup>Carpenter, 1940:238; <sup>4</sup>Oswald, 1988:405).

**Sypherobius pictus** (Banks), 1904a:62 [*Hemerobius*].

Lectotype female: USA, southwestern Colorado, Oslar (MCZ).

TAXONOMY.—To *Sypherobius* (see Banks, 1905c:41); lectotype designated by Carpenter, 1940:235.

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>1</sup>, NM<sup>2</sup>, NV<sup>2</sup> (<sup>1</sup>Banks, 1904a:62; <sup>2</sup>Oswald, 1988:413).

**Sypherobius quadricuspis** Oswald, 1988:421.

Holotype male: USA, Arizona, Cochise Co., Southwest Research Station, 22–30.IV.1963, Vincent Roth (AMNH).

DISTRIBUTION.—USA: AZ (Oswald, 1988:422).

**Sypherobius similis** Carpenter, 1940:236.

Holotype male: USA, Arizona, Santa Rita Mountains, 17.VII.1932, R. H. Beamer (SEM).

DISTRIBUTION.—USA: AZ (Carpenter, 1940:236).

**Sypherobius umbratus** (Banks), 1903b:242 [*Hemerobius*].

Holotype male: USA, Arizona, Williams, 10.VI (USNM).

= *Sypherobius gracilis* Carpenter, 1940:231. Holotype male: USA, North Carolina, Raleigh, 11.VI.1934, R. R. Leiby (MCZ).

TAXONOMY.—To *Sypherobius* (see Banks, 1905c:41); *S. gracilis* = *S. umbratus* (see Oswald, 1988:430).

DISTRIBUTION.—USA: AR<sup>5</sup>, AZ<sup>1</sup>, FL<sup>4</sup>, GA<sup>5</sup>, MO<sup>3</sup>, NC<sup>2</sup>, NM<sup>2</sup>, OK<sup>5</sup>, PA<sup>5</sup>, SC<sup>5</sup>, VA<sup>5</sup> (<sup>1</sup>Banks, 1903b:242; <sup>2</sup>Carpenter, 1940:230, 231; <sup>3</sup>Froeschner, 1947:130; <sup>4</sup>MacLeod & Stange, 1981:2, as *S. gracilis*; <sup>5</sup>Oswald, 1988:431).

## ITHONIDAE

Ithonidae, or moth-like lacewings, are known from only one species in the genus *Oliarces* Banks 1908 in the United States, and are probably encountered less than any other family of Neuroptera. The original type locality is now flooded by the Salton Sea, and only one specimen (a male) was known until 1949. Its known distribution includes the desert of southern California and Arizona, being seen most often between the Gila Mtns. of Arizona and the Salton Sea. Adults are known to have mass emergences, and the larvae appear to feed on the roots of creosote bush (Faulkner, 1990a:18). Another species was recently described from Honduras (Penny, 1996). All other known species of this family are from Australia, although there are other undescribed species known from Mexico, and it is thought that the Mexican species *Narodona mexicana* Navás is actually an ithonid, rather than an osmylid, as originally indicated by Navás

(Adams, 1969). The adult behavior of *Oliarces clara* was described by Faulkner (1990a, 1990b).

### **Oliarces** Banks, 1908b:203

#### **Oliarces clara** Banks, 1908b:203.

Holotype male: USA, California, Imperial Co., Walter's Station, IV, J. B. Smith (MCZ).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, NV<sup>3</sup> (<sup>1</sup>Banks, 1908b:203; <sup>2</sup>Belkin, 1954:67; <sup>3</sup>Penny, NEW STATE RECORD).

### MANTISPIDAE

Mantispidae, or mantid-flies, superficially resemble preying mantids, or in the case of *Climaciella*, *Polistes* wasps. All species have raptorial forelegs as adults. Larvae are hypermetamorphic, being very active triungulins in the first instar, and becoming inactive scarabaeiform larvae in subsequent stages. Primitive members of the family are known to feed on noctuid pupae, solitary bee larvae, and polybiine social wasp larvae. Other species, in the more advanced Mantispinae, are spider egg sac predators, having a number of strategies for seeking out and entering the spider egg sacs (Redborg and MacLeod, 1985). In this family, adult size is dependent on larval food availability, and striking differences in size are common. The higher classification of Mantispidae has been discussed by Penny (1982b) and Lambkin (1986).

### SYMPHRASINAE

#### **Plega** Navás, 1928b:326

##### **Plega banksi** Rehn, 1939:248.

Holotype male: USA, Arizona, Pima Co., Kits Peak Rincon, Balboquivari Mts., 4050', 14.VIII.1916, Lutz & Rehn (AMNH).

DISTRIBUTION.—USA: AZ (Rehn, 1939:250).

##### **Plega dactylota** Rehn, 1939:250.

Holotype male: USA, Arizona, Mohave Co., Kingman, 20.VII.1920, O. C. Poling (ANSP).

= *Plega dactylota lipanica* Rehn, 1939:254. Holotype male: USA, Texas, Brewster Co., hills west of Ord Mountains, 1–15.VI.1926, O. C. Poling (ANSP).

TAXONOMY.—Two subspecies synonymized (see Rice, 1987:342).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NV<sup>1</sup>, TX<sup>1</sup>, UT<sup>1</sup> (<sup>1</sup>Rehn, 1939:253; <sup>2</sup>Penny, NEW STATE RECORD).

#### **Plega fratercula** Rehn, 1939:247.

Holotype male: USA, Arizona, Gila Co., Captain Mountain, 8.VIII.1933, R. Anderson (MCZ).

DISTRIBUTION.—USA: AZ (Rehn, 1939:248).

#### **Plega signata** (Hagen), 1877:208 [*Symphrasis*].

Holotype female: USA, California, Kern Co., Fort Tejon, 1879, F. Brown (MCZ).

TAXONOMY.—To *Plega* (see Navás, 1928b:326).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup> (<sup>1</sup>Hagen, 1877:208; <sup>2</sup>Rehn, 1939:246).

### CALOMANTISPINAE

#### **Nolima** Navás, 1914f:100

##### **Nolima dine** Rehn, 1939:261.

Holotype male: USA, Arizona, Pima Co., Santa Catalina Mts., Pepper Sauce Canyon, 16.VIII.1924, J. O. Martin (CAS).

DISTRIBUTION.—USA: AZ (Rehn, 1939:262).

##### **Nolima kantsi** Rehn, 1939:260.

Holotype male: USA, Texas, Brewster Co., Chisos Mts., 16.VII.1921, C. D. Duncan, (CAS).

DISTRIBUTION.—USA: TX (Rehn, 1939:261).

##### **Nolima pinal** Rehn, 1939:257.

Holotype male: USA, Arizona, Gila Co., base of Pinal Mts., IX, D. K. Duncan (MCZ).

DISTRIBUTION.—USA: AZ (Rehn, 1939:259).

### MANTISPINAE

#### **Climaciella** Enderlein, 1910:360

##### **Climaciella brunnea** (Say), 1824:309 [*Mantispa*].

Syntype male: USA, Minnesota, St. Peter's River (depository unknown).

= *Mantispa denarius* Taylor, 1862:494 No types designated.

= *Mantispa burquei* Provancher, 1875:247. Type(s) (sex unknown): Canada, Quebec (depository unknown).

= *Mantispa brunnea* var. *occidentis* Banks, 1911:348. Syntype males: USA, Colorado, Golden, Chimney Gulch, Oslar; Nevada, Ormsby Co., Baker; Washington, Wawawai, Mann (MCZ).

= *Climaciella rubescens* Stitz, 1913:37. Holotype (sex unknown): Mexico, Tampico (HMB).

= *Climaciella rubescens* var. *unicolor* Stitz, 1913:39. Holotype (sex unknown): Mexico, Sierra Mixteca (HMB).

= *Climaciella rubescens* var. *laciniata* Stitz, 1913:39. Holotype (sex unknown): Mexico, Mexico (HMB).

**TAXONOMY.** — To *Climaciella* (see Enderlein, 1910:360); *M. burquei* = *M. brunnea* (see Provancher, 1877:174); *C. rubescens* and varieties *unicolor* and *laciniata* = *C. brunnea* (see Handschin, 1960:546547); *M. denarius* = *C. brunnea* (see Kevan, 1989:8); *C. brunnea* var. *occidentis* = *C. brunnea* (see Welch & Kondratieff, 1991:70).

**DISTRIBUTION.** — CAN: BC<sup>10</sup>, ON<sup>12</sup>, PQ<sup>4</sup>, USA: AZ<sup>14</sup>, CA<sup>16</sup>, CO<sup>15</sup>, FL<sup>2</sup>, GA<sup>2</sup>, ID<sup>15</sup>, IL<sup>3</sup>, IN<sup>8</sup>, KS<sup>6</sup>, MN<sup>1</sup>, MO<sup>11</sup>, NC<sup>9</sup>, NJ<sup>5</sup>, NV<sup>15</sup>, NY<sup>7</sup>, PA<sup>3</sup>, TX<sup>14</sup>, UT<sup>15</sup>, WA<sup>15</sup>, WS<sup>12</sup> (<sup>1</sup>Say, 1824:309; <sup>2</sup>Walker, 1853:214; <sup>3</sup>Hagen, 1861:208; <sup>4</sup>Provancher, 1875:247; <sup>5</sup>Smith, 1900:54; <sup>6</sup>Smith, 1925:166; <sup>7</sup>Leonard, 1928:39; <sup>8</sup>Montgomery & Trippel, 1933:259; <sup>9</sup>Brimley, 1938:29; <sup>10</sup>Spencer, 1942:24; <sup>11</sup>Froeschner, 1947:124; <sup>12</sup>Robert, 1949:10; <sup>13</sup>Throne, 1972:119; <sup>14</sup>Redborg & MacLeod, 1983:63; <sup>15</sup>Welch & Kondratieff 1991:71; <sup>16</sup>Penny, NEW STATE RECORD).

### ***Mantispa* Illiger in Kugleann, 1798:499**

#### ***Mantispa floridana* Banks, 1897a:23.**

Holotype (sex unknown): USA, Florida, Lake Worth, A. T. Slosson (MCZ).

**DISTRIBUTION.** — USA: FL (Banks, 1897a:23).

***Mantispa interrupta*** Say, 1825: unpaginated text to plate 25.

Type(s) (sex unknown): USA, Pennsylvania, Philadelphia (specimen probably lost).

= *Mantispa aureus* Taylor, 1862:494. No types designated.

= *Mantispa cincticornis* Banks, 1911:347. Holotype female: USA, Texas, near Brownsville, Snow (MCZ).

**TAXONOMY.** — *M. aureus* = *M. interrupta* (see Kevan, 1989:8); *M. cincticornis* = *M. interrupta* (see Welch & Kondratieff, 1991:72).

**DISTRIBUTION.** — CAN: PQ<sup>6</sup>, USA: CO<sup>10</sup>, IN<sup>9</sup>, KS<sup>3</sup>, MN<sup>7</sup>, MO<sup>5</sup>, NJ<sup>2</sup>, NY<sup>4</sup>, PA<sup>1</sup>, SC<sup>11</sup>, TX<sup>1</sup>, VA<sup>1</sup>, WS<sup>8</sup> (<sup>1</sup>Banks, 1892:358; <sup>2</sup>Smith, 1900:54; <sup>3</sup>Smith, 1925:166; <sup>4</sup>Leonard, 1928:39; <sup>5</sup>Froeschner, 1947:124; <sup>6</sup>Robert, 1949:12; <sup>7</sup>Parfin, 1952:423; <sup>8</sup>Throne, 1972:120; <sup>9</sup>Lawson & McCafferty, 1984:130; <sup>10</sup>Welch & Kondratieff, 1991:72; <sup>11</sup>Hoffman & Brushwein, 1992:161).

#### ***Mantispa moesta* Hagen, 1861:210.**

Type(s) (sex unknown): USA, Tennessee (MCZ).

**DISTRIBUTION.** — USA: TN (Hagen, 1861:210).

#### ***Mantispa pulchella* (Banks), 1912:179 [*Mantisippa*].**

Holotype female: USA, Utah, Eureka (MCZ).

**TAXONOMY.** — Transferred to *Mantispa* (see Redborg, 1976:17).

**DISTRIBUTION.** — USA: CO<sup>4</sup>, GA<sup>3</sup>, IL<sup>2</sup>, NC<sup>3</sup>, SC<sup>3</sup>, UT<sup>1</sup> (<sup>1</sup>Banks, 1912:179; <sup>2</sup>Redborg, 1976:18; <sup>3</sup>Hoffman & Brushwein, 1989:8; <sup>4</sup>Welch & Kondratieff, 1991:74).

#### ***Mantispa sayi* Banks, 1897a:23.**

Lectotype male: USA, Texas, Brazos Co. (MCZ).

= *Mantispa fuscicornis* Banks, 1911:347. Lectotype male: USA, Florida, Kissimmee (MCZ).

= *Mantispa uhleri* Banks, 1943:79. Holotype female: USA, Pennsylvania (MCZ).

**TAXONOMY.** — *M. fuscicornis* = *M. sayi*, *M. uhleri* = *M. sayi*; lectotypes designated for *M. sayi* and *M. fuscicornis* (see Hoffman, 1989:638, 639).

**DISTRIBUTION.** — USA: AR<sup>9</sup>, AZ<sup>9</sup>, CO<sup>10</sup>, CT<sup>9</sup>, FL<sup>1</sup>, GA<sup>9</sup>, IL<sup>5</sup>, IN<sup>8</sup>, KS<sup>2</sup>, MD<sup>9</sup>, MI<sup>3</sup>, MN<sup>7</sup>, MO<sup>6</sup>, MS<sup>9</sup>, NC<sup>4</sup>, NE<sup>9</sup>, NM<sup>9</sup>, OH<sup>9</sup>, OK<sup>9</sup>, PA<sup>5</sup>, SC<sup>9</sup>, TX<sup>1</sup>, UT<sup>9</sup>, WS<sup>5</sup> (<sup>1</sup>Banks, 1897a:23; <sup>2</sup>Smith, 1925:166; <sup>3</sup>Kaston, 1938:148, as *M. fuscicornis*; <sup>4</sup>Brimley, 1938:29; <sup>5</sup>Banks, 1943:79; <sup>6</sup>Froeschner, 1947:124; <sup>7</sup>Parfin, 1952:423; <sup>8</sup>Lawson & McCafferty, 1984:130, as *M. uhleri*; <sup>9</sup>Hoffman, 1989:637; <sup>10</sup>Welch & Kondratieff, 1991:72).

#### ***Mantispa scabrosa* (Banks), 1912:179 [*Mantisippa*].**

Holotype female: USA, New Mexico, Mesilla (MCZ).

**TAXONOMY.** — Transferred to *Mantispa* (see Welch & Kondratieff, 1991:75).

DISTRIBUTION.—USA: AZ<sup>2</sup>, NM<sup>1</sup>, TX<sup>2</sup>  
(<sup>1</sup>Banks, 1912:179; <sup>2</sup>Welch & Kondratieff, 1991:75).

**Mantispa viridis** Walker, 1853:227.

Holotype (sex unknown): USA, East Florida (BMNH).

DISTRIBUTION.—USA: FL<sup>1</sup>, SC<sup>5</sup>, TX<sup>2</sup>, VA<sup>3</sup>, WS<sup>4</sup> (<sup>1</sup>Walker, 1853:227; <sup>2</sup>Banks, 1907c:23; <sup>3</sup>Parfin, 1958:203; <sup>4</sup>Throne, 1972:121; <sup>5</sup>Hoffman & Brushwein, 1992:161).

REMARKS.—*Mantispa gulosus* Taylor, 1862:294 (nomen dubium); *Mantispa brevicollis* Banks, 1905a:88 (nomen nudum).

### MYRMELEONTIDAE

Myrmeleontidae, or ant-lions and doodlebugs, are among the largest, and in southern parts of the country, probably the most familiar neuropterans to most people. The larvae of one genus, *Myrmeleon*, form pits in sheltered areas of dry soil and can only move backward. They wait at the bottom of the pits with only their heads exposed, waiting for insects to fall in. To make sure that the prey cannot escape up the loose sides of the funnel, the ant-lion larva throws loose sand or soil at it with its head. Then, the larva drags the insect under the soil surface where the victim's movements are inhibited. The prey is drained of body liquids, and the hollow body thrown out of the pit by the powerful head and mandibles. Other genera of ant-lions move forward and backward (except *Vella*) and live just under the soil surface, in rock crevices, in caves, or in tree-holes. Larvae pupate under the soil surface, and a damselfly-like adult later emerges. Adults generally rest quietly on twigs and grass stems during the day and become active at night. However, some species are active during the daytime and have rather bright pigment patterns on the wings. Ant-lions are most abundant in hot, dry regions, but can also be found in moist, forested regions. North American species of Myrmeleontidae were reviewed by Banks (1927) and Stange and Miller (1990). Current views on higher categories of Myrmeleontidae are to be found in Stange (1994). Keys to genera of Glenurini are to be found in Stange (1970b) and genera and species of Nemoleontini in Stange (1970a).

### ACANTHAACLISINI

**Paranthaclisis** Banks, 1907a:275

**Paranthaclisis congener** (Hagen), 1861:224 [*Acanthaclisis*].

Holotype female: USA, "Western Texas," Pecos River (now New Mexico) (MCZ).

TAXONOMY.—To *Paranthaclisis* (see Banks, 1907a:275).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>3</sup>, NM<sup>1</sup>, OR<sup>3</sup>, UT<sup>3</sup>, WA<sup>3</sup> (<sup>1</sup>Hagen, 1861:224; <sup>2</sup>Currie, 1903:274; <sup>3</sup>Banks, 1927:81).

**Paranthaclisis hageni** (Banks), 1899a:170 [*Acanthaclisis*].

Lectotype male: USA, Arizona, Phoenix, V (MCZ).

TAXONOMY.—To *Paranthaclisis* (see Banks, 1907a:275); lectotype designated by Stange, 1961a:674.

DISTRIBUTION.—USA: AZ<sup>1</sup>, NM<sup>2</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1899a:170; <sup>2</sup>Banks, 1927:81).

**Paranthaclisis nevadensis** Banks, 1939:4.

Holotype male: USA, Nevada, Esmeralda Co., 2 mi. S Silver Peak, 4350', 22.VIII.1924, Rehn & Hebard (ANSP).

DISTRIBUTION.—USA: CA<sup>2</sup>, NV<sup>1</sup> (<sup>1</sup>Banks, 1939:4; <sup>2</sup>Stange, NEW STATE RECORD).

**Vella** Navás, 1913e:46

**Vella americana** (Drury), 1773: unpaginated index [*Myrmeleon*].

Type(s) (sex unknown): USA, New York (depository unknown).

= *Acanthaclisis striata* Hagen, 1861:324 (nomen nudum). Type(s) (sex unknown): Colombia (depository unknown).

TAXONOMY.—To *Myrmecoleon* (see Burmeister, 1839:996); to *Acanthaclisis* (see Rambur, 1842:380); to *Myrmeleon* (see Walker, 1853:317); returned to *Acanthaclisis* (see Hagen, 1861:223); to *Heoclisis* (see Taschenberg, 1879:186); *A. striata* = *A. americana* (see Hagen 1887:136); to *Grypoplectron* (see Esben-Petersen, 1928:75); to *Vella* (see Stange, 1980:3).

DISTRIBUTION.—USA: FL<sup>3</sup>, GA<sup>3</sup>, NC<sup>3</sup>, NJ<sup>2</sup>, NY<sup>1</sup>, SC<sup>1</sup>, VA<sup>3</sup> (<sup>1</sup>Hagen, 1861:223; <sup>2</sup>Smith, 1900:56; <sup>3</sup>Banks, 1927:83).

**Vella fallax haitiensis** Smith, 1931:815.

Holotype female: Haiti, Jacmel, V.1927, G. N. Wolcott (MCZ).

= *Vella fallax cubana* Hagen, 1860:363 [*Acanthaclisis*] (nomen nudum). Type(s) (sex unknown): unknown locality, unknown depository.

TAXONOMY.—To *Vella* (see Navás, 1913e:46); *Vella fallax cubana* = *Vella fallax haitiensis* (Stange, NEW SYNONYMY).

DISTRIBUTION.—USA: FL (Stange, NEW NATIONAL RECORD).

**Vella fallax texana** (Hagen), 1887:147 [*Acanthaclisis*].

Lectotype male: USA, Texas, Dimmit Co., Carrizo Springs (MCZ).

= *Acanthaclisis hesperus* Banks, 1914b:618.

Syntype males: USA, Utah, Eureka and New Mexico, Jemez Mts., 28.VII and 4.VIII, Spalding and Woodgate (MCZ). Stange, NEW SYNONYMY.

= *Vella texana minor* Banks, 1943:75. Holotype (sex unknown): USA, Arizona, Phoenix (MCZ). Stange, NEW SYNONYMY.

TAXONOMY.—To *Vella* (see Navás, 1913e:46); lectotype designated by Stange, 1961:674.

DISTRIBUTION.—USA: AZ<sup>3</sup>, CA<sup>3</sup>, NM<sup>2</sup>, TX<sup>1</sup>, UT<sup>2</sup> (Hagen, 1887:147; Banks, 1914b:618; Stange, NEW STATE RECORDS).

## DENDROLEONTINI

**Dendroleon** Brauer, 1866:42**Dendroleon obsoletus** (Say), 1839:44 [*Formicaleo*].

Type(s) (sex unknown): USA (specimen lost).

= *Myrmeleon nigrocinctus* Rambur, 1842:398. Holotype (sex unknown): no locality data (depository unknown).

TAXONOMY.—To *Myrmecoleon* as *M. ocellatus* Borkhausen:1791:161 (see Burmeister, 1839:995); to *Glenurus* (see Hagen, 1866:405); *D. obsoletus* not = *D. ocellatus* Borkhausen (see Hagen, 1888:185) to *Dendroleon* (see Hagen, 1888:187); *M. nigrocinctus* = *M. obsoletus* (see Hagen, 1866:445).

DISTRIBUTION.—USA: CT<sup>3</sup>, FL<sup>3</sup>, IA<sup>7</sup>, IL<sup>3</sup>, IN<sup>4</sup>, KS<sup>2</sup>, MA<sup>3</sup>, MD<sup>3</sup>, MI<sup>3</sup>, MO<sup>5</sup>, NC<sup>3</sup>, NH<sup>3</sup>, NJ<sup>1</sup>, NY<sup>3</sup>, VA<sup>3</sup>, WS<sup>6</sup> (Smith, 1900:56; Smith,

1925:169; Banks, 1927:7; Montgomery & Trippel, 1933:260; Froeschner, 1947:128; Throne, 1972:124; Penny, NEW STATE RECORD).

**Dendroleon speciosus** Banks, 1905b:7.

Holotype female: USA, Colorado, Boulder (MCZ).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>1</sup>, NM<sup>2</sup> (Banks, 1905b:7; Banks, 1927:8).

## NEMOLEONTINI

**Eremoleon** Banks, 1901a:366**Eremoleon femoralis** (Banks), 1942:146 [*Psammoleon*].

Holotype (sex unknown): 20 mi N. W. La Paz, Lower California, 16.VII.1938, Michelbacher & Ross (CAS).

TAXONOMY.—To *Eremoleon* (see Stange, 1970b:21).

DISTRIBUTION.—USA: AZ (Stange, NEW NATIONAL RECORD)

**Eremoleon gracile** Adams, 1957b:90.

Holotype female: USA, California, Riverside, 31.VIII.1939, P. DeBach (CAS).

DISTRIBUTION.—USA: CA (Adams, 1957b:90).

**Eremoleon insipidus** Adams, 1957b:88.

Holotype male: 5 mi S. San Miguel, Lower California, 20.VII.1938, Michelbacher & Ross (MCZ).

DISTRIBUTION.—USA: CA (Adams, 1957b:89).

**Eremoleon macer** (Hagen), 1861:236 [*Myrmeleon*].

Holotype (sex unknown): Mexico (Vienna).

= *Segura vitreus* Navás, 1914g:18. Holotype male: Mexico, Morelos, Cuernavaca, 1871, Bilimek (MNHN).

= *Novulga mexicana* Navás, 1925c:189. Holotype male: Mexico, Veracruz, Veracruz, 1921 (MNHN).

= *Hesperoleon atomarius* Navás, 1933c:105. Holotype (sex unknown): Mexico, Morelos, Cuernavaca, 1929–30 (Hamburg, specimen destroyed).

TAXONOMY.—To *Eremoleon* (see Banks, 1901:366); *S. vitreus* = *E. macer* (see Banks, 1927:71); *N. mexicana* = *E. macer*, *H. atomarius* = *E. macer* (see Stange, 1970:21).

DISTRIBUTION.—USA: AZ (Banks, 1927:71).

**Eremoleon nigribasis** Banks, 1920:329.

Syntypes male and female: USA, Utah, St. George, VI 5,6 (MCZ).

= *Eremoleon affine* Banks, 1942:144. Holotype male: Mexico, Baja California Sur, Miraflores, VII.8.1938 (CAS).

TAXONOMY.—*E. affine* = *E. nigribasis* (see Adams, 1957b:91).

DISTRIBUTION.—USA: AZ<sup>2</sup>, NM<sup>2</sup>, UT<sup>1</sup> (Banks, 1920:329; <sup>2</sup>Adams, 1957b:91).

**Eremoleon pallens** Banks, 1941b:101.

Syntypes (sex unknown): USA, Arizona, Pichaco Peak, VII.23, Bequaert, Tinkham, Flock (MCZ).

DISTRIBUTION.—USA: AZ (Banks, 1941b:101).

**Glenurus** Hagen, 1866:372, 405

**Glenurus gratus** (Say), 1839:45 [*Formicaleo*].

Holotype (sex unknown): USA, Indiana, Evansville, Corson (depository unknown).

= *Myrmecoleon roseipennis* Burmeister, 1839:995. Holotype (sex unknown): Nord Amerika (Halle).

TAXONOMY.—To *Myrmecoleon* (see Walker, 1853:392); to *Glenurus* (see Hagen, 1866:405); to *Dendroleon* (see Banks, 1892:360); returned to *Glenurus* (see Banks, 1907c:30); *M. roseipennis* = *M. gratus* (see Hagen, 1861:225).

DISTRIBUTION.—USA: FL<sup>3</sup>, IN<sup>1</sup>, KY<sup>3</sup>, MO<sup>4</sup>, NJ<sup>2</sup>, OH<sup>3</sup>, TN<sup>3</sup> (<sup>1</sup>Say, 1839:45; <sup>2</sup>Smith, 1900:56; <sup>3</sup>Banks, 1927:68; <sup>4</sup>Froeschner, 1947:128).

REMARKS.—Carpenter (1992:538) has indicated that Burmeister's *Handbuch der Entomologie* was apparently published in two parts. The first 400 pages were published in 1839, while the remainder, including all Neuroptera, were published in 1838. If this is found to be correct, it could lead to serious nomenclatorial instability. The synonymy of *Glenurus gratus* and *Glenurus roseipennis* would be reversed. Almost certainly the International Commission on Zoological Nomenclature would be petitioned to declare, for nomenclatorial purposes, that the entire book was published in 1839.

**Glenurus luniger** Gerstaecker, 1894:125.

Holotype female: Panama, Chiriqui (Greifswald).

DISTRIBUTION.—USA: AZ (Stange, 1970b:22).

**Glenurus snowii** Banks, 1907b:100.

Holotype female: USA, Arizona, Baboquivari Mountains (MCZ).

DISTRIBUTION.—USA: AZ (Banks, 1907b:100).

**Psammoleon** Banks, 1899b:69

**Psammoleon albovaria** (Banks), 1942:146 [*Puren*].

Holotype (sex unknown): Mexico, Baja California, Venancio, 17.VII.1938, Michelbacher & Ross (CAS).

TAXONOMY.—To *Psammoleon* (see Stange, 1970b:23).

DISTRIBUTION.—USA: CA (Stange, NEW NATIONAL RECORD).

**Psammoleon arizonensis** Banks, 1935:53.

Three syntype females: USA, Arizona, Tucson (one syntype); Phoenix (two syntypes) (MCZ).

DISTRIBUTION.—USA: AZ (Banks, 1935:53).

**Psammoleon bistictus** (Hagen), 1861:235 [*Myrmeleon*].

Holotype (sex unknown): Cuba, Poey (depository unknown).

TAXONOMY.—To *Nelees* (see Navás, 1921:117); to *Psammoleon* (see Banks, 1927:61).

DISTRIBUTION.—USA: FL (Banks, 1927:62).

**Psammoleon connexus** (Banks), 1920:329 [*Puren*].

Holotype male: USA, California, San Jacinto Mountains, 25.VI (MCZ).

TAXONOMY.—To *Psammoleon* (see Banks, 1927:64).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1920:329; <sup>2</sup>Banks, 1927:65).

**Psammoleon decipiens** Banks, 1935:53.

Two syntypes: USA, Georgia, Scriven Co., Millin (one male); Louisiana, Shreveport (one of unknown sex) (MCZ).

DISTRIBUTION.—USA: GA, LA (Banks, 1935:53).

**Psammoleon guttipes** Banks, 1906a:99.

Holotype male: USA, North Carolina, Tryon (MCZ).

DISTRIBUTION.—USA: FL<sup>3</sup>, GA<sup>3</sup>, KS<sup>2</sup>, LA<sup>3</sup>, MO<sup>4</sup>, NC<sup>1</sup>, NJ<sup>3</sup>, TX<sup>3</sup> (<sup>1</sup>Banks, 1906a:99;

<sup>2</sup>Smith, 1925:170; <sup>3</sup>Banks, 1928:64; <sup>4</sup>Froeschner, 1947:128).

**Psammoleon inscriptus** (Hagen), 1861:230 [*Myrmeleon*].

Holotype male: USA, western Texas (now New Mexico), Pecos River, Pope (MCZ).

TAXONOMY.—To *Psammoleon* (see Banks, 1904b:106); to *Puren* (see Banks, 1927:66); returned to *Psammoleon* (see Stange, 1970b:24).

DISTRIBUTION.—USA: CA<sup>1</sup>, CO<sup>2</sup>, NM<sup>1</sup>, UT<sup>3</sup> (<sup>1</sup>Hagen, 1851:230; <sup>2</sup>Hagen, 1875:922; <sup>3</sup>Banks, 1927:66; <sup>4</sup>Stange, NEW STATE RECORD).

**Psammoleon minor** Banks, 1927:62.

Lectotype female: USA, Florida, Loggerhead Key, Dry Tortugas, VI.1917 (MCZ).

TAXONOMY.—Lectotype designated by Stange, 1961:674.

DISTRIBUTION.—USA: FL (Banks, 1927:63).

**Psammoleon normalis** Banks, 1942:145.

Holotype female: Mexico, Baja California, Venancio, 17.VII.1938 (CAS).

DISTRIBUTION.—USA: AZ, CA (Stange, NEW NATIONAL RECORDS).

**Psammoleon sinuatus** Currie, 1903:275.

Holotype male: USA, Arizona, Santa Rita Mountains, 31.V (USNM).

= *Psammoleon serpentinus* Navás, 1923:185. Holotype (sex unknown): USA, New Mexico, Jemez Springs, VII.1916, Woodgate (MZB).

TAXONOMY.—*P. serpentinus* = *P. sinuatus* (see Banks, 1927:5).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CO<sup>1</sup>, NM<sup>2</sup>, UT<sup>2</sup> (<sup>1</sup>Currie, 1903:275; <sup>2</sup>Banks, 1927:64).

#### BRACHYNEMURINI

**Abatoleon** Banks, 1924:436

**Abatoleon dorsalis** (Banks), 1903c:240 [*Brachynemurus*].

Holotype female: USA, Texas, Laredo (MCZ).

= *Brachynemurus curriei* McClendon, 1906a:93. Holotype male: USA, Texas, Brownsville (USNM).

= *Netroneurus pygmaeus* Esben-Petersen, 1933:111. Holotype male: USA, Texas, Dallas, Boll (Hamburg, specimen destroyed)

TAXONOMY.—To *Austroleon* (see Banks, 1928:54); to *Abatoleon* (see Stange, 1994:80); *B.*

*curriei* = *B. dorsalis* (see Banks, 1927:54); *N. pygmaeus* = *B. dorsalis* (see Stange, 1970a:66).

DISTRIBUTION.—USA: TX (Banks, 1903c:240).

**Abatoleon indiges** (Walker), 1860:189 [*Myrmeleon*].

Holotype male: Haiti (BMNH).

= *Nelees cubitalis* Navás, 1921:117. Holotype female: Cuba, Habana, Playa Chivo, 7.VII.1919, Cervera (MCZ).

TAXONOMY.—To *Clathroneuria* (see Navás, 1922b:168); to *Austroleon* (see Smith, 1931:816); to *Abatoleon* (see Stange, 1994:80); *N. cubitalis* = *A. indiges* (see Smith, 1931:816).

DISTRIBUTION.—USA: FL (Stange, 1970a:68).

**Atricholeon** Stange, 1994:75, 82

**Atricholeon tuberculatus** (Banks), 1899b:70 [*Brachynemurus*].

Holotype female: USA, New Mexico, Mesilla (MCZ).

TAXONOMY.—To *Atricholeon* (see Stange, 1994:82).

DISTRIBUTION.—USA: AZ<sup>2</sup>, NM<sup>1</sup> (<sup>1</sup>Banks, 1899b:70; <sup>2</sup>Banks, 1928:49).

**Brachynemurus** Hagen, 1888:34

**Brachynemurus abdominalis** (Say), 1823:163 [*Myrmeleon*].

Neotype male: USA, Arkansas, Hope, 8.VI, Knobel (MCZ).

= *Myrmeleon iniquus* Walker, 1853:330.

Holotype male: (no locality data) (BMNH).

= *Myrmeleon salvus* Hagen, 1861:227. Lectotype male: USA, D. C., Washington (MCZ).

= *Myrmeleon juvencus* Hagen, 1861:234. Holotype female: USA, western Texas (now New Mexico), Pecos River, VII, Pope (MCZ).

= *Brachynemurus tenuis* Banks, 1898:204.

Lectotype male: USA, New Mexico, Mesilla, 28.VI.1897, Pope (MCZ).

= *Hesperoleon placidus* Navás, 1915a:51.

Holotype male: USA, New Jersey, Pt. Pleasant (depository unknown).

TAXONOMY.—To *Macronemurus* (see Hagen, 1866:424); to *Brachynemurus* (see Hagen, 1888:57); to *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see

Stange, 1970a:76); *M. juvencus* = *M. abdominalis* (see Hagen, 1888:5759); *M. salvus* = *B. abdominalis* (see Banks, 1907c:31); *H. placidus* = *B. abdominalis* (see Banks, 1927:5); *M. iniquis* = *B. abdominalis*, *B. tenuis* = *B. abdominalis*; neotype for *M. abdominalis* and lectotype for *M. salvus* designated (see Banks, 1938d:413–414); lectotype for *B. tenuis* designated (see Stange, 1961a:677).

DISTRIBUTION.—CAN: BC<sup>6</sup>, MB<sup>4</sup>, ON<sup>4</sup>, USA: AR<sup>8</sup>, AZ<sup>4</sup>, CA<sup>4</sup>, CO<sup>4</sup>, CT<sup>4</sup>, DC<sup>4</sup>, GA<sup>4</sup>, IA<sup>8</sup>, ID<sup>8</sup>, IL<sup>8</sup>, IN<sup>5</sup>, KS<sup>3</sup>, LA<sup>8</sup>, MA<sup>4</sup>, MD<sup>4</sup>, MI<sup>8</sup>, MN<sup>4</sup>, MO<sup>7</sup>, NC<sup>4</sup>, ND<sup>4</sup>, NE<sup>4</sup>, NJ<sup>2</sup>, NM<sup>1</sup>, NV<sup>8</sup>, NY<sup>4</sup>, OK<sup>8</sup>, OR<sup>4</sup>, PA<sup>8</sup>, SD<sup>8</sup>, TN<sup>8</sup>, TX<sup>4</sup>, UT<sup>4</sup>, VA<sup>4</sup>, WA<sup>4</sup>, WS<sup>9</sup>, WY<sup>8</sup> (<sup>1</sup>Hagen, 1861:234; <sup>2</sup>Smith, 1900:56; <sup>3</sup>Smith, 1925:169; <sup>4</sup>Banks, 1928:39; <sup>5</sup>Montgomery & Trippel, 1933:260; <sup>6</sup>Spencer, 1942:28; <sup>7</sup>Froeschner, 1947:128; <sup>8</sup>Stange, 1970a:map 7; <sup>9</sup>Throne, 1972:123).

**Brachynemurus blandus** (Hagen), 1861:235 [*Myrmeleon*].

Holotype female: USA, western Texas (now New Mexico), Pecos River, Pope (MCZ).

TAXONOMY.—To *Brachynemurus* (see Hagen, 1888:73); to *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see Stange, 1970a:77).

DISTRIBUTION.—USA: AZ<sup>4</sup>, CA<sup>4</sup>, CO<sup>3</sup>, KS<sup>2</sup>, NM<sup>1</sup>, NV<sup>3</sup>, TX<sup>4</sup>, UT<sup>3</sup>, WA<sup>4</sup>, WY<sup>3</sup> (<sup>1</sup>Hagen, 1861:235; <sup>2</sup>Smith, 1925:170; <sup>3</sup>Banks, 1927:29; <sup>4</sup>Stange, 1970a:78).

**Brachynemurus californicus** Banks, 1895a:519.

Lectotype male: Mexico, Baja California Sur, San José del Cabo (MCZ).

= *Brachynemurus fraternus* Banks, 1895a:520. Lectotype male: Mexico, Baja California Sur, San José del Cabo (MCZ).

= *Comptesa ambitiosa* Navás, 1915b:464. Holotype male: Mexico, Baja California (MNHN).

= *Belluga implexa* Navás, 1920:195. Holotype female: Mexico, Baja California Sur, La Paz, 1914, Diguet (MNHN).

TAXONOMY.—*B. fraternus* = *B. californicus*, *C. ambitiosa* = *B. californicus*, *B. implexa* = *B. californicus* (see Banks, 1942:139); lectotypes designated for *B. californicus* and *B. fraternus* (see Stange, 1961:675, 676).

DISTRIBUTION.—USA: AZ, CA (Stange, 1970a:93).

**Brachynemurus carolinus** Banks, 1911:349.

Lectotype female: USA, North Carolina, Southern Pines (MCZ).

TAXONOMY.—To *Netroneurus* (see Banks, 1927:42); returned to *Brachynemurus* (see Stange, 1961a:675); lectotype designated (see Stange, 1961a:675).

DISTRIBUTION.—USA: FL, GA, NC (Banks, 1927:43).

**Brachynemurus elongatus** Banks, 1904b:105.

Lectotype male: USA, New Mexico, Mesilla, 30.VI.1897, Morse (MCZ).

TAXONOMY.—Lectotype designated (see Stange, 1961:676).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, CO<sup>1</sup>, NE<sup>2</sup>, NM<sup>1</sup>, NV<sup>2</sup>, UT<sup>1</sup> (<sup>1</sup>Banks, 1928:48; <sup>2</sup>Stange, 1970a:88).

**Brachynemurus ferox** (Walker), 1853:332 [*Myrmeleon*].

Holotype male: USA, California, Hartweg (BMNH).

= *Brachynemurus brunneus* Currie, 1898:273.

Holotype male: USA, Wyoming, Yellowstone National Park, Fountain, 10.VIII.1896, Currie (USNM).

= *Brachynemurus centralis* Banks, 1898:204.

Lectotype female: USA, Colorado, Estes Park (MCZ).

TAXONOMY.—To *Macronemurus* (see Hagen, 1866:424); to *Hesperoleon* (see Banks, 1913b:64); to *Brachynemurus* (see Stange, 1970a:81); *B. centralis* = *B. brunneus* (see Banks, 1907c:31); *B. brunneus* = *B. ferox* (see Stange, 1970a:81).

DISTRIBUTION.—CAN: BC<sup>4</sup>, USA: AZ<sup>3</sup>, CA<sup>1</sup>, CO<sup>3</sup>, NM<sup>3</sup>, NV<sup>3</sup>, OR<sup>5</sup>, UT<sup>3</sup>, WY<sup>2</sup> (<sup>1</sup>Walker, 1853:332; <sup>2</sup>Currie, 1898:204; <sup>3</sup>Banks, 1928:25, as *H. brunneus*; <sup>4</sup>Spencer, 1942:28; <sup>5</sup>Stange, 1970a:81).

**Brachynemurus fuscus** (Banks), 1905b:6 [*Calinemurus*].

Lectotype male: USA, Arizona, Nogales, 6.VII.1903, Oslar (MCZ).

TAXONOMY.—To *Brachynemurus* (see Stange, 1970a:93); lectotype designated (see Stange, 1961a:677).

DISTRIBUTION.—USA: AZ (Banks, 1905b:6).

**Brachynemurus henshawi** (Hagen), 1887:216 [*Maracanda*].

Holotype male: USA, Oregon, Umatilla, 24.VI.1882, Henshaw (MCZ).

TAXONOMY.—To *Cryptoleon* (see Banks, 1907c:32); to *Brachynemurus* (see Stange, 1970a:82).

DISTRIBUTION.—USA: CA<sup>2</sup>, NV<sup>2</sup>, OR<sup>1</sup> (Hagen, 1887:216; <sup>2</sup>Stange, 1970a:82).

**Brachynemurus hubbardi** Currie, 1898:241.

Lectotype male: USA, Arizona, Fort Grant, 22.VII.1897, Hubbard (MCZ).

= *Brachynemurus cockerelli* Banks, 1902:86. Holotype female: USA, New Mexico, Lone Mountain (MCZ).

= *Brachynemurus hubbardi nubeculipennis* Currie, 1903:277. Lectotype male: USA, Arizona, Phoenix (USNM).

= *Brachynemurus hubbardi curtus* Banks, 1927:31. Lectotype male: USA, Kansas, Seward Co., 18.VIII.1911, Williams (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913:65); returned to *Brachynemurus* (see Stange, 1970a:88); *B. cockerelli* = *B. hubbardi* (see Currie, 1903:277); *B. h. nubeculipennis* = *B. h. hubbardi* (see Banks, 1907c:31); *B. h. curtus* = *B. h. hubbardi* (see Stange, 1970a:88); lectotype designated for *B. h. curtus* (see Stange, 1961a:677); for *B. hubbardi* and *B. h. nubeculipennis* (see Stange, 1970a:88).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CO<sup>5</sup>, KS<sup>3</sup>, NM<sup>2</sup>, NV<sup>5</sup>, OK<sup>5</sup>, TX<sup>4</sup>, UT<sup>4</sup> (Currie, 1898:241; <sup>2</sup>Banks, 1902:86; <sup>3</sup>Smith, 1925:169; <sup>4</sup>Banks, 1927:30, 31; <sup>5</sup>Stange, 1970a:89).

**Brachynemurus irregularis** Currie, 1906:186.

Holotype female: USA, Texas, Columbus, VI.1879, Schwarz (USNM).

= *Mastega texanus* Navás, 1914b:13. Holotype female: USA, Texas (Vienna).

TAXONOMY.—*B. irregularis* to *Hesperoleon* and *M. texanus* to *Cryptoleon* (see Banks, 1927:5, 36); returned to *Brachynemurus*; *M. texanus* = *B. irregularis* (see Stange, 1970a:78).

DISTRIBUTION.—USA: IL<sup>1</sup>, KS<sup>2</sup>, OK<sup>3</sup>, TX<sup>1</sup> (Currie, 1905:186; <sup>2</sup>Smith, 1925:170; <sup>3</sup>Stange, 1970a:79).

**Brachynemurus longicaudus** (Burmeister), 1839:994 [*Myrmecoleon*].

Syntypes male and female: "Nord-Amerika," Winthem Coll. (MCZ).

= *Myrmecoleon irroratus* Burmeister, 1839:995. Two syntype females: USA, South Carolina (Halle).

TAXONOMY.—To *Myrmeleon* (see Hagen, 1860:365); to *Macronemurus* (see Hagen, 1866:424); to *Brachynemurus* (see Hagen,

1888:35); *M. longicaudus* = *M. abdominalis*, *M. irroratus* = *M. conspersus* (see Hagen, 1860:365); *M. irroratus* = *M. talpinus* (see Hagen, 1861:226); *M. irroratus* = *M. conspersus* (see Taschenberg, 1879:213); *B. longicaudus* not = *B. abdominalis*, *B. irroratus* not = *B. conspersus*, *B. irroratus* = *B. longicaudus* (see Hagen, 1888:3536).

DISTRIBUTION.—USA: FL<sup>3</sup>, GA<sup>2</sup>, IN<sup>5</sup>, MS<sup>4</sup>, NC<sup>3</sup>, NJ<sup>3</sup>, SC<sup>1</sup>, VA<sup>4</sup> (<sup>1</sup>Burmeister, 1839:995; <sup>2</sup>Hagen, 1861:227; <sup>3</sup>Banks, 1927:46; <sup>4</sup>Stange, 1970a:83; <sup>5</sup>Lawson & McCafferty, 1984:130).

**Brachynemurus mexicanus** Banks, 1895a:520.

Lectotype female: Mexico, Tepic (MCZ).

TAXONOMY.—*B. mexicanus* = *B. versutus* (see Banks, 1901:365); *B. mexicanus* not = *B. versutus* (see Banks, 1927:32); lectotype designated by Stange, 1961:676.

DISTRIBUTION.—USA: AZ<sup>1</sup>, NM<sup>2</sup> (<sup>1</sup>Banks, 1927:33; <sup>2</sup>Stange, 1970a:91).

**Brachynemurus nebulosus** (Olivier), 1811:127 [*Myrmeleon*].

Type(s) (sex unknown): USA, New York, M. Bofe (depository unknown).

= *Myrmeleon talpinus* Hagen, 1866:448 (nomen nudum). Type(s) (sex unknown): USA, South Carolina, Zimmerman (depository unknown).

= *Myrmecoleon contaminatus* Burmeister, 1839:995. Syntype females: USA, South Carolina, Zimmerman (Halle).

= *Myrmeleon conspersus* Rambur, 1842:387. Holotype female: "l'Amerique septentrionale" (IRSNB).

TAXONOMY.—To *Macronemurus* (see Hagen, 1866:424); to *Maracanda* (see Hagen, 1887:212); to *Cryptoleon* (see Banks, 1901:330); to *Brachynemurus* (see Stange, 1970a:84); *M. irroratus* (?) = *M. conspersus* (see Walker, 1853:329); *M. irroratus* = *M. abdominalis*, *M. conspersus* = *M. abdominalis*, *M. talpinus* = *M. abdominalis* (see Hagen, 1861:226); *M. talpinus* = *M. nebulosus*, *M. contaminatus* = *M. nebulosus*, *M. irroratus* = *M. nebulosus*, *M. conspersus* = *M. nebulosus* (see Hagen, 1887:212).

DISTRIBUTION.—CAN: ON<sup>4</sup>, USA: FL<sup>4</sup>, GA<sup>4</sup>, IL<sup>6</sup>, IN<sup>7</sup>, LA<sup>6</sup>, MI<sup>4</sup>, MN<sup>5</sup>, NC<sup>4</sup>, NE<sup>6</sup>, NJ<sup>3</sup>, NY<sup>1</sup>, OH<sup>4</sup>, OK<sup>6</sup>, PA<sup>6</sup>, SC<sup>2</sup>, TX<sup>6</sup>, WS<sup>6</sup> (<sup>1</sup>Olivier, 1811:127; <sup>2</sup>Burmeister, 1839:995; <sup>3</sup>Smith, 1900:56, as *M. conspersa*; <sup>4</sup>Banks, 1927:11; <sup>5</sup>Parfin, 1952:425; <sup>6</sup>Stange, 1970a:84; <sup>7</sup>Lawson & McCafferty, 1984:130).

**Brachynemurus pulchellus** Banks, 1911:348.

Holotype female: USA, California, Brown, 10.X, Grinnell (MCZ).

= *Clathroneuria exigua* Navás, 1920:197.

Holotype male: Mexico, Baja California Sur, La Paz, 1914, Diguet (MNHN).

= *Netroneurus pulchellus pallescens* Banks, 1942:140. Holotype male: Mexico, Baja California, San Domingo (CAS).

TAXONOMY.—To *Netroneurus* (see Banks, 1927:42); returned to *Brachynemurus* (see Stange, 1970a:102); *C. exigua* = *N. pulchellus* (see Banks, 1942:138); *N. p. pallescens* = *B. pulchellus* (see Stange, 1970a:102).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, NV<sup>2</sup> (Banks, 1911:348; Stange, 1970a:104).

**Brachynemurus ramburi** Banks, 1907c:31 (replacement name).

Holotype female: "l'Amerique septentrionale" (IRSNB).

= *Myrmeleon nebulosus* Rambur, 1842:387. A junior primary homonym of *M. nebulosus* Olivier, 1811:127.

TAXONOMY.—To *Macronemurus* (see Hagen, 1866:444); to *Brachynemurus* (see Hagen, 1888:36).

DISTRIBUTION.—USA: FL, GA, NC (Banks, 1927:47).

**Brachynemurus sackeni** Hagen, 1888:94.

Lectotype female: USA, Texas, Dallas, Boll (MCZ).

= *Bollenga dinamitensis* Navás, 1932:14.

Holotype female: Mexico, Dinamita (Turin).

= *Hesperoleon brevipilis* Banks, 1938d:414.

Holotype male: USA, Arizona, Colorado River (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see Stange, 1970a:144); *B. dinamensis* = *B. sackeni*, *H. brevipilis* = *B. sackeni* (see Stange, 1970a:144); lectotype designated by Stange, 1961a:676.

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>3</sup>, CO<sup>3</sup>, KS<sup>4</sup>, NE<sup>4</sup>, NM<sup>3</sup>, NV<sup>3</sup>, OR<sup>4</sup>, TX<sup>1</sup>, UT<sup>3</sup>, WY<sup>4</sup> (Hagen, 1888:94; Banks, 1938d:414; Banks, 1928:34; Stange, 1970a: map 28).

**Brachynemurus seminolae** Stange, 1970a:86.

Holotype male: USA, Florida, Bay Co., St. Andrews State Park, 30.VIII.1960, L. A. Stange (UCD).

DISTRIBUTION.—USA: FL, NC (Stange, 1970a:86).

**Brachynemurus signatus** (Hagen), 1887:215 [*Maracanda*].

Lectotype female: USA, Michigan, Lake Superior, Whitefish Point (MCZ).

TAXONOMY.—To *Cryptoleon* (see Banks, 1907c:32); to *Brachynemurus* (see Stange, 1970a:79); lectotype designated by Banks, 1928:12.

DISTRIBUTION.—USA: CO<sup>5</sup>, DE<sup>5</sup>, IL<sup>3</sup>, KS<sup>2</sup>, MI<sup>1</sup>, MN<sup>4</sup>, ND<sup>5</sup>, NE<sup>5</sup>, NJ<sup>5</sup>, NY<sup>3</sup>, OH<sup>3</sup>, PA<sup>5</sup>, RI<sup>3</sup>, WS<sup>6</sup> (Hagen, 1887:215; Smith, 1925:170; Banks, 1928:12; Parfin, 1952:425; Stange, 1970a:80; Throne, 1972:123).

**Brachynemurus versutus** (Walker), 1853:331 [*Myrmeleon*].

Holotype male: Mexico (BMNH).

= *Hesperoleon spegazzinius* Navás, 1934:72. Two syntype females: Mexico, Texcoco, 20.X.1930; Ixcapalapa, 15.IX.1931, Spegazzini (depository unknown).

TAXONOMY.—To *Macronemurus* (see Hagen, 1866:449); to *Hesperoleon* (see Banks, 1928:13); to *Brachynemurus* (see Stange, 1970a:91); *H. spegazzinius* = *B. versutus* (see Stange 1970a:91).

DISTRIBUTION.—USA: AZ, CO, NM (Stange, 1970a:91).

**Chaetoleon** Banks, 1920:328**Chaetoleon pumilis** (Burmeister), 1839:995 [*Myrmecoleon*].

Type(s) (sex unknown): USA, South Carolina (Halle).

= *Myrmeleon angusticollis* Rambur, 1842:399. Holotype female: "Bombay" (IRSNB).

TAXONOMY.—To *Macronemurus* (see Hagen, 1866:424); to *Creagris* (see McLachlan, 1873b:137); to *Brachynemurus* (see Banks, 1892:361); to *Dendroleon* (see Banks, 1907c:30); to *Chaetoleon* (see Banks, 1920:328); returned to *Brachynemurus* (see Stange, 1970a:100); returned to *Chaetoleon* (see Stange, 1994:84); *M. angusticollis* = *B. pumilis* (see Stange, 1970a:10).

DISTRIBUTION.—USA: FL<sup>3</sup>, GA<sup>3</sup>, NC<sup>3</sup>, NY<sup>2</sup>, SC<sup>1</sup> (Burmeister, 1839:995; Smith, 1900:56; Banks, 1927:57).

**Chaetoleon pusillus** (Currie), 1899:363 [*Brachynemurus*].

Lectotype female: USA, New Mexico, Mesilla Park, 13.VI, Cockerell (USNM).

**TAXONOMY.**—To *Scotoleon* (see Banks, 1913b:65); to *Chaetoleon* (see Banks, 1920:328); returned to *Brachynemurus* (see Stange, 1970a:100); returned to *Chaetoleon* (see Stange, 1994:84); lectotype designated by Stange, 1970a:101.

**DISTRIBUTION.**—**USA:** AZ<sup>2</sup>, CA<sup>3</sup>, CO<sup>3</sup>, NM<sup>1</sup>, NV<sup>3</sup>, TX<sup>3</sup>, UT<sup>2</sup> (<sup>1</sup>Currie, 1899:363; <sup>2</sup>Banks, 1927:58; <sup>3</sup>Stange, 1970a:101).

**Chaetoleon tripunctatus** (Banks), 1922:60 [*Hesperoleon*].

Lectotype male: USA, Arizona, Palmerlee, VI (MCZ).

**TAXONOMY.**—To *Austroleon* (see Banks, 1928:55); to *Brachynemurus* (see Stange, 1970a:98); to *Chaetoleon* (see Stange, 1994:84); lectotype designated by Stange, 1961:677.

**DISTRIBUTION.**—**USA:** AZ (Banks, 1922:60).

#### **Clathroneuria** Banks, 1913b:65

**Clathroneuria arapahoe** Banks, 1938d:418.

Holotype male: USA, Arizona, Tucson (MCZ).

**TAXONOMY.**—To *Brachynemurus* (see Stange, 1970a:105); returned to *Clathroneuria* (see Stange, 1994:85).

**DISTRIBUTION.**—**USA:** AZ (Banks, 1938d:418).

**Clathroneuria coquillettii** (Currie), 1898:93 [*Brachynemurus*].

Holotype male: USA, Arizona, San Simon, 5.VII.1897, Hubbard (USNM).

= *Brachynemurus maculosus* Banks, 1899a:170. Holotype male: USA, California, Tehama (MCZ).

**TAXONOMY.**—To *Hesperoleon* (see Banks, 1913b:65); to *Brachynemurus* (see Stange, 1970a:106); to *Clathroneuria* (see Stange, 1994:85); *B. coquillettii* = *B. blandus* (see Banks, 1899a:171); *B. coquillettii* not = *B. blandus* (see Currie, 1903:276); *B. maculosus* = *B. coquillettii* (see Stange, 1970a:106).

**DISTRIBUTION.**—**USA:** AZ<sup>1</sup>, CA<sup>1</sup>, KS<sup>2</sup>, NM<sup>2</sup>, NV<sup>3</sup>, TX<sup>2</sup>, UT<sup>2</sup>, WA<sup>2</sup>, WY<sup>3</sup> (<sup>1</sup>Currie, 1898:93; <sup>2</sup>Banks, 1927:32, 37; <sup>3</sup>Stange, 1970a:Map 17).

**Clathroneuria navajo** Banks, 1938d:418.

Holotype female: USA, Arizona, Pinery Canyon (MCZ).

**TAXONOMY.**—To *Brachynemurus* (see Stange, 1970a:107); returned to *Clathroneuria* (see Stange, 1994:85).

**DISTRIBUTION.**—**USA:** AZ<sup>1</sup>, CA<sup>2</sup> (<sup>1</sup>Banks, 1938d:418; <sup>2</sup>Stange, 1970a:107).

**Clathroneuria schwarzi** (Currie), 1903:280 [*Brachynemurus*].

Holotype male: USA, Arizona, Flagstaff (USNM).

**TAXONOMY.**—To *Clathroneuria* (see Banks, 1913b:63); returned to *Brachynemurus* (see Stange, 1970a:107); returned to *Clathroneuria* (see Stange, 1994:85).

**DISTRIBUTION.**—**USA:** AZ<sup>1</sup>, CA<sup>2</sup>, CO<sup>3</sup>, ID<sup>3</sup>, NM<sup>2</sup>, NV<sup>3</sup>, OK<sup>3</sup>, OR<sup>3</sup>, TX<sup>3</sup>, UT<sup>2</sup>, WY<sup>3</sup> (<sup>1</sup>Currie, 1903:280; <sup>2</sup>Banks, 1927:52; <sup>3</sup>Stange, 1970a:107).

**Clathroneuria westcotti** (Stange), 1970a:109 [*Brachynemurus*].

Holotype male: USA, Arizona, Cochise Co., west side of Willcox Dry Lake, 7.VIII.1959, R. L. Westcott (LACM).

**TAXONOMY.**—To *Clathroneuria* (see Stange, 1994:85).

**DISTRIBUTION.**—**USA:** AZ (Stange, 1970a:110).

#### **Mexoleon** Stange, 1994:75, 86

**Mexoleon papago** (Currie), 1899:361 [*Brachynemurus*].

Holotype male: USA, Arizona, Santa Rita Mountains, Madera Canyon, 7.VI.1898, Schwarz (USNM).

= *Clathroneuria arioles* Banks, 1941b:103.

Holotype male: USA, California, Riverside, 25.IX.1940, G. & H. Sperry (MCZ).

**TAXONOMY.**—To *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see Stange, 1970a:97); to *Mexoleon* (see Stange, 1994:86); *C. arioles* = *B. papago* (see Stange, 1970a:97).

**DISTRIBUTION.**—**USA:** AZ<sup>1</sup>, CA<sup>2</sup>, OR<sup>3</sup>, TX<sup>3</sup> (<sup>1</sup>Currie, 1899:361; <sup>2</sup>Banks, 1941b:103; <sup>3</sup>Stange, 1970a:97).

#### **Scotoleon** Banks, 1913b:65

**Scotoleon carizzoanus** (Hagen), 1888:93 [*Brachynemurus*].

Lectotype male: USA, Texas, Dimmit Co., Carrizo Springs (MCZ).

= *Hesperoleon douglasi* Banks, 1927:20. Lectotype male: USA, Arizona, Nogales, 6.VII.1903, Oslar (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913b:64); returned to *Brachynemurus* (see Stange, 1970a:117); to *Scotoleon* (see Stange, 1994:87); *H. carrizonus* = *H. peregrinus* (see Banks, 1913b:64); *H. carrizonus* not = *H. peregrinus* (see Banks, 1927:16); *H. douglasi* = *B. carrizonus* (see Stange, 1970a:117); lectotype designated by Stange, 1961a:675.

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>3</sup>, NM<sup>2</sup>, NV<sup>3</sup>, OK<sup>3</sup>, TX<sup>1</sup>, UT<sup>3</sup> (<sup>1</sup>Hagen, 1888:93; <sup>2</sup>Banks, 1927:20; <sup>3</sup>Stange, 1970a:120).

**Scotoleon deflexus** (Adams), 1957b:98 [*Hesperoleon*].

Holotype male: USA, California, San Bernardino Co., Baker, 23.VIII.1952 (CAS).

TAXONOMY.—To *Brachynemurus* (see Stange, 1970a:120); to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: CA<sup>1</sup>, NV<sup>2</sup> (<sup>1</sup>Adams, 1957b:100; <sup>2</sup>Stange, 1970a:121).

**Scotoleon dissimilis** (Banks), 1903d:175 [*Brachynemurus*].

Lectotype male: Mexico, Baja California Sur, San José del Cabo (MCZ).

= *Brachynemurus plectus* Navás, 1913e:49.

Lectotype male: Mexico, Basse-Californie, Diguet (MNHN).

= *Hesperoleon planus* Navás, 1920:200. Holotype male: Mexico, Baja California Sur, La Paz, 1914, Diguet (MNHN).

= *Hesperoleon curriei* Banks, 1938d:416.

Holotype female: USA, Arizona, Santa Catalina Mountains (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1942:136); to *Scotoleon* (see Stange, 1994:87); *B. plectus* and *H. planus* probably = *B. dissimilis* (see Banks, 1942:135); *B. curriei* = *B. dissimilis* (see Stange, 1970a:121); lectotypes designated for *B. dissimilis* (see Stange, 1961a:675) and *B. plectus* (see Stange, 1970a:121).

DISTRIBUTION.—USA: AZ<sup>1</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1938d:416, as *H. curriei*; <sup>2</sup>Stange, 1970a:122).

**Scotoleon eiseni** Banks, 1908c:33 [*Brachynemurus*].

Lectotype male: Mexico, Baja California Sur, San José del Cabo (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1942:137); to *Scotoleon* (see Stange, 1994:87); lectotype designated (see Stange, 1961a:676).

DISTRIBUTION.—USA: AZ, CA, NV (Stange, 1970a:123).

**Scotoleon expansus** (Navás), 1913e:48 [*Brachynemurus*].

Holotype female: Mexico, Baja California, Diguet (MNHN).

= *Hesperoleon apache* Banks, 1938d:417. Holotype female: USA, Arizona, Globe (MCZ).

TAXONOMY.—To *Hesperoleon* via synonymy with *H. sackeni* (see Banks, 1942:138); returned to *Brachynemurus* (see Stange, 1970a:125); to *Scotoleon* (see Stange, 1994:87); *B. expansus* = *H. sackeni* (see Banks, 1942:138); *B. expansus* not = *B. sackeni*, *H. apache* = *B. expansus* (see Stange, 1970a:125).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NM<sup>2</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1938d:417 as *H. apache*; <sup>2</sup>Stange, 1970a:125).

**Scotoleon fidelitas** (Adams), 1957b:94 [*Hesperoleon*].

Holotype male: USA, Arizona, Santa Rita Mountains, Box Canyon, 21.VIII.1949 (MCZ).

TAXONOMY.—To *Brachynemurus* (see Stange, 1970a:126); to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: AZ, CA (Adams, 1957b:97).

**Scotoleon infuscatus** (Adams), 1957b:104 [*Hesperoleon*].

Holotype male: USA, California, Contra Costa Co., Antioch, 24.V.1949, MacSwain (CAS).

TAXONOMY.—To *Brachynemurus* (see Stange, 1970a:127); to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: CA (Adams, 1957b:105).

**Scotoleon intermedius** (Currie), 1903:283 [*Brachynemurus*].

Holotype male: USA, Arizona, Phoenix (USNM).

TAXONOMY.—To *Hesperoleon* (see Banks, 1927:18); returned to *Brachynemurus* (see Stange, 1970a:127); to *Scotoleon* (see Stange, 1994:87); *B. intermedius* = *B. texanus* (see Banks, 1907c:31); *B. intermedius* not = *B. texanus* (see Banks, 1927:18).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NV<sup>3</sup>, UT<sup>3</sup> (<sup>1</sup>Currie, 1903:283; <sup>2</sup>Banks, 1927:19; <sup>3</sup>Stange, 1970a:128).

**Scotoleon longipalpis** (Hagen), 1888:95  
[*Brachynemurus*].

Lectotype male: Mexico, Baja California Sur, Cabo San Lucas, 1860, Xanthus (MCZ).

= *Scotoleon congener* Banks, 1942:138. Holotype male: Mexico, Baja California, Chapala Dry Lake, 21.VI, Michelbacher & Ross (CAS).

TAXONOMY.—To *Scotoleon* (see Banks, 1913b:64); returned to *Brachynemurus* (see Stange, 1970a:128); returned to *Scotoleon* (see Stange, 1994:87); *B. congener* = *B. longipalpis* (see Stange, 1970a:129); lectotype designated (see Stange, 1961a:676).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NM<sup>2</sup>, NV<sup>2</sup>, TX<sup>2</sup> (<sup>1</sup>Currie, 1903:276; <sup>2</sup>Stange, 1970a:Map 19).

**Scotoleon marshi** (Stange), 1970a:130  
[*Brachynemurus*].

Holotype male: USA, Texas, Brewster Co., 22 miles south of Marathon, 3.IX.1950, Stange & Marsh (UCD).

TAXONOMY.—To *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: TX (Stange, 1970a:130).

**Scotoleon minusculus** (Banks), 1898:203  
[*Brachynemurus*].

Lectotype male: USA, California, Lancaster, VII, Morge (MCZ).

= *Brachynemurus texanus* Banks, 1903d:175  
Lectotype male: USA, Texas, Laredo, VIII, McClendon (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1927:24); returned to *Brachynemurus* (see Stange, 1961a:676); to *Scotoleon* (see Stange, 1994:87); *B. texanus* = *B. minusculus* (see Stange, 1970a:131); lectotype designated for *B. minusculus* (see Stange, 1961a:676), and for *B. texanus* (see Stange, 1970a:131).

DISTRIBUTION.—USA: AZ<sup>3</sup>, CA<sup>1</sup>, KS<sup>5</sup>, NE<sup>5</sup>, NM<sup>4</sup>, NV<sup>5</sup>, OK<sup>5</sup>, TX<sup>2</sup>, UT<sup>5</sup> (<sup>1</sup>Banks, 1898:203; <sup>2</sup>Banks, 1903d:175; <sup>3</sup>Currie, 1903:278; <sup>4</sup>Banks, 1928:24; <sup>5</sup>Stange, 1970a:131).

**Scotoleon minutus** (Adams), 1957b:100 [*Hesperoleon*].

Holotype male: USA, California, Cathedral City, 24.VII.1950, Isaak (CAS).

TAXONOMY.—To *Brachynemurus* (see Stange, 1970a:132); to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>1</sup>, NV<sup>2</sup> (<sup>1</sup>Adams, 1957b:102; <sup>2</sup>Stange, 1970a:133).

**Scotoleon niger** (Currie), 1898:134 [*Brachynemurus*].

Holotype female: USA, Arizona, Fort Grant, 20.VII.1897, Hubbard (USNM).

= *Hesperoleon huachuca* Banks, 1938d:415.  
Holotype male: USA, Arizona, Reef (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1927:26); returned to *Brachynemurus* (see Stange, 1970a:133); to *Scotoleon* (see Stange, 1994:87); *B. huachuca* = *B. niger* (see Stange, 1970a:133).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NV<sup>2</sup> (<sup>1</sup>Currie, 1898:134; <sup>2</sup>Stange, 1970a:134).

**Scotoleon nigrescens** (Stange), 1970a:135  
[*Brachynemurus*].

Holotype male: USA, Arizona, Cochise Co., 13 miles east of Douglas, San Bernardino Ranch, 12.VI.1959, L. Stange (UCD).

TAXONOMY.—To *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: AZ, TX (Stange, 1970a:136).

**Scotoleon nigrilabris** (Hagen), 1888:72  
[*Brachynemurus*].

Lectotype female: USA, Colorado, Manitou (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see Stange, 1970a:136); to *Scotoleon* (see Stange, 1994:87); lectotype designated (see Banks, 1927:27).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>6</sup>, CO<sup>1</sup>, ID<sup>6</sup>, KS<sup>3</sup>, MN<sup>5</sup>, MT<sup>6</sup>, NE<sup>4</sup>, NM<sup>4</sup>, NV<sup>4</sup>, SD<sup>4</sup>, TX<sup>6</sup>, UT<sup>4</sup>, WY<sup>6</sup> (<sup>1</sup>Hagen, 1888:72; <sup>2</sup>Currie, 1903:275; <sup>3</sup>Smith, 1925:169; <sup>4</sup>Banks, 1927:28; <sup>5</sup>Parfin, 1952:425; <sup>6</sup>Stange, 1970a:137).

**Scotoleon nivatensis** (Navás), 1915a:52 [*Hesperoleon*].

Holotype male: USA, Nevada, Lyon Co., Yerington (despository unknown).

TAXONOMY.—To *Brachynemurus* (see Stange, 1970a:138); to *Scotoleon* (see Stange, 1994:87); *H. nivatensis* = *H. singularis* (see Banks, 1927:5); *B. nivatensis* not = *B. singularis* (see Stange, 1970a:138).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>2</sup>, ID<sup>2</sup>, NV<sup>1</sup>, OR<sup>2</sup> (<sup>1</sup>Navás, 1915a:52; <sup>2</sup>Stange, 1970a:138).

**Scotoleon pallidus** (Banks), 1899a:171  
[*Brachynemurus*].

Holotype female: USA, Arizona, Phoenix, IX, Kunze (MCZ).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913:65); returned to *Brachynemurus* (see Stange, 1970a:139); to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NM<sup>2</sup>, NV<sup>2</sup>, TX<sup>2</sup> (<sup>1</sup>Banks, 1899a:171; <sup>2</sup>Stange, 1970a:139).

**Scotoleon peregrinus** (Hagen), 1861:234 [*Myrmeleon*].

Lectotype female: USA, California, Havilah (MCZ).

= *Brachynemurus assimilis* Banks, 1903d:174

Lectotype male: USA, California, Tehama, 28.VIII.1897 (MCZ).

TAXONOMY.—To *Brachynemurus* (see Hagen, 1888:59); to *Hesperoleon* (see Banks, 1913b:64 as synonym of *H. ferox*); returned to *Brachynemurus* (see Stange, 1970a:141); to *Scotoleon* (see Stange, 1994:87); lectotype designated for *H. peregrinus* (see Banks, 1927:16); for *B. assimilis* (see Stange, 1961:675); *B. peregrinus* = *B. ferox* (see Banks, 1903d:175); *B. peregrinus* = *B. ferox* (see Currie, 1903:276); *H. carrizonus* = *H. peregrinus* (see Banks, 1938d:414); *H. carrizonus* not = *H. peregrinus* (see Stange, 1961a:676); *B. ferox* not = *B. peregrinus*; *B. assimilis* = *B. peregrinus* (see Stange, 1970a:141).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: AZ<sup>2</sup>, CA<sup>1</sup>, CO<sup>2</sup>, NE<sup>2</sup>, NM<sup>2</sup>, NV<sup>2</sup>, UT<sup>2</sup>, WY<sup>2</sup> (<sup>1</sup>Hagen, 1861:325; <sup>2</sup>Stange, 1970a:142).

**Scotoleon quadripunctatus** (Currie), 1898:136 [*Brachynemurus*].

Holotype female: USA, California, San Bernardino Co., Coquillet (USNM).

TAXONOMY.—To *Hesperoleon* (see Banks, 1927:21); returned to *Brachynemurus* (see Stange, 1970a:143); returned to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, NV<sup>3</sup> (<sup>1</sup>Currie, 1898:136; <sup>2</sup>Banks, 1927:22; <sup>3</sup>Stange, 1970a:144).

**Scotoleon singularis** (Currie), 1903:284 [*Brachynemurus*].

Holotype male: USA, Arizona, Phoenix (USNM).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see Stange, 1970a:145); to *Scotoleon* (see Stange, 1994:87); *H. nivatensis* = *H. singularis* (see Banks, 1927:5); *B. nivatensis* not = *B. singularis* (see Stange, 1970a:145).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>3</sup>, NV<sup>3</sup>, NM<sup>3</sup>, UT<sup>2</sup> (<sup>1</sup>Currie, 1903:284; <sup>2</sup>Banks, 1927:18; <sup>3</sup>Stange, 1970a: 146).

**Scotoleon yavapai** (Currie), 1903:281 [*Brachynemurus*].

Holotype male: USA, Arizona, Yavapai Co., Hot Springs (USNM).

TAXONOMY.—To *Hesperoleon* (see Banks, 1913b:65); returned to *Brachynemurus* (see Stange, 1970a:148); returned to *Scotoleon* (see Stange, 1994:87).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>3</sup>, NV<sup>3</sup>, UT<sup>2</sup> (<sup>1</sup>Currie, 1903:281; <sup>2</sup>Banks, 1927:35; <sup>3</sup>Stange, 1970a:148).

#### GNOPHOLEONTINI

**Gnopholeon** Stange, 1970a:148

**Gnopholeon barberi** (Currie), 1903:282 [*Brachynemurus*].

Holotype female: USA, Arizona, Hot Springs (USNM).

TAXONOMY.—To *Scotoleon* (see Banks, 1913b:65); to *Austroleon* (see Banks, 1927:55); to *Gnopholeon* (see Stange, 1970a:150).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup> (<sup>1</sup>Currie, 1903:282; <sup>2</sup>Stange, 1970a:150).

**Gnopholeon delicatulus** (Currie), 1903:279 [*Brachynemurus*].

Holotype male: USA, Arizona, Phoenix, 11.V.1898, Kunze (USNM).

TAXONOMY.—To *Scotoleon* (see Banks, 1913b:65); to *Clathroneuria* (see Banks, 1927:52); to *Gnopholeon* (see Stange, 1970a:151).

DISTRIBUTION.—USA: AZ<sup>1</sup>, CA<sup>2</sup>, NV<sup>2</sup> (<sup>1</sup>Currie, 1903:279; <sup>2</sup>Stange, 1970a:152).

**Menkeleon** Stange, 1970a:154

**Menkeleon bellulus** (Banks), 1905b:7 [*Maracandula*].

Lectotype female: USA, California, Three Rivers (MCZ).

= *Maracandula bellula* Banks, 1905:90 (no-men nudum).

= *Maracandula minima* Banks, 1942:140.

Holotype male: USA, Mexico, Baja California, San Miguel (CAS).

TAXONOMY.—To *Menkeleon* (see Stange, 1970a:155); *M. minima* = *M. bellula* (see Stange, 1970a:155); lectotype designated (see Stange, 1970a:155).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup>, NM<sup>3</sup>, UT<sup>2</sup> (<sup>1</sup>Banks, 1905b:7; <sup>2</sup>Banks, 1927:59; <sup>3</sup>Stange, 1970a:157).

### *Tyttholeon* Adams, 1957b:106

#### *Tyttholeon puerilis* Adams, 1957b:107.

Holotype male: USA, California, Riverside Co., Palm Springs, 27.VI.1939, Isaacs (CAS).

DISTRIBUTION.—USA: AZ, CA (Adams, 1957b:108).

### MYRMELEONTINI

#### *Myrmeleon* Linnaeus, 1767:913

##### *Myrmeleon arizonicus* Banks, 1943:76.

Syntypes (sex unknown): USA, Arizona, Tucson, 6,20.VI; Texas, Brownsville, IV (MCZ).

DISTRIBUTION.—USA: AZ, TX (Banks, 1943:76).

##### *Myrmeleon californicus* Banks, 1943:77.

Holotype female: USA, California, Gavilan, 11.VII, G. & J. Sperry (MCZ).

DISTRIBUTION.—USA: CA (Banks, 1943:77).

##### *Myrmeleon carolinus* Banks, 1943:75.

Syntypes (sex unknown): USA, North Carolina, Southern Pines, VVIII, Manee; Georgia, Scriben Co., Millin, VII, Morrison (MCZ).

DISTRIBUTION.—USA: GA, NC (Banks, 1943:75).

##### *Myrmeleon crudelis* Walker, 1853:388.

Holotype (sex unknown): USA, E. Florida, St. John's Bluff, E. Doubleday (BMNH).

= *Myrmeleon tectus* Walker, 1853:378. Lectotype male: USA, E. Florida, St. John's Bluff, E. Doubleday (BMNH).

TAXONOMY.—*M. tectus* = *M. crudelis* Stange, NEW SYNONYMY.

DISTRIBUTION.—USA: FL<sup>1</sup>, GA<sup>2</sup>, MO<sup>3</sup>, NC<sup>2</sup>, NJ<sup>2</sup>, NY<sup>2</sup>, TX<sup>2</sup>, VA<sup>2</sup> (<sup>1</sup>Walker, 1853:378; <sup>2</sup>Banks, 1927:78; <sup>3</sup>Froeschner, 1947:128).

##### *Myrmeleon exitialis* Walker, 1853:376.

Holotype (sex unknown): USA, California, Hartweg's coll. (BMNH).

= *Myrmeleon immaculatus occidentalis* Currie, 1903:274. Holotype male: USA, Arizona, Williams (USNM).

= *Myrmeleon immaculatus coloradensis* Banks, 1905a:90 (nomen nudum). Holotype (sex unknown): USA, Nevada, Ormsby Co. (MCZ).

TAXONOMY.—*M. immaculatus occidentalis* = *M. exitialis*, *M. immaculatus coloradensis* = *M. exitialis*) Stange, NEW SYNONYMY.

DISTRIBUTION.—CAN: BC<sup>4</sup>, USA: AZ<sup>2</sup>, CA<sup>1</sup>, NV<sup>3</sup>, (<sup>1</sup>Walker, 1853:376; <sup>2</sup>Currie, 1903:274; <sup>3</sup>Banks, 1905a:90; <sup>4</sup>Spencer, 1942:28, as *M. immaculatus occidentalis*).

##### *Myrmeleon immaculatus* DeGeer, 1773:564.

Type(s) (sex unknown): USA, Pennsylvania, Acrelius (NRS).

= *Myrmeleon melanocephalum* Olivier, 1811:127. Type(s) (sex unknown): USA, New York, environs of New York, M. Bofe (depository unknown).

TAXONOMY.—To *Neleon* (see Navás, 1915a:53); returned to *Myrmeleon* (see Banks, 1928:74); *M. melanocephalum* = *M. immaculatus* (see Hagen, 1888:188).

DISTRIBUTION.—USA: DC<sup>3</sup>, FL<sup>5</sup>, GA<sup>3</sup>, IN<sup>6</sup>, KS<sup>4</sup>, MD<sup>5</sup>, MI<sup>5</sup>, MN<sup>8</sup>, MO<sup>7</sup>, NC<sup>5</sup>, NH<sup>5</sup>, NY<sup>2</sup>, OH<sup>5</sup>, OK<sup>5</sup>, PA<sup>1</sup>, VA<sup>3</sup>, WS<sup>9</sup> (<sup>1</sup>DeGeer, 1773:564; <sup>2</sup>Olivier, 1811; <sup>3</sup>Hagen, 1861:232; <sup>4</sup>Smith, 1925:170; <sup>5</sup>Banks, 1927:74 [western records excluded as probably pertaining to *M. exitialis*]; <sup>6</sup>Montgomery & Trippel, 1933:260; <sup>7</sup>Froeschner, 1947:128; <sup>8</sup>Parfin, 1952:425; <sup>9</sup>Throne, 1972:123).

##### *Myrmeleon insertus* Hagen, 1861:233.

Syntypes (sex unknown): Cuba, Poey; Haiti, Port-au-Prince (MCZ).

DISTRIBUTION.—USA: FL (Lucas & Stange, 1981:213).

##### *Myrmeleon invisus* Walker, 1853:379.

Holotype (sex unknown): no locality data (BMNH).

= *Myrmeleon heriocles* Banks, 1914b:619. Syntypes (sex unknown): USA, North Carolina, Southern Pines, May (MCZ).

TAXONOMY.—*M. heriocles* = *M. invisus* Stange, NEW SYNONYMY.

DISTRIBUTION.—USA: FL<sup>2</sup>, NC<sup>1</sup>, NJ<sup>2</sup> (<sup>1</sup>Banks, 1914b:618, as *M. heriocles*; <sup>2</sup>Banks, 1927:76, as *M. heriocles*).

##### *Myrmeleon mexicanus* Banks, 1903c:241.

Syntypes (sex unknown): Mexico, Guadaluja, McClendon (MCZ).

DISTRIBUTION.—USA: AZ (Banks, 1938d:420).

**Myrmeleon mobilis** Hagen, 1888:204.

Syntypes male and female: USA, Alabama, VII.1883, Lyon (MCZ).

= *Myrmeleon mobilis* Hagen, 1860:368 (nomen nudum).

DISTRIBUTION.—USA: AL<sup>1</sup>, FL<sup>5</sup>, GA<sup>2</sup>, MO<sup>4</sup>, NC<sup>3</sup>, (Hagen, 1888:204; Banks, 1927:75; Brimley, 1938:30; Froeschner, 1947:128; Lucas & Stange, 1981:214).

**Myrmeleon rusticus** Hagen, 1861:233.

Lectotype (sex unknown): USA, western Texas (now New Mexico), Pecos River, 4.VIII, Pope (MCZ).

= *Myrmeleon diversus* Hagen, 1873:729. Holotype (sex unknown): USA, Wyoming, Yellowstone National Park (MCZ).

= *Myrmeleon distans* Banks, 1898:206. Holotype (sex unknown): USA, California, Coronado Beach, VII, Morse (MCZ).

= *Myrmeleon agriope* Banks, 1914b:618. Syntypes (sex unknown): USA, California, Claremont; Arizona, Nogales; Arizona, Phoenix (MCZ).

TAXONOMY.—*M. rusticus* = *M. crudelis* (see Banks, 1907c:29); *M. rusticus* not = *M. crudelis*, *M. distans* = *M. rusticus*, *M. agriope* = *M. rusticus* (see Banks, 1927:78); *M. diversus* = *M. rusticus* Stange, NEW SYNONYMY; lectotype designated by Banks, 1927:78.

DISTRIBUTION.—USA: AZ<sup>4</sup>, CA<sup>3</sup>, NM<sup>1</sup>, UT<sup>4</sup>, WY<sup>2</sup> (Hagen, 1861:233; Hagen, 1873:729; Banks, 1898:206; Banks, 1927:78). (A New Jersey record by Smith, 1900:56 must certainly be a misidentification).

**Myrmeleon texanus** Banks, 1900:596.

Holotype (sex unknown): USA, Texas, Galveston, VI.1900 (MCZ).

DISTRIBUTION.—USA: TX (Banks, 1900:596).

### POLYSTOECHOTIDAE

Polystoechotids, or giant lacewings, are among the largest of North American neuroptera, and among the least known. Only the first instar larva has been described (Welch, 1914), and no one has yet been able to get the young larvae to feed. Pupal skins of *Platystoechotes* have been found around the roots of incense cedar trees in the Sierra Nevada Mountains of California. Adult *Platystoechotes* emerge in

early summer, while adult *Polystoechotes* emerge in late summer and early fall. Adult *Polystoechotes* are most frequently seen around the smoke of campfires, and sometimes are attracted to lights. The geographical distribution of *Platystoechotes* is entirely in the Sierra Nevada of California. The range of *Polystoechotes* at one time included almost all of northern North America and much of California south to Panama. However, this genus cannot now be found over much of its former range, and today appears confined to remote mountainous areas.

### **Platystoechotes** Carpenter, 1940:270

**Platystoechotes lineatus** Carpenter, 1940:271.

Holotype male: USA, California, Wolverton, Sequoia National Park, 7000–9000', 14.VI.1929, E. C. Van Dyke (CAS).

DISTRIBUTION.—USA: CA (Carpenter, 1940:271).

### **Polystoechotes** Burmeister, 1839:982

**Polystoechotes punctatus** (Fabricius), 1793:73 [Semblis].

Holotype (sex unknown): without locality data (BMNH).

= *Hemerobius nebulosus* Fabricius, 1798:202. Type(s) (sex unknown): (locality unknown), (depository unknown).

= *Hemerobius irroratus* Say, 1824:306. Type(s) (sex unknown): USA (depository unknown).

= *Hemerobius vittatus* Say, 1824:307. Type(s) (sex unknown): USA: New Jersey; Pennsylvania (ANSP).

= *Polystoechotes sticticus* Burmeister, 1839:982. Type(s) (sex unknown): North America (Halle).

= *Osmylus validus* Walker, 1853:233. 5 syntypes (sex unknown): USA (BMNH).

TAXONOMY.—To *Polystoechotes* (see Hagen, 1861:206); *H. nebulosa* = *P. punctatus*, *H. irroratus* = *P. punctatus*, *P. sticticus* = *P. punctatus*, *O. validus* = *P. punctatus* (see Hagen, 1861:206); *H. vittatus* = *P. punctatus* (see Carpenter, 1940:268269).

DISTRIBUTION.—CAN: AB<sup>7</sup>, BC<sup>7</sup>, ON<sup>7</sup>, PQ<sup>7</sup>, USA: AK<sup>11</sup>, AZ<sup>7</sup>, CA<sup>7</sup>, CO<sup>2</sup>, GA<sup>7</sup>, IA<sup>11</sup>, ID<sup>7</sup>, IN<sup>10</sup>, KS<sup>4</sup>, MA<sup>7</sup>, MD<sup>7</sup>, ME<sup>7</sup>, MI<sup>11</sup>, MN<sup>8</sup>,

MT<sup>7</sup>, NC<sup>6</sup>, NE<sup>11</sup>, NH<sup>7</sup>, NJ<sup>12</sup>, NM<sup>2</sup>, NV<sup>7</sup>, NY<sup>5</sup>, OH<sup>11</sup>, OR<sup>7</sup>, PA<sup>3</sup>, SD<sup>11</sup>, TX<sup>1</sup>, UT<sup>7</sup>, WA<sup>7</sup>, WS<sup>9</sup>, WY<sup>7</sup> (<sup>1</sup>Hagen, 1861:206; <sup>2</sup>Hagen, 1875:920; <sup>3</sup>Smith, 1900:55; <sup>4</sup>Smith, 1925:167; <sup>5</sup>Leonard, 1928:40; <sup>6</sup>Brimley, 1938:30; <sup>7</sup>Carpenter, 1940:269; <sup>8</sup>Parfin, 1952:425; <sup>9</sup>Throne, 1971b:86; <sup>10</sup>Lawson & McCafferty, 1984:130; <sup>11</sup>Penny, NEW STATE RECORDS; <sup>12</sup>S. Marshall, NEW STATE RECORD).

#### SISYRIDAE

Sisyrids, or spongilla-flies, are larval predators of freshwater sponges, sucking out the liquid contents of the cells with their long, straight, hollow mandibles. This is the only truly aquatic family of Neuroptera, although some Osmylidae are semi-aquatic. The larvae bear external, abdominal gills in second and third instars. When mature, larvae leave the water and climb nearby vegetation, where they spin a silken cocoon. Often, the cocoon is surrounded by a distinctive, outer, mesh-like covering. Adults usually emerge in late summer, and are often attracted to lights, sometimes at a considerable distance from water. Eggs are laid above water, and when they hatch, the young larvae fall directly into the water. The taxonomy of North American spongilla-flies was dealt with by Parfin and Gurney (1956).

#### *Climacia* McLachlan, 1869a:21

***Climacia areolaris*** (Hagen), 1861:199 [*Micromus*].

Neotype male: USA Oklahoma, Grant, 1.VII.1937, Standish and Kaiser (MCZ).

= *Climacia dictyona* Needham, 1901:558.

Lectotype male: USA, New York, Saranac Inn, 28.VI.1900 (Cornell).

= *Sisyra lampra* Navás, 1914c:60. Type(s) (sex unknown): USA, New Jersey, Lakehurst (depository unknown).

TAXONOMY.—To *Climacia* (see McLachlan, 1869:21); *S. lampra* = *C. areolaris* (see Navás, 1935:33); *C. dictyona* = *C. areolaris*, *S. lampra* = *S. vicaria* (see Carpenter, 1940:254, 255); *S. lampra* = *C. areolaris* (see Parfin & Gurney, 1956:486). Hagen's original type was apparently lost in shipment from Europe, and Carpenter (1940:256) designated a neotype. Parfin and

Gurney (1956:488) designated a lectotype from Needham's syntype series of *C. dictyona*.

DISTRIBUTION.—CAN: ON<sup>4</sup>, PQ<sup>4</sup>, USA: AL<sup>7</sup>, AR<sup>7</sup>, CO<sup>7</sup>, CT<sup>4</sup>, FL<sup>1</sup>, GA<sup>4</sup>, IL<sup>4</sup>, IN<sup>7</sup>, LA<sup>4</sup>, MA<sup>4</sup>, MD<sup>4</sup>, ME<sup>4</sup>, MI<sup>4</sup>, MN<sup>6</sup>, MO<sup>5</sup>, MS<sup>4</sup>, NC<sup>2</sup>, NH<sup>4</sup>, NJ<sup>4</sup>, NM<sup>4</sup>, NY<sup>3</sup>, OH<sup>4</sup>, OK<sup>4</sup>, PA<sup>4</sup>, TX<sup>4</sup>, VA<sup>4</sup>, VT<sup>4</sup>, WS<sup>7</sup> (<sup>1</sup>Hagen, 1861:199; <sup>2</sup>Brimley, 1938:29; <sup>3</sup>Leonard, 1928:39; <sup>4</sup>Carpenter, 1940:256; <sup>5</sup>Froeschner, 1947:132; <sup>6</sup>Parfin, 1952:424; <sup>7</sup>Parfin & Gurney, 1956:489).

#### *Climacia californica* Chandler, 1953:183.

Holotype male: USA, California, Lake Co., Clear Lake, 19.V.1949, 1318 ft., H. P. Chandler (CAS).

DISTRIBUTION.—USA: CA<sup>1</sup>, ID<sup>2</sup> (<sup>1</sup>Chandler, 1953:182; <sup>2</sup>Clark, 1985:391).

#### *Climacia chapini* Parfin and Gurney, 1956:495.

Holotype male: USA, Texas, Columbus, E. A. Schwarz (USNM).

DISTRIBUTION.—USA: NM<sup>1</sup>, OK<sup>2</sup>, TX<sup>1</sup> (<sup>1</sup>Parfin & Gurney, 1956:498; <sup>2</sup>Penny, NEW STATE RECORD).

#### *Sisyra* Burmeister, 1839:975

##### *Sisyra apicalis* Banks, 1908a:261.

Holotype female: Cuba, Hayana (MCZ).

DISTRIBUTION.—USA: FL<sup>1</sup>, GA<sup>1</sup>, LA<sup>2</sup>, MS<sup>3</sup> (<sup>1</sup>Parfin & Gurney, 1956:472; <sup>2</sup>Poirier, 1969:574; <sup>3</sup>Lago, 1981:28).

##### *Sisyra fuscata* (Fabricius), 1793:84 [*Hemerobius*].

Holotype female: Denmark (ZMC).

= *Hemerobius confinis* Stephens, 1836:115.

Type(s) (sex unknown): England (BMNH).

= *Sisyra morio* Burmeister, 1839:976. Type(s) (sex unknown): Germany (Halle).

= *Sisyra nigripennis* Wesmael, 1841:213.

Type(s) (sex unknown): Belgium (IRSNB).

= *Branchiota spongillae* Westwood, 1842:108. Type(s) (sex unknown): larva, England (Oxford).

= *Hemerobius fumatus* Motschulsky, 1853:20.

Type(s) (sex unknown): Russia (Moscow).

TAXONOMY.—To *Sisyra* (see Burmeister, 1839:976); *H. nitidulus* = *S. fuscata*, *H. confinis* = *S. fuscata*, *S. morio* = *S. fuscata*, *S. nigripennis* = *S. fuscata*, *B. spongillae* = *S. fuscata*, *H. fumatus* = *S. fuscata* (see Parfin & Gurney, 1956:468).

DISTRIBUTION.—CAN: BC<sup>1</sup>, ON<sup>1</sup>, PQ<sup>1</sup>, USA: AK<sup>1</sup>, IN<sup>2</sup>, MA<sup>1</sup>, ME<sup>1</sup>, MI<sup>1</sup>, MN<sup>1</sup>, NY<sup>1</sup>,

WS<sup>1</sup> (<sup>1</sup>Parfin & Gurney, 1956:470; <sup>2</sup>Lawson & McCafferty, 1984:130).

**Sisyra vicaria** (Walker), 1853:297 [*Hemero-bius*].

Lectotype male: USA, Georgia, John Abbot(BMNH).

= *Sisyra umbrata* Needham, 1901:555. Lectotype male: USA, Illinois, Lake Forest, 2.VI.1899 (Cornell).

TAXONOMY.—To *Sisyra* (see Hagen, 1861:197); *S. umbrata* = *S. vicaria*, *S. lampra* = *S. vicaria* (see Carpenter, 1940:254); *S. lampra* not = *S. vicaria* (see Parfin & Gurney, 1956:486).

DISTRIBUTION.—CAN: BC<sup>5</sup>, NS<sup>5</sup>, ON<sup>5</sup>, PQ<sup>5</sup>, USA: AZ<sup>9</sup>, CT<sup>5</sup>, DC<sup>5</sup>, FL<sup>5</sup>, GA<sup>1</sup>, IL<sup>5</sup>, IN<sup>9</sup>, KS<sup>2</sup>, KY<sup>9</sup>, MA<sup>5</sup>, MD<sup>5</sup>, ME<sup>6</sup>, MI<sup>5</sup>, MN<sup>8</sup>, MO<sup>7</sup>, MS<sup>10</sup>, NC<sup>4</sup>, NY<sup>3</sup>, OK<sup>9</sup>, OR<sup>5</sup>, PA<sup>5</sup>, RI<sup>9</sup>, TN<sup>9</sup>, TX<sup>5</sup>, VA<sup>9</sup>, WA<sup>9</sup>, WS<sup>5</sup>, (<sup>1</sup>Walker, 1853:297; <sup>2</sup>Smith, 1925:166; <sup>3</sup>Leonard, 1928:39; <sup>4</sup>Brimley, 1938:29; <sup>5</sup>Carpenter, 1940:254; <sup>6</sup>Procter, 1946:42; <sup>7</sup>Froeschner, 1947:132; <sup>8</sup>Parfin, 1952:424; <sup>9</sup>Parfin & Gurney, 1956:464; <sup>10</sup>Poirier & Holzenthal, 1980:1).

#### RAPHIDIOPTERA

Snake flies are elongate as larvae, living as predators under the surface of tree bark, under fallen logs, or in leaf litter. Adults may be associated with a variety of coniferous and broad-leaved trees and shrubs. Life histories of most of our species are still unknown. In America north of Mexico, they do not occur east of the Rocky Mountains, except in Texas. The taxonomy of all species of Raphidioptera was treated by Aspöck (1974, 1975), and H. and U. Aspöck and Rausch (1991).

#### INOCELLIDAE

Inocellid snake-flies can be separated from raphidiids by the absence of ocelli, longer thicker antennae, and darker thicker pterostigmal spot. Only a single genus, *Negha* Navás 1916b, is found in North America.

**Negha** Navás, 1916b:510

**Negha inflata** (Hagen), 1861:196 [*Raphidia*].

Neotype male: USA, California, Yosemite National Park, Mariposa Grove, 12.VII.1927, F. M. Carpenter (MCZ).

= *Inocellia hageni* Albarda, 1891:171. Holotype female: USA, California, San Francisco (BMNH).

TAXONOMY.—To *Inocellia* (see Albarda, 1891:167); to *Negha* (see Navás, 1916b:510); returned to *Inocellia* (see Carpenter, 1936:138); to *Inocellia* (*Negha*) (see Aspöck, 1975:540); returned to *Negha* (see Aspöck, 1988:107); *I. hageni* = *I. inflata* (see Carpenter, 1936:139); *I. longicornis* = *I. inflata* (see Aspöck, 1975:540).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: CA<sup>1</sup>, ID<sup>3</sup>, MT<sup>3</sup>, NV<sup>2</sup>, OR<sup>2</sup>, UT<sup>3</sup>, WA<sup>3</sup> (<sup>1</sup>Hagen, 1861:196; <sup>2</sup>Carpenter, 1936:140; <sup>3</sup>Aspöck, 1975:540).

**Negha longicornis** (Albarda), 1891:169 [*Inocellia*].

Holotype male: USA, California (BMNH).

TAXONOMY.—To *Negha* (see Navás, 1919a:75); returned to *Inocellia* (see Carpenter, 1936:140); returned to *Negha* (see Aspöck, 1988:107); *I. longicornis* = *I. inflata* (see Aspöck, 1975:540); *I. longicornis* not = *I. inflata* (see Aspöck, 1988:107).

DISTRIBUTION.—CAN: BC<sup>2</sup>, USA: CA<sup>1</sup>, NV<sup>2</sup>, OR<sup>2</sup>, WA<sup>2</sup> (<sup>1</sup>Albarda, 1891:169; <sup>2</sup>Carpenter, 1936:141).

**Negha meridionalis** U. Aspöck, 1988:107.

Holotype male: USA, California, Riverside Co., 35 miles south of Palm Springs, 26.II.1977, S. Johnson (SDNRM).

DISTRIBUTION.—USA: CA (Aspöck, 1988:107).

#### RAPHIDIIDAE

The biology of raphidiids is similar to that of inocellids. For subgeneric characters, see H. and U. Aspöck and Rausch (1991).

**Agulla** (*Agulla*) Navás, 1914d:66

**Agulla** (*Agulla*) **arnaudi** (U. Aspöck), 1973:234 [*Raphidia* (*Agulla*)].

Holotype male: Mexico, Baja California, San Pedro Martir, trail La Joya to La Zanja, 10.6.1953, P. H. Arnaud, Jr. (CAS).

TAXONOMY.—To *Agulla* (see Aspöck, 1986:25).

DISTRIBUTION.—USA: CA (Aspöck, 1973:234).

**Agulla** (*Agulla*) **assimilis** (Albarda), 1891:144 [*Raphidia*].

Holotype female: Canada, British Columbia, Vancouver Island (BMNH).

= *Raphidia arizonica* (Banks), 1911:338. Holotype male: USA, Arizona, Phoenix, Oslar (MCZ).

= *Agulla bagnalli* Navás, 1914d:67. Holotype male: Canada, British Columbia, Vancouver Island, Victoria (Oxford).

TAXONOMY.—To *Agulla* (see Navás, 1914d:66); to *Raphidia* (*Agulla*) (see Aspöck, 1975:538); returned to *Agulla* (see Aspöck, 1986:25); *A. bagnalli* = *R. assimilis* (Navás, 1918c:26); *R. arizonica* = *R. assimilis* (Aspöck, 1975:538).

DISTRIBUTION.—CAN: AB<sup>3</sup>, BC<sup>1</sup> USA: AZ<sup>2</sup>, CA<sup>3</sup>, CO<sup>4</sup>, ID<sup>4</sup>, MT<sup>4</sup>, OR<sup>3</sup>, UT<sup>4</sup>, WA<sup>4</sup>, WY<sup>4</sup> (<sup>1</sup>Albarda, 1891:144; <sup>2</sup>Banks, 1911:338, as *R. arizonica*; <sup>3</sup>Carpenter, 1936:128; <sup>4</sup>Carpenter, 1936:118, as *R. arizonica*).

**Agulla (Agulla) astuta** (Banks), 1911:338 [*Raphidia*].

Three syntype males: USA, California, Pasadena, 31.V.1919, F. Grinnell, Jr.; San Gabriel Mountains, 17.VI; Mount Wilson, 5.VI.1904 (MCZ).

= *Agulla singularis* Carpenter, 1936:132. Holotype male: USA, California, Banning, 29.V.1928, E. C. Van Dyke (CAS).

TAXONOMY.—To *Glavia* (see Navás, 1919a:30); to *Agulla* (*Agulla*) (see Carpenter, 1936:131); to *Raphidia* (*Agulla*) (see Aspöck, 1975:538); returned to *Agulla* (see Aspöck, 1986:25); *A. singularis* = *A. astuta* (see Aspöck, 1975:538).

DISTRIBUTION.—USA: AZ<sup>2</sup>, CA<sup>1</sup> (<sup>1</sup>Banks, 1911:338; <sup>2</sup>Aspöck, 1975:538).

**Agulla (Agulla) barri** (U. Aspöck), 1973:238 [*Raphidia* (*Agulla*)].

Holotype male: USA, California, Inyo Co., Westgard Pass, 11.6.1969, W. F. Barr, on pinyon pine (CAS).

TAXONOMY.—To *Agulla* (see Aspöck, 1986:25).

DISTRIBUTION.—USA: CA, NV (Aspöck, 1973:239).

**Agulla (Agulla) bicolor** (Albarda), 1891:152 [*Raphidia*].

Ten syntypes (four males, six females): USA, Colorado (BMNH).

= *Raphidia occulta* Banks, 1905a:88 (nomen nudum).

= *Raphidia occulta* Banks, 1905b:4. Syntypes (sex unknown): USA, California, Claremont, Baker; Arizona, Prescott, Oslar; New Mexico, Pecos (MCZ).

TAXONOMY.—To *Raphidilla* (see Navás, 1919a:51); to *Agulla* (*Agulla*) (see Carpenter, 1936:126); to *Raphidia* (*Agulla*) (see Aspöck, 1975:538); returned to *Agulla* (see Aspöck, 1986:25); *R. occulta* = *A. bicolor* (see Carpenter, 1936:126).

DISTRIBUTION.—CAN: BC<sup>3</sup>, USA: AZ<sup>2</sup>, CA<sup>2</sup>, CO<sup>1</sup>, NM<sup>2</sup>, NV<sup>3</sup>, OR<sup>3</sup>, UT<sup>3</sup>, WA<sup>3</sup> (<sup>1</sup>Albarda, 1891:152; <sup>2</sup>Banks, 1905b:4; <sup>3</sup>Carpenter, 1936:126).

**Agulla (Agulla) bractea** Carpenter, 1936:130.

Holotype male: USA, California, San Gabriel Mountains, 3500', 11.VI.1910, F. Grinnell, Jr. (MCZ).

= *Agulla neglecta* Carpenter, 1936:132. Holotype male: USA, California, East Lake, VI.1883, Turner (USNM).

TAXONOMY.—To *Raphidia* (*Agulla*) (see Aspöck, 1975:538); returned to *Agulla* (see Aspöck, 1986:25); *A. neglecta* = *A. bractea* (see Aspöck, 1975:538).

DISTRIBUTION.—USA: CA (Carpenter, 1936:130).

**Agulla (Agulla) crotchi** (Banks), 1924:429 [*Raphidia*].

Holotype male: USA, California, Crotch (MCZ).

TAXONOMY.—To *Agulla* (*Agulla*) (see Carpenter, 1936:123).

DISTRIBUTION.—USA: CA (Banks, 1924:429).

**Agulla (Agulla) faulkneri** U. Aspöck, 1987:1.

Holotype male: USA, California, Riverside Co., Idylwild-Pine Cove, 26.VI.1977, J. W. Brown (SDNHM).

DISTRIBUTION.—USA: CA (Aspöck, 1987:1).

**Agulla (Agulla) flexa** Carpenter, 1936:125.

Holotype male: USA, New Mexico, Jemez Mountains, 18.VI, Woodgate (MCZ).

TAXONOMY.—To *Raphidia* (*Agulla*) (see Aspöck, 1975:539); returned to *Agulla* (see Aspöck, 1986:25).

DISTRIBUTION.—USA: AZ, NM (Carpenter, 1936:125).

**Agulla (Agulla) herbsti** (Esben-Petersen), 1912:273 [*Raphidia*].

Holotype male: "Chile, Concepcion," 29.XII.1908, P. Herbst (HMB). Note: This locality is certainly in error, the species is only found in North America.

= *Raphidia bifurca* Banks, 1920:327. Syntypes male and female: Canada, British Columbia, Wellington, 9.V.1898; 3.V.1897 (MCZ).

= *Agulla occidentis* Carpenter, 1936:129. Holotype male: USA, California, Pinecrest, Johnson (CAS).

TAXONOMY.—To *Raphidia (Agulla)* (see Aspöck, 1975:538); *R. bifurca* = *R. herbsti*, *A. occidentis* = *R. herbsti* (see Aspöck, 1975:538).

DISTRIBUTION.—CAN: AB<sup>3</sup>, BC<sup>1</sup>, USA: CA<sup>2</sup>, ID<sup>3</sup>, OR<sup>2</sup>, MT<sup>3</sup>, NV<sup>3</sup>, WA<sup>3</sup> (Banks, 1920:327, as *R. bifurca*; Carpenter, 1936:130, as *A. occidentis*; Aspöck, 1975:539).

**Agulla (Californoraphidia)** Aspöck et al., 1991:464

**Agulla (Californoraphidia) nigrinotum** Woglum and McGregor, 1964:201.

Holotype male: USA, California, Kern Co., Mill Potrero, 1.VII.1963, R. S. Woglum (CAS).

TAXONOMY.—To *Raphidia (Agulla)* (see Aspöck, 1975:539); to *Agulla* (see Aspöck, 1986:25); to *Agulla (Californoraphidia)* by Aspöck et al., 1991:464.

DISTRIBUTION.—USA: CA (Woglum & McGregor, 1964:202).

**Agulla (Franciscoraphidia)** Aspöck et al., 1991:462

**Agulla (Franciscoraphidia) directa** Carpenter, 1936:124 [*Agulla (Agulla)*].

Holotype male: USA, California, Oroville, 17.IV.1928, H. Kiefer (CAS).

TAXONOMY.—To *Raphidia (Agulla)* (see Aspöck, 1975:539); returned to *Agulla* (see Aspöck, 1986:25); to *Agulla (Franciscoraphidia)* (see Aspöck et al., 1991:462).

DISTRIBUTION.—USA: CA (Carpenter, 1936:125).

**Agulla (Glavia)** Navás, 1916b:?

**Agulla (Glavia) adnixa** (Hagen), 1861:195 [*Raphidia*].

Lectotype female: USA, California; Oregon, Willcox (HMB).

= *Raphidia oblita* Hagen, 1861:195. Holotype male (destroyed): USA, California.

TAXONOMY.—To *Glavia* (see Navás, 1916b:509); to *Agulla (Agulla)* (see Carpenter, 1936:116); to *Raphidia (Agulla)* (see Aspöck, 1975:539); to *Agulla (Glavia)* (see Aspöck et al., 1991:450); *R. oblita* = *A. adnixa* (see Carpenter, 1936:117); lectotype designated (see Aspöck, et al., 1991:452).

DISTRIBUTION.—CAN: AB<sup>2</sup>, BC<sup>2</sup>, USA: AZ<sup>2</sup>, CA<sup>1</sup>, CO<sup>2</sup>, ID<sup>2</sup>, MT<sup>2</sup>, NM<sup>2</sup>, NV<sup>2</sup>, OR<sup>1</sup>, UT<sup>2</sup>, WA<sup>2</sup>, WY<sup>2</sup> (Hagen, 1861:195; Carpenter, 1936:116).

**Agulla (Glavia) modesta adryte** U. Aspöck, 1982:98.

Holotype male: USA, California, Inyo Co., 7 miles north of Parcher's Camp, 30.VI.1961, J. K. Drew (UCB).

DISTRIBUTION.—USA: CA (Aspöck, 1982:98).

**Agulla (Glavia) modesta aphynphte** U. Aspöck, 1982:102.

Holotype male: USA, California, Los Angeles Co., Westwood Hills, IV-V.1949 (LACM).

DISTRIBUTION.—USA: CA (Aspöck, 1982:102).

**Agulla (Glavia) modesta aphyrite** U. Aspöck, 1982:100.

Holotype male: USA, Arizona, Coconino Co., Oak Creek Canyon, 26.VI.1950, L. D. Beamer (SEM).

DISTRIBUTION.—USA: AZ, NM (Aspöck, 1982:100).

**Agulla (Glavia) modesta banksi** Carpenter, 1936:119 [*Agulla (Agulla)*].

Holotype male: USA, California, Big Bear Lake, 26.VII.1932, R. H. Beamer (MCZ).

TAXONOMY.—To *Raphidia (Agulla)* (see Aspöck, 1975:539); to *Agulla (Glavia)* (see Aspöck, 1988:98); *A. banksi* = *R. modesta* (see Aspöck, 1975:539); *A. banksi* = subspecies of *A. modesta* (see Aspöck, 1988:98).

DISTRIBUTION.—USA: CA (Carpenter, 1936:120).

**Agulla (Glavia) modesta modesta** Carpenter, 1936:119 [*Agulla (Agulla)*].

Holotype male: USA, Utah, Beaver Canyon (USNM).

**TAXONOMY.**—To *Raphidia (Agulla)* (see Aspöck, 1975:539); to *Agulla (Glavia)* (see Aspöck, 1982:97).

**DISTRIBUTION.**—USA: CO<sup>2</sup>, UT<sup>1</sup> (<sup>1</sup>Carpenter, 1936:120; <sup>2</sup>Aspöck, 1982:98).

**Agulla (Glavia) paramerica** U. Aspöck, 1982:104.

Holotype male: USA, California, Plumas Co., Johnsville, 11.VII–27.VIII.1964, H. Pini (UCD).

**DISTRIBUTION.**—USA: CA (Aspöck, 1982:104).

**Agulla (Glavia) unicolor** Carpenter, 1936:121 [*Agulla (Agulla)*].

Holotype male: USA, California, Yosemite National Park, Mariposa Grove, 20.VII.1932, F. M. Carpenter (MCZ).

**TAXONOMY.**—To *Raphidia (Agulla)* (see Aspöck, 1975:539); to *Agulla (Glavia)* (see Aspöck, 1982:102).

**DISTRIBUTION.**—CAN: BC, USA: CA, ID, MT, NV, OR, UT, WA (Carpenter, 1936:121).

**Alena (Alena)** Navás, 1916b:510

**Alena (Alena) distincta** (Banks), 1911:338 [*Raphidia*].

Holotype male: USA, California, Kern Co., Mt. Pinos, 6.VI.1904, F. Grinnell, Jr. (MCZ).

**TAXONOMY.**—To *Alena* (see Navás, 1916b:510); to *Agulla (Alena)* (see Carpenter, 1936:134); to *Raphidia (Alena)* (see Aspöck, 1975:539); returned to *Alena* (see Aspöck, 1986:25).

**DISTRIBUTION.**—USA: CA (Banks, 1911:338).

**Alena (Aztekoraphidia)** U. Aspöck and H. Aspöck, 1970:709

**Alena (Aztekoraphidia) minuta** (Banks), 1903b:239 [*Raphidia*].

Holotype male: USA, Arizona, Williams, 1.VI, Barber and Schwarz (USNM).

**TAXONOMY.**—To *Alena* (see Navás, 1916b:510); to *Agulla (Alena)* (see Carpenter, 1936:133); to *Raphidia (Sombreroraphidia)* (see Aspöck, 1975:539); to *Alena (Aztekoraphidia)* (see Aspöck et al., 1991:471).

**DISTRIBUTION.**—USA: AZ, NM (Banks, 1903b:239).

## MEGALOPTERA CORYDALIDAE

Corydalidae, or dobson-flies and fish-flies, are among our largest neuropteroids. Males of the dobson-fly *Corydalus cornutus* can have a wing-span of five and one-half inches (14 cm) and bear elongate, sickle-shaped mandibles more than one and one-half inches (4 cm) long. The larvae are aquatic predators known as hellgrammites, which feed on aquatic insects, tadpoles, and small fish. Larvae may go through seven or eight instars and take two or three years or longer to reach maturity. Larvae of some species of *Protochauliodes* live in intermittent streams and form a cell in the soil of the stream bed during the months when the stream is dry. Despite their ferocious appearance, adults either do not feed, or feed on small quantities of nectar and fruit juices. The taxonomy of this group has been treated by Chandler (1956), Contreras-Ramos (1996), Evans (1984), Flint (1965), and Merritt and Cummins (1978).

## CORYDALINAE

**Corydalus** Latreille, 1802:290

**Corydalus cornutus** (Linnaeus), 1758:551 [*Heremobius*].

Type(s) (sex unknown): USA, Pennsylvania, DeGeer (London).

= *Corydalus cognatus* Hagen, 1861:193 (as *Corydalis [sic] cognata*). Holotype female: USA, western Texas (now New Mexico), Pecos River, 4.VII, Pope (MCZ).

= *Corydalus luteus* Hagen, 1861:193 (as *Corydalis [sic] lutea*). Syntype males: Mexico, Salle; Cordova, Saussure, one in Selys Collection (IRSNB).

= *Corydalus crassicornis* McLachlan, 1867:233 (as *Coryalis [sic] crassicornis*). Holotype (sex unknown): USA, Texas (MNHN).

= *Corydalus inamabilis* McLachlan, 1867:235 (as *Corydalis [sic] inamabilis*). Type(s) (sex unknown): USA, Texas (BMNH).

= *Nevromus pallidus* Davis, 1903:470 (as *Neuromus [sic] pallidus*). Holotype male: (locality and depository unknown).

= *Corydalus texanus* Banks, 1903:239 (as *Corydalis [sic] texana*). Holotype female: USA, Texas (depository unknown).

TAXONOMY.—To *Raphidia* (see Linnaeus, 1767:916); returned to *Hemerobius* (see DeGeer, 1773:559); to *Corydalus* (see Latreille, 1802:290); *C. crassicornis* = *C. cornutus*, *C. inamabilis* = *C. cornutus*, *C. cognata* = *C. cornutus*, *C. lutea* = *C. cornutus*, *C. pallidus* = *C. cornutus*, *C. texana* = *C. cornutus* (see Weele, 1910:13); *C. cognatus* not = *C. cornutus* (see Evans, 1972:77).

DISTRIBUTION.—CAN: PQ<sup>4</sup>, USA: AZ<sup>5</sup>, CA<sup>7</sup>, DC<sup>4</sup>, IA<sup>8</sup>, IL<sup>2</sup>, IN<sup>6</sup>, KS<sup>4</sup>, MD<sup>2</sup>, MN<sup>4</sup>, MO<sup>4</sup>, NC<sup>4</sup>, NM<sup>5</sup>, NV<sup>7</sup>, NY<sup>2</sup>, OH<sup>4</sup>, PA<sup>1</sup>, SC<sup>4</sup>, TX<sup>3</sup>, UT<sup>7</sup>, VA<sup>4</sup> (<sup>1</sup>Linnaeus, 1758:551; <sup>2</sup>Hagen, 1861:192; <sup>3</sup>McLachlan, 1867:233; <sup>4</sup>Davis, 1903:478, as *C. cornutus*; <sup>5</sup>Davis, 1903:479, as *C. cognata*; <sup>6</sup>Montgomery & Trippel, 1933:259; <sup>7</sup>Evans, 1972:78, as *C. cognata*; <sup>8</sup>Penny, NEW STATE RECORD).

#### CHAULIODINAE

##### **Chauliodes** Latreille, 1796:102

**Chauliodes pectinicornis** (Linnaeus), 1763:412 [*Hemerobius*].

Type(s) (sex unknown): (locality unknown) (Tring).

= *Hemerobius virginensis* Drury, 1773: unpaginated index. Type(s) (sex unknown):(locality unknown) (BMNH).

TAXONOMY.—To *Semblis* (see Fabricius, 1781:386); to *Chauliodes* (see Latreille, 1802:290); *H. virginensis* = *C. pectinicornis* (see Davis, 1903:461).

DISTRIBUTION.—CAN: BC<sup>7</sup>, USA: DE<sup>2</sup>, FL<sup>4</sup>, IA<sup>9</sup>, IL<sup>9</sup>, IN<sup>5</sup>, KS<sup>9</sup>, LA<sup>4</sup>, MA<sup>2</sup>, MD<sup>3</sup>, ME<sup>8</sup>, MI<sup>4</sup>, MO<sup>4</sup>, MS<sup>9</sup>, NC<sup>6</sup>, NE<sup>9</sup>, NH<sup>9</sup>, NJ<sup>3</sup>, NY<sup>4</sup>, OH<sup>4</sup>, PA<sup>3</sup>, SC<sup>4</sup>, RI<sup>9</sup>, TX<sup>9</sup>, VA<sup>1</sup>, VT<sup>9</sup>, WS<sup>4</sup> (<sup>1</sup>Drury, 1773:2; <sup>2</sup>Walker, 1853:198; <sup>3</sup>Hagen, 1861:189; <sup>4</sup>Davis, 1903:462; <sup>5</sup>Montgomery & Trippel, 1933:259; <sup>6</sup>Brimley, 1938:28; <sup>7</sup>Spencer, 1942:23; <sup>8</sup>Procter, 1946:42; <sup>9</sup>Tarter, Watkins et al., 1976:223)

**Chauliodes rastricornis** Rambur, 1842:444.

Holotype female: no published locality data, Selys Collection (IRSNB).

= *Hermes indecisus* Walker, 1853:204. Type(s) (sex unknown): no locality data (BMNH).

TAXONOMY.—*H. indecisus* = *C. rastricornis* (see Davis, 1903:460).

DISTRIBUTION.—USA: AL<sup>7</sup>, FL<sup>2</sup>, GA<sup>2</sup>, IA<sup>7</sup>, IL<sup>1</sup>, IN<sup>2</sup>, KS<sup>3</sup>, ME<sup>5</sup>, MN<sup>6</sup>, MO<sup>2</sup>, MS<sup>7</sup>, NC<sup>4</sup>, NE<sup>7</sup>, NH<sup>7</sup>, NY<sup>2</sup>, PA<sup>2</sup>, RI<sup>7</sup>, SC<sup>2</sup>, SD<sup>7</sup>, TN<sup>7</sup>, VA<sup>7</sup>, WS<sup>7</sup> (<sup>1</sup>Hagen, 1863:181; <sup>2</sup>Davis, 1903:460; <sup>3</sup>Smith, 1925:166; <sup>4</sup>Brimley, 1938:28; <sup>5</sup>Procter, 1946:42; <sup>6</sup>Parfin, 1952:422; <sup>7</sup>Tarter, Watkins, et al., 1976:224).

##### **Dysmicohermes** Munroe, 1953:191

**Dysmicohermes disjunctus** (Walker), 1866:334 [*Chauliodes*].

Holotype female: Canada, British Columbia, Vancouver Island, Chalukweyuh Lake, IX.1859 (BMNH).

TAXONOMY.—To *Neohermes* (see Weele, 1910:54); to *Dysmicohermes* (see Munroe, 1953:190).

DISTRIBUTION.—CAN: BC<sup>1</sup>, USA: CA<sup>2</sup>, OR<sup>2</sup>, WA<sup>2</sup> (<sup>1</sup>Walker, 1866:334; <sup>2</sup>Evans, 1972:87).

**Dysmicohermes ingens** Chandler, 1954:105.

Holotype male: USA, California, Mariposa Co., Miami Ranger Station, 27.VII.1946, 5000 ft., H. P. Chandler (CAS).

DISTRIBUTION.—USA: CA (Chandler, 1954:106).

##### **Neohermes** Banks, 1908d:29

**Neohermes angusticollis** (Hagen), 1861:191 [*Chauliodes*].

Neotype male: USA, Georgia, Atlanta, 11.VI.1939, P. W. Fattig (USNM).

TAXONOMY.—To *Neohermes* (see Banks, 1908d:29); *N. angusticollis* = *N. californicus* (see Weele, 1910:53); *N. angusticollis* is valid species (see Flint, 1965:259); neotype designated (see Flint, 1965:260).

DISTRIBUTION.—USA: GA<sup>1</sup>, SC<sup>2</sup> (<sup>1</sup>Flint, 1965:260; <sup>2</sup>Tarter et al., 1976:225).

**Neohermes californicus** (Walker), 1853:199 [*Chauliodes*].

Holotype female: USA, California (BMNH).

TAXONOMY.—To *Neohermes* (see Banks, 1908d:29).

DISTRIBUTION.—USA: CA<sup>1</sup>, NV<sup>2</sup> (<sup>1</sup>Walker, 1853:199; <sup>2</sup>Flint, 1965:258).

**Neohermes concolor** (Davis), 1903:462 [*Chauliodes*].

Lectotype female: (no locality data) (Cornell).

**TAXONOMY.** — *N. concolor* = *N. californicus* (see Weele, 1910:53); *N. concolor* is valid species (see Flint, 1965:262); lectotype designated (see Flint, 1965:262).

**DISTRIBUTION.** — USA: AR<sup>2</sup>, DC<sup>2</sup>, DE<sup>2</sup>, GA<sup>3</sup>, IL<sup>3</sup>, IN<sup>1</sup>, KS<sup>4</sup>, KY<sup>2</sup>, MA<sup>2</sup>, MD<sup>2</sup>, MO<sup>2</sup>, MS<sup>3</sup>, NC<sup>2</sup>, NJ<sup>2</sup>, NY<sup>2</sup>, OH<sup>3</sup>, OK<sup>3</sup>, PA<sup>2</sup>, TN<sup>3</sup>, VA<sup>2</sup>, VT<sup>3</sup> (<sup>1</sup>Montgomery & Trippel, 1933:259; <sup>2</sup>Flint, 1965:262; <sup>3</sup>Tarter, Watkins et al., 1976:225; <sup>4</sup>Roble, 1984:69).

**Neohermes filicornis** (Banks), 1903b:238 [*Chauliodes*].

Holotype male: USA, Arizona, Jerome, 24.VI.1902, Oslar (MCZ).

**TAXONOMY.** — To *Neohermes* (see Banks, 1908d:29); *N. filicornis* = *N. californicus* (see Weele, 1910:53); *N. filicornis* is valid name (see Van Dyke, 1945:110).

**DISTRIBUTION.** — USA: AZ<sup>1</sup>, CA<sup>2</sup>, NM<sup>2</sup> (<sup>1</sup>Banks, 1903b:238; <sup>2</sup>Flint, 1965:257).

**Neohermes matheri** Flint, 1965:260.

Holotype male: USA, Mississippi, Hinds Co., Clinton, 12.VI.1960, Bryant Mather (USNM).

**DISTRIBUTION.** — USA: MS (Flint, 1965:260).

**Nigronia** Banks, 1908d:30

**Nigronia fasciata** (Walker), 1853:201 [*Chauliodes*].

Two syntype males: USA, New York (BMNH).

= *Chauliodes lunatus* Hagen, 1863:180. Syntypes (sex unknown): USA, Illinois; Maryland; New York; Pennsylvania (depository unknown).

**TAXONOMY.** — To *Nigronia* (see Banks, 1908d:30); *C. lunatus* = *C. fasciatus* (see Davis, 1903:458).

**DISTRIBUTION.** — USA: AL<sup>6</sup>, AR<sup>3</sup>, DC<sup>3</sup>, DE<sup>6</sup>, IL<sup>2</sup>, LA<sup>6</sup>, MD<sup>2</sup>, ME<sup>5</sup>, MN<sup>3</sup>, MO<sup>3</sup>, MS<sup>1</sup>, NC<sup>4</sup>, NH<sup>6</sup>, NY<sup>1</sup>, OH<sup>3</sup>, PA<sup>2</sup>, SC<sup>6</sup> (<sup>1</sup>Walker, 1853:201; <sup>2</sup>Hagen, 1863:181; <sup>3</sup>Davis, 1903:458; <sup>4</sup>Brimley, 1938:28; <sup>5</sup>Procter, 1946:42; <sup>6</sup>Tarter, Watkins et al., 1976:224; <sup>7</sup>Stark & Lago, 1983:357).

**Nigronia serricornis** (Say), 1824:307 [*Chauliodes*].

Type(s) (sex unknown): USA: Arkansas; Minnesota, Lake of the Woods; Pennsylvania; Missouri; Canada: Saskatchewan, Red River of Lake Winnipeg (depository unknown).

= *Nevromus maculatus* Rambur, 1842:442. Holotype female: (no locality data) (IRSNB).

**TAXONOMY.** — To *Nigronia* (see Banks, 1908d:30); *C. maculatus* = *C. serricornis* (see Hagen, 1863:181).

**DISTRIBUTION.** — CAN: MB<sup>1</sup>, USA: AL<sup>9</sup>, AR<sup>1</sup>, CT<sup>8</sup>, DC<sup>2</sup>, FL<sup>11</sup>, GA<sup>1</sup>, IN<sup>4</sup>, KS<sup>3</sup>, LA<sup>8</sup>, MA<sup>1</sup>, MD<sup>1</sup>, ME<sup>6</sup>, MN<sup>7</sup>, MO<sup>1</sup>, MS<sup>10</sup>, NC<sup>5</sup>, NY<sup>1</sup>, OH<sup>2</sup>, PA<sup>1</sup>, RI<sup>8</sup>, SC<sup>8</sup> (<sup>1</sup>Hagen, 1861:191; <sup>2</sup>Davis, 1903:460; <sup>3</sup>Smith, 1925:166; <sup>4</sup>Montgomery & Trippel, 1933:259; <sup>5</sup>Brimley, 1938:28; <sup>6</sup>Procter, 1946:42; <sup>7</sup>Parfin, 1952:422; <sup>8</sup>Tarter, Watkins et al., 1976:224; <sup>9</sup>Scheiring, 1979: 176; <sup>10</sup>Stark & Lago, 1983:358; <sup>11</sup>Stange, NEW STATE RECORD).

**Orohermes** Evans, 1984:1

**Orohermes crepusculus** (Chandler), 1954:107 [*Dysmicohermes*].

Holotype male: USA, California, El Dorado Co., Pyramid Ranger Station, 20.VIII.1952, J. W. MacSwain (CAS).

**TAXONOMY.** — To *Orohermes* (see Evans, 1984:1).

**DISTRIBUTION.** — USA: CA, OR (Chandler, 1954:108).

**Protochauliodes** Weele, 1909b:257

**Protochauliodes aridus** Maddux, 1954:70.

Holotype male: USA, California, Butte Co., Neal Road, 7 miles southeast of Chico, 15.V.1951, D. Maddux (CAS).

**DISTRIBUTION.** — USA: CA (Maddux, 1954:70).

**Protochauliodes cascadius** Evans, 1984:1.

Holotype male: USA, Oregon, Marion Co., ca. 8 miles west of Mill City, along Little Santiam River, 26.VII.1963, S. Jewett (CAS).

**DISTRIBUTION.** — USA: OR (Evans, 1984:2).

**Protochauliodes minimus** (Davis), 1903:463 [*Chauliodes*].

Holotype male: USA, California, San Rafael, 26.V, O. Sacken (MCZ).

= *Neohermes infuscatus* Caudell, 1933:125. Holotype female: USA, California, Solano Co., Vacaville, 1.VI.1932, A. S. Harrison (USNM).

= *Neohermes nigrinus* Van Dyke, 1945:110. Holotype male: USA, California, Butte Co.,

Richardson's Spring, 12.V.1944, E. C. Van Dyke (CAS).

**TAXONOMY.**—To *Protochauliodes* (see Chandler, 1954:110); *C. minimus* = *N. californicus* (see Weele, 1910:53); *P. minimus* not = *N. californicus* (see Chandler, 1954:110); *N. infuscatus* = *P. minimus*, *N. nigrinus* = *N. minimus* (see Evans, 1972:128).

**DISTRIBUTION.**—**USA:** CA (Davis, 1903:463).

**Protochauliodes montivagus** Chandler, 1954:111.

Holotype male: USA, California, Sierra Co., St. Charles Hill, 7.VII.1921, E. H. Nast (CAS).

**DISTRIBUTION.**—**USA:** CA (Chandler, 1954:111).

**Protochauliodes simplus** Chandler, 1954:110.

Holotype male: USA, California, Los Angeles Co., San Dimas Experimental Forest, Tanbark Flat, 13.VII.1950, J. D. Paschke (CAS).

**DISTRIBUTION.**—**USA:** CA (Chandler, 1954:110).

**Protochauliodes spenceri** Munroe, 1953:190.

Holotype male: Canada, British Columbia, Duncan, 1.VIII.1918, W. Downes (CNC).

**DISTRIBUTION.**—**CAN:** BC<sup>1</sup>, **USA:** CA<sup>2</sup>, OR<sup>2</sup>, WA<sup>2</sup> (<sup>1</sup>Munroe, 1953:190; <sup>2</sup>Evans, 1972:134).

#### SIALIDAE

Sialidae, or alder-flies, as adults are rather uniform insects about one inch long, with smoky black wings. Head markings can vary from orange to black. They live along the margins of rivers, streams, and lakes, never getting more than several yards away from water. Larvae live in the bottom substrate of these same water courses, and often survive in heavily polluted waters. The immatures are predators, feeding on other aquatic arthropods. The taxonomy of *Sialis* Latreille, 1802, was treated by Ross (1937).

**Sialis** Latreille, 1802:290

**Sialis aequalis** Banks, 1920:326.

Lectotype male: USA, Virginia, Falls Church, 25.IV, N. Banks (MCZ).

**TAXONOMY.**—Lectotype designated (see Ross, 1937:77).

**DISTRIBUTION.**—**USA:** CT<sup>4</sup>, DE<sup>4</sup>, GA<sup>5</sup>, KY<sup>5</sup>, MD<sup>2</sup>, ME<sup>5</sup>, MI<sup>5</sup>, MN<sup>4</sup>, NC<sup>2</sup>, NJ<sup>2</sup>, NY<sup>4</sup>,

OH<sup>4</sup>, PA<sup>2</sup>, SC<sup>4</sup>, VA<sup>1</sup>, VT<sup>5</sup>, WV<sup>3</sup> (<sup>1</sup>Banks, 1920:326; <sup>2</sup>Ross, 1937:77; <sup>3</sup>Tarter & Woodrum, 1973b:147; <sup>4</sup>Tarter, et al., 1978:231; <sup>5</sup>Whiting, 1991b:51).

**Sialis americana** Rambur, 1842:447 [*Semblis*].

Holotype female: (no locality data), Selys Collection (IRSNB).

= *Sialis ferrugineus* Walker, 1853:195. Holotype (sex unknown): USA, Georgia (BMNH).

**TAXONOMY.**—To *Sialis* (see Hagen, 1861:188); to *Protosialis* (see Weele, 1909b: 263); returned to *Sialis* (see Ross, 1937:70); *S. ferrugineus* = *S. americana* (see Banks, 1907c:22).

**DISTRIBUTION.**—**USA:** CT<sup>9</sup>, DC<sup>6</sup>, FL<sup>9</sup>, GA<sup>1</sup>, IN<sup>8</sup>, LA<sup>9</sup>, MD<sup>4</sup>, MO<sup>11</sup>, MS<sup>10</sup>, NC<sup>5</sup>, NH<sup>11</sup>, NJ<sup>11</sup>, OH<sup>3</sup>, PA<sup>2</sup>, SC<sup>9</sup>, TX<sup>9</sup>, VA<sup>6</sup>, WS<sup>7</sup> (<sup>1</sup>Walker, 1853:195 as *S. ferrugineus*; <sup>2</sup>Hagen, 1861:188; <sup>3</sup>Banks, 1907c:22; <sup>4</sup>Ross, 1937:71; <sup>5</sup>Brimley, 1938:29; <sup>6</sup>Flint, 1964:12; <sup>7</sup>Tennessee, 1968:185; <sup>8</sup>Lawson & McCafferty, 1984:129; <sup>9</sup>Tarter, et al., 1978:231; <sup>10</sup>Stark & Lago, 1983:117; <sup>11</sup>Whiting, 1991b:51).

**Sialis arvalis** Ross, 1937:68.

Holotype male: USA, California, Mokelumne Hill, IV, F. E. Blaisdell (CAS).

**DISTRIBUTION.**—**USA:** CA<sup>1</sup>, OR<sup>2</sup> (<sup>1</sup>Ross, 1937:68; <sup>2</sup>Evans, 1972:23).

**Sialis bilobata** Whiting, 1991a:411.

Holotype male: USA, California, Los Angeles Co., Brents Mountain Crags, 20.V.1939 (LACM).

**DISTRIBUTION.**—**USA:** CA (Whiting, 1991a:412).

**Sialis californica** Banks, 1920:326.

Lectotype male: USA, California, Kern Co., San Emigdio Cañon, 3.VI.1904, F. Grinnell, Jr. (MCZ).

**TAXONOMY.**—Lectotype designated (see Ross, 1937:69).

**DISTRIBUTION.**—**CAN:** AB<sup>2</sup>, BC<sup>2</sup>, **USA:** CA<sup>1</sup>, OR<sup>2</sup>, WA<sup>2</sup> (<sup>1</sup>Banks, 1920:326; <sup>2</sup>Ross, 1937:69).

**Sialis concava** Banks, 1897a:22.

Holotype male: USA, New York, Ithaca (MCZ).

**DISTRIBUTION.**—**CAN:** BC<sup>6</sup>, ON<sup>2</sup>, **USA:** MD<sup>2</sup>, ME<sup>6</sup>, NC<sup>3</sup>, NY<sup>1</sup>, VA<sup>5</sup>, WV<sup>4</sup> (<sup>1</sup>Banks, 1897a:22; <sup>2</sup>Ross, 1937:73; <sup>3</sup>Brimley, 1938:29; <sup>4</sup>Tarter et al., 1977:104; <sup>5</sup>arter et al., 1978:232; <sup>6</sup>Whiting, 1991b:51).

**Sialis contigua** Flint, 1964:10.

Holotype male: USA, Virginia, Highland Co., bridge on route 220 over East Branch Potomac River, 19.V.1963, O. S. Flint, Jr. & W. D. Field (USNM).

DISTRIBUTION.—USA: TN<sup>2</sup>, VA<sup>1</sup> (<sup>1</sup>Flint, 1964:12; <sup>2</sup>Tarter et al., 1978:232).

**Sialis cornuta** Ross, 1937:69.

Holotype male: USA, Oregon, Blue Mts., Horseshoe Lake, 7500', 26.VII.1929, H. A. Scullen (INHS).

DISTRIBUTION.—CAN: AB<sup>1</sup>, USA: ID<sup>1</sup>, MT<sup>2</sup>, OR<sup>1</sup>, UT<sup>1</sup>, WA<sup>2</sup>, WY<sup>2</sup> (<sup>1</sup>Ross, 1937:70; <sup>2</sup>Evans, 1972:32).

**Sialis dreisbachi** Flint, 1964:9.

Holotype male: USA, Michigan, Schoolcraft Co., 5.VI.1959, R. & K. Dreisbach (USNM).

DISTRIBUTION.—USA: MI<sup>1</sup>, MN<sup>2</sup>, WS<sup>3</sup> (<sup>1</sup>Flint, 1964:10; <sup>2</sup>Tarter et al., 1978:232; <sup>3</sup>Whiting, 1991b:52).

**Sialis glabella** Ross, 1937:71.

Holotype male: USA, Illinois, Mt. Carmel, 28.V.1884 (INHS).

DISTRIBUTION.—USA: IL<sup>1</sup>, Y<sup>2</sup>, MS<sup>3</sup> (<sup>1</sup>Ross, 1937:71; <sup>2</sup>Canterbury & Neff, 1980:415; <sup>3</sup>Stark & Lago, 1980:118).

**Sialis hamata** Ross, 1937:70.

Holotype male: USA, Utah, Logan, 16.IV.1933, J. A. Meacham (INHS).

DISTRIBUTION.—CAN: AB<sup>1</sup>, BC<sup>1</sup>, USA: ID<sup>2</sup>, MT<sup>1</sup>, NV<sup>2</sup>, OR<sup>1</sup>, UT<sup>1</sup>, WA<sup>2</sup>, WY<sup>1</sup> (<sup>1</sup>Ross, 1937:70; <sup>2</sup>Evans, 1972:34).

**Sialis hasta** Ross, 1937:74.

Holotype male: USA, Michigan, Lovells, along Au Sable River, 22.V.1936, Frison & Ross (INHS).

DISTRIBUTION.—AR<sup>2</sup>, IN<sup>3</sup>, KY<sup>4</sup>, MI<sup>1</sup>, MO<sup>3</sup>, PA<sup>1</sup> (<sup>1</sup>Ross, 1937:74; <sup>2</sup>Flint, 1964:12; <sup>3</sup>Tarter et al., 1978:232; <sup>4</sup>Canterbury & Neff, 1980:413).

**Sialis infumata** Newman, 1838:500.

Holotype female: USA, New Jersey, Trenton Falls, Doubleday (BMNH).

DISTRIBUTION.—CAN: ON<sup>10</sup>, USA: AR<sup>7</sup>, IL<sup>2</sup>, IN<sup>7</sup>, KS<sup>2</sup>, KY<sup>8</sup>, ME<sup>4</sup>, MI<sup>2</sup>, MN<sup>7</sup>, MO<sup>7</sup>, NC<sup>3</sup>, NJ<sup>1</sup>, NY<sup>2</sup>, OH<sup>2</sup>, OK<sup>9</sup>, PA<sup>2</sup>, SC<sup>7</sup>, VA<sup>5</sup>, WS<sup>6</sup>, WV<sup>10</sup> (<sup>1</sup>Newman, 1838:500; <sup>2</sup>Ross, 1937:73; <sup>3</sup>Brimley, 1938:29; <sup>4</sup>Procter, 1946:42; <sup>5</sup>Flint, 1964:13; <sup>6</sup>Tennessee, 1968:185; <sup>7</sup>Tarter et al., 1978:232; <sup>8</sup>Canterbury & Neff, 1980:414; <sup>9</sup>Stark & Lago, 1980:121; <sup>10</sup>Whiting, 1991b:52).

**Sialis iola** Ross, 1937:68.

Holotype male: USA, Pennsylvania, Pittsburgh (INHS).

DISTRIBUTION.—CAN: PQ<sup>1</sup>, USA: CT<sup>3</sup>, DC<sup>1</sup>, GA<sup>6</sup>, IN<sup>5</sup>, ME<sup>3</sup>, MS<sup>4</sup>, NC<sup>3</sup>, NH<sup>1</sup>, NJ<sup>1</sup>, NY<sup>1</sup>, OH<sup>3</sup>, PA<sup>1</sup>, SC<sup>3</sup>, VA<sup>2</sup> (<sup>1</sup>Ross, 1937:68; <sup>2</sup>Flint, 1964:13; <sup>3</sup>Tarter et al., 1978:232; <sup>4</sup>Stark & Lago, 1980:118; <sup>5</sup>Lawson & McCafferty, 1984:129; <sup>6</sup>Whiting, 1991b:53).

**Sialis itasca** Ross, 1937:72.

Holotype male: USA, Illinois, Momence, along Kankakee River, 1.VI.1937, B. D. Burks (INHS).

DISTRIBUTION.—CAN: ON<sup>1</sup>, PQ<sup>1</sup>; USA: AR<sup>5</sup>, DC<sup>1</sup>, GA<sup>2</sup>, IL<sup>1</sup>, IN<sup>5</sup>, KS<sup>1</sup>, MD<sup>2</sup>, MI<sup>1</sup>, MN<sup>2</sup>, MO<sup>1</sup>, NC<sup>5</sup>, ND<sup>1</sup>, NY<sup>1</sup>, OH<sup>1</sup>, OK<sup>5</sup>, PA<sup>1</sup>, TN<sup>5</sup>, TX<sup>5</sup>, VA<sup>1</sup>, WS<sup>3</sup>, WV<sup>4</sup> (<sup>1</sup>Ross, 1937:72; <sup>2</sup>Flint, 1964:13; <sup>3</sup>Tennessee, 1968:185; <sup>4</sup>Tarter, Ashley et al., 1976:32; <sup>5</sup>Tarter et al., 1978:232).

**Sialis joppa** Ross, 1937:67.

Holotype male: USA, North Carolina, Great Smoky Mountains National Park, Newfound Gap, 28.V.1934, T. H. Frison (INHS).

DISTRIBUTION.—CAN: ON<sup>6</sup>, USA: CT<sup>4</sup>, DE<sup>4</sup>, FL<sup>6</sup>, IL<sup>1</sup>, KY<sup>5</sup>, LA<sup>4</sup>, MD<sup>2</sup>, ME<sup>1</sup>, MI<sup>4</sup>, NC<sup>1</sup>, NH<sup>1</sup>, NY<sup>1</sup>, OH<sup>3</sup>, PA<sup>1</sup>, VA<sup>2</sup>, VT<sup>4</sup>, WS<sup>4</sup>, WV<sup>3</sup> (<sup>1</sup>Ross, 1937:68; <sup>2</sup>Flint, 1964:13; <sup>3</sup>Tarter & Woodrum, 1973a:165; <sup>4</sup>Tarter et al., 1978:232; <sup>5</sup>Canterbury & Neff, 1980:413; <sup>6</sup>Stange, 1990:2).

**Sialis mohri** Ross, 1937:74.

Holotype male: USA, Wisconsin, Boulder Junction, on Trout River, 20.VI.1934, Frison & Mohr (INHS).

DISTRIBUTION.—CAN: NB<sup>1</sup>, ON<sup>1</sup>, PQ<sup>1</sup>, USA: AR<sup>2</sup>, CT<sup>1</sup>, IL<sup>1</sup>, IN<sup>1</sup>, KS<sup>3</sup>, KY<sup>1</sup>, MA<sup>1</sup>, ME<sup>3</sup>, MI<sup>1</sup>, MN<sup>1</sup>, MO<sup>3</sup>, MS<sup>4</sup>, NE<sup>5</sup>, NH<sup>3</sup>, NJ<sup>1</sup>, NY<sup>1</sup>, OH<sup>1</sup>, OK<sup>2</sup>, PA<sup>1</sup>, RI<sup>3</sup>, TN<sup>3</sup>, VT<sup>5</sup>, WS<sup>1</sup> (<sup>1</sup>Ross, 1937:74; <sup>2</sup>Flint, 1964:13; <sup>3</sup>Tarter et al., 1978:232; <sup>4</sup>Stark & Lago, 1980:118; <sup>5</sup>Whiting, 1991b:53).

**Sialis nevadensis** Davis, 1903:450.

Two syntypes, male and female: USA, Nevada, Reno (USNM).

= *Sialis morrisoni* Davis, 1903:450. Holotype female: USA, Nevada, Reno (MCZ).

TAXONOMY.—*S. morrisoni* = *S. nevadensis* (see Banks, 1907c:22).

DISTRIBUTION.—CA<sup>2</sup>, NV<sup>1</sup> (<sup>1</sup>Davis, 1903:450; <sup>2</sup>Ross, 1937:76).

**Sialis nina** Townsend, 1939:224.

Holotype male: USA, Kentucky, Lexington, along North Elkhorn Creek, 1.IV.1938, P. O. Ritcher and L. H. Townsend (USNM).

DISTRIBUTION.—KY (Townsend, 1939:225).

**Sialis occidens** Ross, 1937:69.

Holotype male: USA, California, Sequoia National Park, Wolverton, 7000–9000 ft., 25.VI.1929, E. C. Van Dyke (CAS).

DISTRIBUTION.—CA<sup>1</sup>, NV<sup>2</sup> (Ross, 1937:69; Evans, 1972:37).

**Sialis rotunda** Banks, 1920:327.

Lectotype male: Canada, British Columbia, Bon Accord, 20.V (MCZ).

TAXONOMY.—Lectotype designated (see Ross, 1937:75).

DISTRIBUTION.—CAN: BC<sup>1</sup>, USA: CA<sup>3</sup> OR<sup>2</sup>, WA<sup>2</sup>, WS<sup>4</sup> (Banks, 1920:327; Ross, 1937:75; Evans, 1972:43; Tarter et al., 1978:233).

**Sialis spangleri** Flint, 1964:12.

Holotype male: USA, Maryland, Garrett Co., Swallow Falls State Park near Oakland, 14.V.1963, P. J. Spangler (USNM).

DISTRIBUTION.—MD (Flint, 1964:12).

**Sialis vagans** Ross, 1937:76.

Holotype male: USA, Indiana, Columbia City, along Eel River, 19.V.1936, Frison & Ross (INHS).

DISTRIBUTION.—CAN: NB<sup>1</sup>, NS<sup>1</sup>, ON<sup>1</sup>, PQ<sup>1</sup>; USA: AR<sup>4</sup>, CT<sup>3</sup>, FL<sup>6</sup>, GA<sup>3</sup>, IL<sup>1</sup>, IN<sup>1</sup>, KS<sup>7</sup>, KY<sup>5</sup>, MA<sup>1</sup>, MD<sup>1</sup>, ME<sup>1</sup>, MI<sup>1</sup>, MN<sup>2</sup>, MS<sup>4</sup>, NC<sup>4</sup>, NH<sup>1</sup>, NJ<sup>1</sup>, NY<sup>1</sup>, OH<sup>4</sup>, PA<sup>1</sup>, VA<sup>3</sup>, VT<sup>4</sup>, WS<sup>1</sup> (Ross, 1937:77; Parfin, 1952:422; Flint, 1964:13; Tarter et al., 1978:233; Canterbury & Neff, 1980:415; Stange, 1990:2; Whiting, 1991b:54).

**Sialis velata** Ross, 1937:71.

Holotype male: USA, Michigan, Houghton Lake, 15–18.VI.1935, T. H. Frison (INHS).

DISTRIBUTION.—CAN: AB<sup>1</sup>, BC<sup>1</sup>, MB<sup>1</sup>, ON<sup>1</sup>, PQ<sup>1</sup>, SK<sup>1</sup>; USA: CO<sup>5</sup>, CT<sup>5</sup>, ID<sup>4</sup>, IL<sup>1</sup>, IN<sup>2</sup>, KS<sup>1</sup>, MA<sup>1</sup>, MD<sup>1</sup>, ME<sup>1</sup>, MI<sup>1</sup>, MN<sup>1</sup>, MO<sup>1</sup>, MT<sup>5</sup>, NC<sup>3</sup>, ND<sup>1</sup>, NE<sup>3</sup>, NH<sup>1</sup>, NJ<sup>5</sup>, NY<sup>1</sup>, TN<sup>3</sup>, TX<sup>1</sup>, UT<sup>5</sup>, VA<sup>1</sup>, VT<sup>3</sup>, WS<sup>1</sup>, WV<sup>1</sup>, WY<sup>5</sup> (Ross, 1937:71; Lawson & McCafferty, 1984:129; Tarter et al., 1978:233; Stark & Lago, 1980:121; Whiting, 1991b:55).

## ACKNOWLEDGMENTS

The authors wish to thank John D. Oswald for pointing out that the names *Megalomus speciosus* and *Ululodes quadrimaculata* were junior primary homonyms. He also is to be thanked for very carefully reviewing an earlier version of this manuscript, spending an enormous amount of time helping the authors. The Ernst Mayr Fund at Harvard University is to be thanked for a grant to allow one of us (NDP) to study type material in the Museum of Comparative Zoology.

## LITERATURE CITED

- ADAMS, P. A. 1957a. A new genus and new species of Chrysopidae from the western United States, with remarks on the wing venation of the family. *Psyche* (Camb., Mass.) 63:67–74, 5 figs.
- . 1957b. New ant-lions from the southwestern United States (Neuroptera: Myrmeleontidae). *Psyche* (Camb., Mass.) 63:82–108, 46 figs.
- . 1958. A synonym in the genus *Chrysopa* (Neuroptera: Chrysopidae). *Psyche* (Camb., Mass.) 63(2):45.
- . 1962. A stridulatory structure in Chrysopidae (Neuroptera). *Pan-Pacific Entomologist* 38(3):178–180.
- . 1963. Taxonomy of Hawaiian *Chrysopa* (Neuroptera: Chrysopidae). *Proceedings of the Hawaiian Entomological Society* 18(2):221–223.
- . 1967. A review of the Mesochrysinae and Nothochrysinae (Neuroptera: Chrysopidae). *Bulletin of the Museum of Comparative Zoology*, Harvard University 135(4):215–238.
- . 1969. New species and synonymy in the genus *Meleoma* (Neuroptera, Chrysopidae), with a discussion of genitalic homologies. *Postilla* 136:1–18.
- . 1970. A review of the New World Dilaridae. *Postilla* 148:1–30, 14 figs.
- . 1975. Status of the genera *Ungla* and *Mallada* Navás (Neuroptera: Chrysopidae). *Psyche* (Camb., Mass.) 82:167–173.

- . 1977. Taxonomy of United States *Leucochrysa* (Neuroptera: Chrysopidae). *Psyche* (Camb., Mass.) 84(1):92–102.
- . 1982a. *Ceraeochrysa*, a new genus of Chrysopinae (Neuroptera) (Studies in New World Chrysopidae, Part II). *Neuroptera International* 2:69–75.
- . 1982b. *Plesiochrysa*, a new subgenus of *Chrysopa* (Neuroptera) (Studies in New World Chrysopidae, Part I). *Neuroptera International* 2:27–32.
- ADAMS, P. A. AND J. A. GARLAND. 1981. A new species of *Chrysopiella* Banks from western North America (Neuroptera: Chrysopidae). *Canadian Entomologist* 113:1–4, 13 figs.
- . 1982. A review of the genus *Mallada* in the United States and Canada, with a new species (Neuroptera: Chrysopidae). *Psyche* (Camb., Mass.) 89(3–4):239–248, 20 figs.
- ADAMS, P. A. AND N. D. PENNY. [1987] 1985. Neuroptera of the Amazon Basin. Part 11a. Introduction and Chrysopini. *Acta Amazonica* 15:413–479.
- AGNEW, C. W., W. L. STERLING AND D. A. DEAN. 1981. Notes on the Chrysopidae and Hemerobiidae of eastern Texas with keys for their identification. *The Southwestern Entomologist*. Supplement 4. 20 pp.
- ALAYO, P. 1968. Los Neurópteros de Cuba. *Poeyana Ser. B*(2):1–127.
- ALBARDA, H. 1891. Révision des Rhaphidides. *Tijdschrift voor Entomologie* 34:65–184.
- ASHMEAD, W. H. 1880. Orange insects; a treatise on the injurious and beneficial insects found on the orange trees of Florida. Ashmead Bros., Jacksonville, FL.
- . 1895. Notes on the cotton insects found in Mississippi. *Insect Life* 7:25–29.
- ASPÖCK, H., U. ASPÖCK AND H. RAUSCH. 1991. Die Raphidiopteren der Erde. Eine monographische Darstellung der Systematik, Taxonomie, Biologie, Ökologie und Chorologie der rezenten Raphidiopteren der Erde, mit einer zusammenfassenden Übersicht der fossilen Raphidiopteren (Insecta: Neuropteroidea). 2 vols., 730 and 550 pp. Goecke and Evers, Krefeld, Germany.
- ASPÖCK, U. 1973. Zwei neue Raphidiiden-Spezies aus Nordamerika (Neuroptera, Raphidioptera). *Polskie Pismo Entomologiczne* 43:233–240, 4 figs.
- . 1974. Die Raphidiopteren der Nearktis (Insecta, Neuropteroidea). Dissertation from The University of Wien. 2 vols., 238 pp., 777 figs, 20 maps.
- . 1975. The present state of knowledge on the Raphidioptera of America (Insecta, Neuropteroidea). *Polskie Pismo Entomologiczne* 45:537–546.
- . 1982. Polymorphismus und polytypie bei nearktischen Raphidiiden: *Agulla (Glavia) modesta* Carpenter, *Agulla (Glavia) unicolor* Carpenter und *Agulla (Glavia) paramerica* n.sp. (Neuropteroidea: Raphidioptera). *Zeitschrift der Arbeitsgemeinschaft österreichischer Entomologen* 33(3/4):95–112.
- . 1987. *Agulla faulknei* n.sp. – eine neue Raphidiiden-Spezies aus Kalifornien (Neuropteroidea: Raphidioptera: Raphidiidae). *Zeitschrift der Arbeitsgemeinschaft österreichischer Entomologen* 39(1/2):1–4.
- . 1988. *Negha meridionalis* n.sp. - eine neue Inocelliden-Spezies aus Süd-Kalifornien (U. S. A.) und Baja California (Mexiko) (Neuropteroidea: Raphidioptera: Inocelliidae). *Zeitschrift der Arbeitsgemeinschaft österreichischer Entomologen* 39(3/4):107–112.
- ASPÖCK, U. AND H. ASPÖCK. 1970. Untersuchungen über die Raphidiopteren Mexikos (Insecta, Neuropteroidea). *Polskie Pismo Entomologiczne* 40:705–725.
- BAGNALL, R. S. 1915. *Conwentzia cryptoneuris* sp.n., a neuropteron (Coniopterygidae) new to the British Fauna. *Entomologists Monthly Magazine* 51:192–193.
- BANKS, N. 1892. A synopsis, catalogue, and bibliography of the neuropteroid insects of temperate North America. *Transactions of the American Entomological Society* 19:327–373, 6 figs.
- . 1895a. Some Mexican Neuroptera. *Proceedings of the California Academy of Sciences* (2)5:515–522.

- . 1895b. New neuropteroid insects. Transactions of the American Entomological Society 22:313-316.
- . 1896. A new species of *Meleoma*. Entomological News 7:95-96.
- . 1897a. New North American neuropteroid insects. Transactions of the American Entomological Society 24:21-31.
- . 1897b. Three new species of Chrysopidae. Proceedings of the Entomological Society of Washington 4:173-175.
- . 1897c. A *Leucochrysa* from Florida. Entomological News 8:183-184.
- . 1898 [1899]. Descriptions of new North American neuropteroid Insects. Transactions of the American Entomological Society 25:199-218.
- . 1899a. New Myrmeleonidae. Entomological News 10:170-172.
- . 1899b. A Classification of the North American Myrmeleonidae. Canadian Entomologist 31:67-71.
- . 1900. A new species of *Myrmeleon* from Texas. Entomological News 11:596.
- . 1901a. A List of neuropteroid insects from Mexico. Transactions of the American Entomological Society 27:361-371, 1 pl.
- . 1901b. Some insects of the Hudsonian zone in New Mexico. VI. Neuropteroid insects. Psyche (Camb., Mass.) 9:286-287.
- . 1902. A new species of *Brachynemurus*. Entomological News 13:86.
- . 1903a. A revision of the Nearctic Chrysopidae. Transactions of the American Entomological Society 29:137-162, 1 pl.
- . 1903b. Neuropteroid insects from Arizona. Proceedings of the Entomological Society of Washington 5:237-245, 1 pl.
- . 1903c. Some new neuropteroid Insects. Journal of the New York Entomological Society 11:236-243.
- . 1903d. Notes on *Brachynemurini* of the *B. ferox* group. Proceedings of the Entomological Society of Washington 5:173-177, 1 pl.
- . 1904a. New species of *Hemerobius*. Canadian Entomologist 36:61-63.
- . 1904b. Neuropteroid insects from New Mexico. Transactions of the American Entomological Society 30:97-110, 1 pl.
- . 1904c. A list of neuropteroid insects, exclusive of Odonata, from the vicinity of Washington, D. C. Proceedings of the Entomological Society of Washington 6:201-217.
- . 1905a. Notes on neuropteroid insects from the Pacific Coast of North America, with descriptions of new species. Invertebrata Pacifica 1:85-92.
- . 1905b. Descriptions of new Nearctic neuropteroid insects. Transactions of the American Entomological Society 32:1-20, 2 pls.
- . 1905c. A revision of the Nearctic Hemerobiidae. Transactions of the American Entomological Society 32:21-51, 3 pls.
- . 1906a. Three new species of Neuroptera. Psyche (Camb., Mass.) 13:98-100, 1 fig.
- . 1906b. A Revision of the Nearctic Coniopterygidae. Proceedings of the Entomological Society of Washington 8:77-86, 2 pls.
- . 1906c. A revision of the Nearctic Coniopterygidae. Proceedings of the Entomological Society of Washington 8:77-86, 2 pls.
- . 1906d. A revision of the Nearctic Hemerobiidae. Transactions of the American Entomological Society 32:21-51, 3 pls.
- . 1907a. A new genus and a new species of Neuroptera. Entomological News 18:275.
- . 1907b. A new ant lion-fly from Arizona. Entomological News 18:100-101.
- . 1907c. Catalogue of the neuropteroid insects (except Odonata) of the United States. American Entomological Society, Philadelphia. 53 pp
- . 1908a. Neuropteroid insects—notes and descriptions. Transactions of the American Entomological Society 34:255-267, 3 pls.

- . 1908b. A new genus and species of Neuroptera. *Entomological News* 19:203–204.
- . 1908c. New tropical American Neuroptera. *Proceedings of the Entomological Society Washington* 9:30–34.
- . 1908d. On the classification of the Corydalinae, with description of a new species. *Proceedings of the Entomological Society of Washington* 10:27–30.
- . 1910. New South American neuropteroid insects. *Proceedings of the Entomological Society of Washington* 12:146–160.
- . 1911. Descriptions of new species of North American neuropteroid insects. *Transactions of the American Entomological Society*, 37:335–360, 3 pls.
- . 1912. Notes on Nearctic Mantispidae. *Proceedings of the Entomological Society of Washington* 14:178–179.
- . 1913a. New exotic neuropteroid insects. *Proceedings of the Entomological Society of Washington* 15:137–143.
- . 1913b. The genus *Brachynemurus* (Neuroptera). *Entomological News* 24:63–65.
- . 1914a. New American Chrysopidae. *Canadian Entomologist* 46:24–27.
- . 1914b. New neuropteroid insects, native and exotic. *Proceedings of the Academy of Natural Sciences of Philadelphia* 66:608–618, 1 pl.
- . 1915a. Two new names in the Ascalaphidae (Neur.). *Entomological News* 26:350.
- . 1915b. New neuropteroid insects, native and exotic. *Proceedings of the Academy of Natural Sciences of Philadelphia* 66:619–632, 1 pl.
- . 1920. New neuropteroid insects. *Bulletin of the Museum of Comparative Zoology, Harvard University* 64:299–362, 7 pls.
- . 1922. South American *Glenurus* and some other Myrmeleontidae. *Canadian Entomologist* 54:58–60.
- . 1924. Descriptions of new neuropteroid insects. *Bulletin of the Museum of Comparative Zoology, Harvard University* 65:421–455, 4 pls.
- . 1927. Revision of the Nearctic Myrmeleontidae. *Bulletin of the Museum of Comparative Zoology, Harvard University* 68(1):1–84, 4 pls.
- . 1930a. New neuropteroid insects from the United States. *Psyche (Camb., Mass.)* 37:223–233, 1 pl.
- . 1930b. Some new neotropical neuropteroid insects. *Psyche (Camb., Mass.)* 37:183–191, 6 figs.
- . 1931. A new species of *Chrysopa*. *Psyche (Camb., Mass.)* 38:174.
- . 1935. A few new North American Neuroptera. *Psyche (Camb., Mass.)* 42:53–57.
- . 1938a. New native neuropteroid insects. *Psyche (Camb., Mass.)* 45:72–79, 1 pl.
- . 1938b. New Chrysopidae and species new to the United States. *Canadian Entomologist* 70:118–122.
- . 1938c. New West Indian neuropteroid insects. *Revista de Entomología, Rio de Janeiro* 9:285–304, 29 figs.
- . 1938d. Notes on native Myrmeleontidae. *Annals of the Entomological Society of America* 31:413–420, 1 pl.
- . 1939. On some new and previously-known Neuroptera in the collection of the Academy of Natural Sciences of Philadelphia. *Notulae Naturae (Philadelphia)*, No. 32. 5 pp., 2 figs.
- . 1941a. New neuropteroid insects from the Antilles. *Memorias de la Sociedad Cubana de Historia Natural—Felipe Poey* 15:385–402, 3 pls.
- . 1941b. Three new species of Myrmeleontidae (Neuroptera). *Psyche (Camb., Mass.)* 48:101–104.
- . 1942. Contributions toward a knowledge of the insect fauna of Lower California. No. 4. Myrmeleontidae. *Proceedings of the California Academy of Sciences (4)* 24:133–152, 17 figs.
- . 1943. New Neuroptera and Trichoptera from the United States. *Psyche (Camb., Mass.)* 50:74–81, 1 fig.

- \_\_\_\_\_. 1944. Neuroptera of northern South America. Part III. Boletín de Entomología Venezolana 3:1–34.
- \_\_\_\_\_. 1946. A review of the Chrysopidae (Nothochrysidae) of Central America. Psyche (Camb., Mass.) 52(3–4):139–174.
- \_\_\_\_\_. 1948. Chrysopidae (Nothochrysidae) collected in Mexico by Dr. A. Dampf (Neuroptera). Psyche (Camb., Mass.) 55(4):151–177, 3 pls.
- \_\_\_\_\_. 1950. Notes and descriptions of western Chrysopidae (Neuroptera). Psyche (Camb., Mass.) 57:45–67, 2 pls.
- BEATTY, G. H. AND A. F. J. M. RUNNER. 1970. The occurrence of Ascalaphidae in Pennsylvania (Neuroptera). Proceedings of the Pennsylvania Academy of Sciences 44:153–155.
- BELKIN, J. N. 1954. Additional records of *Oliarces clara* in California and Arizona (Neuroptera, Ithonidae). Bulletin of the Southern California Academy of Sciences 53(2):65–68.
- BICKLEY, W. E. 1941. Records of Tennessee Chrysopidae (Neuroptera). Proceedings of the Entomological Society of Washington 43(8):187–189.
- BICKLEY, W. E. AND E. G. MACLEOD. 1956. A Synopsis of the Nearctic Chrysopidae with a key to the genera (Neuroptera). Proceedings of the Entomological Society of Washington 58(4):177–202.
- BRAM, R. A. AND W. E. BICKLEY. 1963. The green lacewings of the genus *Chrysopa* in Maryland. Maryland agricultural Experiment Station Bulletin A-124:1–18.
- BRAUER, F. 1864. Entomologische Beiträge. B. Beiträge zur Kenntniss der Neuroptera. I. Gattung *Isoscelipteron* Costa. Verhandlungen der kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien 14:898–899.
- \_\_\_\_\_. 1866. Reise der Österreichischen Fregatte *Novara* um die Erde, in ... 1857–59, unter den Befehl des Commodore B. von Wüllerstorf-Urbair, H.K. von Scherzer, ed. Zoologischer Theil, Bd. 2, No. 4 (Neuroptera). Wien. 104 pp.
- BRIMLEY, C. S. 1938. The insects of North Carolina. North Carolina Department of Agriculture, Raleigh, N.C.
- BROOKS, S. J. AND P. C. BARNARD. 1990. The green lacewings of the world: a generic review (Neuroptera: Chrysopidae). Bulletin of the British Museum of Natural History (Entomology) 59:117–286.
- BRUSHWEIN, J. R. 1987a. Bionomics of *Lomamyia hamata* (Neuroptera: Berothidae). Annals of the Entomological Society of America 80(5):671–679.
- \_\_\_\_\_. 1987b. Observations on *Lomamyia longicollis* (Neuroptera: Berothidae) in South Carolina. Journal of the Kansas Entomological Society 60(1):150–152.
- BURMEISTER, H. C. C. 1839. Handbuch der Entomologie. Bd. 2, Abt. 2 (pp. 397–1050). T. C. F. Enslin, Berlin.
- CANTERBURY, L. E. AND S. E. NEFF. 1980. Eggs of *Sialis* (Sialidae: Megaloptera) in eastern North America. Canadian Entomologist 112:409–419.
- CARPENTER, F. M. 1936. A revision of the Nearctic Raphidiodea (Recent and Fossil). Proceedings of the American Academy of Arts and Sciences 71:89–157, 2 pls.
- \_\_\_\_\_. 1940. A revision of the nearctic Hemerobiidae, Berothidae, Sisyridae, Polystoechotidae and Dilaridae (Neuroptera). Proceedings of the American Academy of Arts and Sciences 74(7):193–280.
- \_\_\_\_\_. 1992. Treatise on invertebrate paleontology. Part R. Arthropoda 4. Vols. 3 and 4: Superclass Hexapoda. The Geological Society of America and the University of Kansas. xxxi + 1–277 (vol. 3), ii + 279–655 (vol. 4). 265 figs., 2 tables.
- CAUDELL, A. N. 1933. *Neohermes infuscatus*, a new sialid from California. Pan-Pacific Entomologist 9(3):125–126.
- CHANDLER, H. P. 1953. A new species of *Climacia* from California (Sisyridae, Neuroptera). Journal of the Washington Academy of Sciences 43:182–184, 1 fig.
- \_\_\_\_\_. 1954. Four new species of dobsonflies from California. Pan-Pacific Entomologist 30(2):105–111, 10 figs.
- \_\_\_\_\_. 1956. Megaloptera. Pp. 229–233 in Aquatic Insects of California R. L. Usinger, ed., University of California Press, Berkeley.

- CLARK, W. H. 1985. First record of *Climacia californica* (Neuroptera: Sisyridae) and its host sponge, *Ephydatia mulleri* (Porifera: Spongillidae), from Idaho with water quality relationships. Great Basin Naturalist 45(3):391–394.
- CONTRERAS-RAMOS, A. 1996. A Revision of the genus *Corydalus*. Ph.D. Thesis from the University of Minnesota.
- COQUILLET, D. W. 1890. "Lace-winged fly" and "brown *Hemerobius*." Annual Report of the California State Board of Horticulture 1890:288–289, 6 figs., 1 pl.
- CURRIE, R. P. 1898. New species of North American Myrmeleonidae. I–IV. Canadian Entomologist 30:93–96, 134–140, 241–243, 273–276.
- \_\_\_\_\_. [1900] 1899. New species of North American Myrmeleonidae. Canadian Entomologist 31:361–365.
- \_\_\_\_\_. [1900] 1899. Neuroptera. In Insects of New Jersey. A list of species occurring in New Jersey, with notes on those of economic importance, J. B. Smith, ed. Trenton, New Jersey. 755 pp. [Insects of New Jersey was published as a supplement to the 27th Annual Report of the New Jersey State Board of Agriculture.]
- \_\_\_\_\_. 1903. Myrmeleonidae from Arizona. Proceedings of the Entomological Society of Washington 5:272–284.
- \_\_\_\_\_. 1904a. Notes on some Hemerobiidae from Arizona and California. Proceedings of the Entomological Society of Washington 6:79–81.
- \_\_\_\_\_. 1904b. Hemerobiidae from the Kootenay District of British Columbia. Proceedings of the Entomological Society of Washington 6:81–90.
- \_\_\_\_\_. 1906. A new Myrmeleond from the United States. Proceedings of the Entomological Society of Washington 7:186–188.
- CURTIS, J. 1834. British entomology, being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from Nature of the most rare and beautiful species, and in many instances of the plants upon which they are found. London. [Issued between 1824 and 1839 in 16 volumes containing 769 color plates with unpaginated descriptive letterpress.] *Coniopteryx psociformis* is in vol. 11: pl. 528.
- DAVIS, K. C. 1903. Sialidae of North and South America. Bulletin 68. New York State Museum of Entomology 18:442–486.
- DEGEER, C. 1773. Mémoires pour servir à l'histoire des Insectes. Vol. 3. Stockholm. 696 pp.
- DRURY, D. 1773. Illustrations of natural history wherein are exhibited upwards of two hundred and forty figures of exotic insects, ... interspersed with remarks and reflections on the nature and properties of many of them. Vol. 2. London. 90 pp.
- EISNER, T., K. HICKS, M. EISNER, AND D. S. ROBSON. 1978. "Wolf-in-sheep's clothing" strategy of a predaceous insect larva. Science, New York 199:790–794, 4 figs.
- ENDERLEIN, G. 1905a. *Conwentzia pineticola* nov. gen., nov. spec., eine neue Neuroptera aus Westpreussen. Bericht des Westpreussischen Botanischen-Zoologischen Vereins 26–27 (Anlagen):10–12.
- \_\_\_\_\_. 1905b. Klassifikation der Neuropteren-Familie Coniopterygidae. Zoologischer Anzeiger 29:225–227.
- \_\_\_\_\_. 1905c. Ein neuer zu den Coniopterygiden gehöriger Neuropteren-Typus aus der Umgebung von Berlin. Wiener entomologische Zeitung 24:197–198.
- \_\_\_\_\_. 1906. Monographie der Coniopterygiden. Zoologische Jahrbücher (Abt. für Systematik, Geographie und Biologie der Tiere) 23:173–242.
- \_\_\_\_\_. 1907. Nachträge zur Monographie der Coniopterygiden. Stettiner entomologische Zeitung 68:10–13.
- \_\_\_\_\_. 1908. Neuroptera: Fam. Coniopterygidae. In Wytsman, P., Genera Insectorum, fasc. 67. 18 pp. Bruxelles. 2 pls., 25 figs.
- \_\_\_\_\_. 1910. Klassifikation der Mantispiden nach dem Material des Stettiner Zoologischen Museums. Stettiner Entomologische Zeitung 71:341–379.
- \_\_\_\_\_. 1930. Die Klassifikation der Coniopterygiden auf Grund der recenten und fossilen Gattungen. Archiv für klassifikatorische und phylogenetische Entomologie 1:98–114.

- ESBEN-PETERSEN, P. 1912. A few new Neuroptera Planipennia. *Entomologische Mitteilungen* 1:267–273, 7 figs.
- . 1928. Neue und wenig bekannte Neuropteren des Hamburger Museums. *Deutsche entomologische Zeitschrift* 1928:73–77.
- . 1933. New and little-known Neuroptera. *Videnskalbelige Meddelelser fra Dansk Naturhistorisk Forening, Kjøbenhavn* 94:109–123, 13 figs.
- . 1934. The Norwegian zoological expedition to the Galapagos Islands, 1925, conducted by Alf Wollebaek. XII. Neuropterous insects from Galapagos Is. *Nytt Magasin for Naturvidenskapen* 74:291–294, 3 figs.
- EVANS, E. D. 1972. A study of the Megaloptera of the Pacific coastal region of the United States. Unpublished Ph.D. thesis from Oregon State University, June 1972. 210 pp.
- . 1984. A new genus and a new species of dobsonfly from the far western United States (Megaloptera: Corydalidae). *Pan-Pacific Entomologist* 60:1–3, 3 figs.
- FABRICIUS, J. C. 1781. *Species insectorum exhibentes eorum differentias specificas, synonyma avctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus. Hamburgi et Kilonii. Tome I.* 552 pp.
- . 1793. *Entomologia systematica emendata et aucta secundum classes, ordines, genera, species adiectis synonymis, locis observationibus, descriptionibus. Hafniae. Vol. 2.* 519 pp.
- FAULKNER, D. K. 1990a. Phantom of the Desert. Biology of the little-known moth lacewing. *Environment West* 1(2):17–19.
- . 1990b. Current knowledge of the biology of the moth-lacewing *Oliarces clara* Banks (Insecta: Neuroptera: Ithonidae). Advances in Neuropterology. Proceedings of the Third International Symposium of Neuropterology, pp. 197–203.
- FITCH, A. [1855] 1854. Report [upon the noxious and other insects of the state of New York]. *Transactions of the New York State Agricultural Society* 14:705–880.
- FLESCHNER, C. A. AND D. W. RICKER. 1953. Food habits of coniopterygids on citrus in southern California. *Journal of Economic Entomology* 46(3):458–461.
- FLINT, O. S., JR. 1964. New species and new state records of *Sialis* (Neuroptera: Sialidae). *Entomological News* 75:9–13, 11 figs.
- . 1965. The genus *Neohermes*. *Psyche* (Camb., Mass.) 72:255–263, 19 figs.
- . 1974. A coniopterygid (*Aleuropteryx juniperi*)—Virginia. *Cooperative Economic Insect Report* 24:703.
- FROESCHNER, R. C. 1947. Notes and keys to the Neuroptera of Missouri. *Annals of the Entomological Society of America* 40:123–136.
- GARLAND, J. A. 1985. Identification of Chrysopidae in Canada, with bionomic notes (Neuroptera). *Canadian Entomologist* 117:737–762.
- GARLAND, J. A. AND B. D. MARSHALL. 1980. Confirmation of *Uluodes quadrimaculata* in Canada (Neuroptera: Ascalaphidae). *Canadian Entomologist* 112:637–638.
- GERSTAECKER, A. 1894. Über neue und weniger bekannte Neuropteren aus der Familie Megaloptera, Burm. *Mitteilungen des Naturwissenschaftlichen Vereins für Neu-Vorpommern und Rügen im Greifswald* 25:93–173.
- GÖSZY, G. VON. 1852. Beobachtungen der Verwandlungsgeschichte und Beschreibung einiger neuen Arten der Gattung *Hemerobius* Leach. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Class., Wien* 8:344–348.
- GULDING, L. 1825. The natural history of *Phasma cornutum*, and the description of a new species of *Ascalaphus*. *Transactions of the Linnean Society of London* 14:137–141.
- GURNEY, A. B. 1948. Notes on Nearctic Hemerobiidae, with descriptions of two new species (Neuroptera). *Annals of the Entomological Society of America* 41:213–222, 23 figs.
- HAGEN, H. A. 1860. Beitrag zur Kenntniss der *Myrmeleon*-arten. *Stettiner Entomologische Zeitung* 21:359–369.

- . 1861. Synopsis of the Neuroptera of North America. With a list of the South American species. Smithsonian Miscellaneous Collections 4:1–347.
- . 1863. Observations on certain N.A. Neuroptera. Proceedings of the Entomological Society of Philadelphia 2:167–182.
- . 1866. Hemerobidarum synopsis synonymica. Stettiner Entomologische Zeitung 27:369–462.
- . 1873. Odonata from the Yellowstone. Annual Report. F. V. Hayden, ed. United States Geological Survey of the Territories 6:727–729.
- . 1875. Report upon the collections of Neuroptera and Pseudo-Neuroptera made in portions of Colorado, New Mexico and Arizona, during the years 1872, 1873 and 1874. Pp. 909–922 in Report upon Geographical and Geological Explorations and Surveys West of the 100th Meridian. Vol. 5 (Zoology). Chapter XIV.
- . 1877. *Symphrasis*, eine neue Mantispiden-Gattung. Stettiner entomologische Zeitung 38:208–211.
- . 1886. Monograph of the Hemerobiidae. Part II. Proceedings of the Boston Society of Natural History 23:276–292.
- . 1887. Stray notes on Myrmeleonidae. Canadian Entomologist 19:89–93, 110–112, 133–136, 147–156, 209–216.
- . 1888. Stray notes on Myrmeleonidae. Canadian Entomologist 20:34–38, 57–60, 72–74, 93–97, 185–191, 204–211.
- HALE CARPENTER, G. O. 1938. Notes on insects collected in West Greenland by the Oxford University Greenland Expedition, 1936. With descriptions of a new species of *Angitia* (Hymenoptera, Ichneumonidae) by A. Roman and of *Fannia* (Diptera, Anthomyiidae) by J. E. Collin. Annals and Magazine of Natural History (11)1:529–553.
- HANDSCHIN, E. 1960. Zur Revision süd-amerikanischer Mantispiden. Revue Suisse de Zoologie 67:523–560.
- HENRY, C. S. 1977. The behavior and life histories of two North American ascalaphids. Annals of the Entomological Society of America, 70:179–195, 10 figs.
- . 1983. Acoustic recognition of sibling species within the holarctic lacewing *Chrysoperla carnea* (Neuroptera: Chrysopidae). Systematic Entomology 8:293–301.
- . 1985. Sibling species, call differences, and speciation in green lacewings (Neuroptera: Chrysopidae: *Chrysoperla*). Evolution 39(5):965–984.
- . 1993. *Chrysoperla johnsoni* (Neuroptera: Chrysopidae): Acoustic evidence for full species status. Annals of the Entomological Society of America 86(1):14–25.
- HENRY, C. S., M. M. WELLS, AND R. J. PUPEDIS. 1993. Hidden taxonomic diversity within *Chrysoperla plorabunda* (Neuroptera: Chrysopidae): Two new species based on courtship songs. Annals of the Entomological Society of America 86(1):1–13.
- HENRY, T. J. 1974. Beneficial insects. A coniopterygid (*Aleuropteryx juniperi* Ohm). Cooperative Economic Insect Report (Plant Protection and Quarantine Programs) 24:659.
- HOFFMAN, K. M. 1989. Taxonomic status of *Mantispa sayi*, *Mantispa fuscicornis* and *Mantispa uhleri* (Neuroptera: Mantispidae). Proceedings of the Entomological Society of Washington 91:637–639.
- . 1990. Distributional notes on North and Central American Dilaridae (Neuroptera). Entomological News 101(3):155–157.
- HOFFMAN, K. M. AND J. R. BRUSHWEIN. 1989. Species of spiders (Araneae) associated with the immature stages of *Mantispa pulchella* (Neuroptera, Mantispidae). Journal of Arachnology 17:7–14.
- . 1992. Descriptions of the larvae and pupae of some North American Mantispinae (Neuroptera: Mantispidae) and development of a system of larval chaetotaxy for Neuroptera. Transactions of the American Entomological Society 118(2):159–196.
- ILLIGER, J. K. W. 1798. [Genus *Mantispa*]. P. 499 in Verzeichnis der Käfer Preussens, entworfen von Johann Gottlieb Kugelann ... ausgearbeitet von Johann Karl Wilhelm Illiger. mit einer Vorrede des Professors und Pagenhofmeisters Helwig in Braunschweig, und dem angehängten Versuche einer natürlichen Ordnungs- und Gattungs-Folge der Insekten. Halle. 510 pp.

- JOHNSON, J. B. AND K. S. HAGEN. 1981. A neuropterous larva uses an allomone to attack termites. *Nature* 289:506-507.
- JOHNSON, V. 1976. A new genus and species of Coniopterygidae (Neuroptera) from New Mexico. *Psyche* (Camb., Mass.), 83(2):192-195, 9 figs.
- \_\_\_\_\_. 1981a. Review of the Coniopterygidae (Neuroptera) of North America with a revision of the genus *Aleuropteryx*. *Psyche* (Camb., Mass.) 87(3-4):259-298, 15 figs.
- \_\_\_\_\_. 1981b. New species of Coniopterygidae (Neuroptera) from North America. *Psyche* (Camb., Mass.) 87(3-4):181-192, 7 figs.
- KASTON, B. J. 1938. Mantispidae parasitic on spider egg sacs. *Journal of the New York Entomological Society* 46:147-153.
- KEVAN, D. K. MCE. 1989. Ms. Taylor's Walking-sticks: "Queen Bess" and odd "orthopterians" of 1862 Mantodea, Phasmatodea, Dermaptera, a reduviid Hemipteran and mantispid Neuroptera). *Proceedings of the Entomological Society of Ontario* 120:1-9.
- KEVAN, D. K. MCE. AND J. KLIMASZEWSKI. 1986. Notes on the *Hemerobiidae* or brown lacewing flies (Neuroptera) of Canada and Alaska. *Neuroptera International* 4(1):7-22.
- \_\_\_\_\_. 1987. The *Hemerobiidae* of Canada and Alaska. Genus *Hemerobius* L. *Giornale Italiano di Entomologia* 16:305-369. 94 figs., 11 maps.
- KILLINGTON, F. J. 1937. A monograph of the British Neuroptera. Vol. 2. Ray Society, London. 306 pp.
- KIMMINS, D. E. 1928. New and little known Neuroptera of Central America. EOS: *Revista española de Entomología* 4:363-370, 7 figs.
- \_\_\_\_\_. 1932. *Hemerobius simulans* Walk., *H. orotypus* Wallingr., and *H. crispus* Steph. Walk. (Neuroptera). *Entomologist* 65:87-88.
- \_\_\_\_\_. 1935. Notes on the genera *Megalomus* Rambur and *Nesobiella*, gen. nov. (Neuroptera), with descriptions of new species. *Annals and Magazine of Natural History* (10)16:602-619, 11 figs.
- \_\_\_\_\_. 1940. Notes on some types of Chrysopidae (Neuroptera) in the British Museum collections. *Annals and Magazine of Natural History* 11(5):442-449.
- KLAPALEK, F. 1894. Is *Aleuropteryx lutea*, Löw, identical with *Coniopteryx lutea* Wallg.? *Entomologist's Monthly Magazine* 5:121-122.
- KLIMASZEWSKI, J. AND D. K. MCE. KEVAN. 1985. The brown lacewing flies of Canada and Alaska (Neuroptera: Hemerobiidae). Part I The Genus *Hemerobius* Linnaeus: Systematics, Bionomics and Distribution. Lyman Entomological Museum and Research Laboratory Memoir No. 15. 119 pp.
- \_\_\_\_\_. 1987a. The brown lacewing flies of Canada and Alaska (Neuroptera; Hemerobiidae). Part IIb. The genus *Wesmaelius* Krüger. *Neuroptera International* 4:245-274.
- \_\_\_\_\_. 1987b. The brown lacewing flies of Canada and Alaska (Neuroptera; Hemerobiidae). Part II[a]. The genus *Wesmaelius* Krüger. *Neuroptera International* 4:153-204. 167 figs., 8 maps.
- \_\_\_\_\_. 1988a. The brown lacewing flies of Canada and Alaska (Neuroptera: Hemerobiidae). Part III. The genus *Micromus* Rambur. *Giornale italiano di Entomologia* 19:31-76, 42 figs., 4 maps.
- \_\_\_\_\_. 1988b. New distribution data for Canadian Hemerobiidae (Neuroptera): Genera: *Hemerobius* Linnaeus, *Micromus* Rambur and *Wesmaelius* Krüger. Supplement [1st]. *Revue d'Entomologie du Québec* 32:9-15.
- \_\_\_\_\_. 1989. Distribution data for some Nearctic species of the genus *Hemerobius* Linnaeus and *Wesmaelius* Krüger (Neuroptera: Hemerobiidae.). Supplement 2. *Neuroptera International* 5(4):205-210.
- \_\_\_\_\_. 1990a. New distribution data for some American species of the genus *Hemerobius* Linneaus [sic], with a redescription of *Hemerobius alpestris* Banks (Neuroptera: Hemerobiidae). *Annals of the Transvaal Museum* 35:151-155, 6 figs.
- \_\_\_\_\_. 1990b. The genus *Micromus* Rambur in North America, with special reference to Canada and Alaska—a synopsis (Insecta: Neuroptera: Hemerobiidae). Pp. 63-66 in *Advances in Neuropterology*. M. W. Mansell and H. Aspöck, eds. Proceedings of the Third International Symposium on Neuropterology, South African Department of Agricultural Development: Pretoria.
- KLUG, J. C. F. [1838] 1836. Versuch einer systematischen Feststellung der Insekten-Familie: Panorpatae

- und Auseinandersetzung ihrer Gattungen und Arten. Abhandlungen der königlichen Akademie der Wissenschaften zu Berlin. 1836:81–108.
- KOLBE, J. 1888. Die geographische Verbreitung der Neuroptera und Pseudoneuroptera der Antillen, nebst einer Übersicht über die von Herrn Consul Krug auf Portoriko gesammelten Arten. Archiv für Naturgeschichte 54:153–178.
- KRÜGER, L. 1922. Hemerobiidae. Beiträge zu einer Monographie der Neuropteren-Familie der Hemerobiiden. Stettiner entomologische Zeitung 83:138–172.
- KUWAYAMA, S. 1956. Further studies on the Neuroptera-Planipennia of the Kurile Islands. Insecta Matsumurana, 20:77–82, 2 figs.
- LACROIX, J. L. 1926. Quelques insectes Névroptères exotiques. Annales de la Société Linnéenne de Lyon 72:67–72.
- LAGO, P. K. 1981. Mississippi spongillaflies [Sisyridae, Neuroptera]: A new state record, additional distribution records and A key to the adults. Journal of the Mississippi Academy of Sciences 26:27–30.
- LAGO, P. K. AND S. TESTA, III. 1989. Records of owlflies (Neuroptera: Ascalaphidae) from Mississippi, with a key to species. Entomological News 100(1):11–17.
- LAMBKIN, K. J. 1986. A revision of the Australian Mantispidae (Insecta: Neuroptera) with a contribution to the classification of the family. I. General and Drepanicinae. Australian Journal of Zoology, Supplementary Series No. 116:1–142, 391 figs.
- LATREILLE, P. A. 1796. Classe Quatrième. Névroptères. Pp. 96–104 in Précis des caractères généraux des insectes disposés dans un ordre naturel. Paris. 201 pp.
- . 1802. Histoire naturelle, générale et particulière de Crustacés et des Insectes. Vol. 3, Dufart, Paris. 467 pp.
- LAWSON, H. R. AND W. P. McCAFFERTY. 1984. A checklist of Megaloptera and Neuroptera (Planipennia) of Indiana. The Great Lakes Entomologist 17(3):129–131.
- LEACH, W. E. 1815. Entomology. Pp. 57–172 in Edinburgh Encyclopaedia, D. Brewster, ed. Vol. 9, part 1, Edinburgh.
- LEONARD, M. D. 1928. A list of the insects of New York. Cornell University Agricultural Experiment Station. Memoir 101.
- LICHTENSTEIN, A. A. H. 1796. Catalogus rerum naturalium rarissimarum Hamburgi - auctionis lege distrahendarum. Gottl. Friedr. Schniebes, Hamburg. Vol. 3. xii + 222 pp.
- LINNAEUS, C. 1758. Systema natura per regna tria naturae secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. 10th Edition. Vol. 1. Salvii, Holmiae, Sweden. 824 pp.
- . 1761. Fauna Svecica, sistens animalia Sveciae regni: Mammalia, Aves, Amphibia, Pices, Insecta, Vermes distributa per classes and ordines, genera and species, cum differentiis specierum, synonymis auctorum, nominibus natalium, descriptionibus insectorum. Salvii, Stockholmiae, Sweden. 578 pp.
- . 1763. Centuria Insectorum, quam, praeside D. D. Car. von Linné, proposuit Boas Johansson, Calmariensis. Pp. 384–415 in Amoenitates Academicæ; seu dissertationes variae physicae, medicæ, botanicae; antehac seorsim editæ, nunc collectæ et auctæ cum tabulis aeneis. Vol. 6 (No. 121). Salvii, Holmiae, Sweden.
- . 1767. Systema natura per regna tria naturae secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. 12th Ed. Vol. 1, pt. 2 (pp. 533–1327). Salvii, Holmiae, Sweden.
- LOW, F. 1885. Beitrag zur Kenntnis der Coniopterygiden. Sitzungsberichte der Akademie der Wissenschaften, Wien. Mathematisch- Naturwissenschaftliche Klasse (Abt. I) 91:73–89.
- LUCAS, J. R. AND L. A. STANGE. 1981. Key and descriptions to the *Myrmeleon* larvae of Florida (Neuroptera: Myrmeleontidae). Florida Entomologist 64(2):207–216, figs. 1–6.
- MACGILLIVRAY, A. D. 1894. New species of *Nothochrysa*. Canadian Entomologist 26:169–171.
- MACLEOD, E. G. AND L. A. STANGE. 1981. The brown lacewings of Florida. Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Entomology Circular 227: 4 unnumbered pages, 17 figs.

- MACNEILL, C. D. 1962. Observations on the flight behavior of an ascalaphid of the genus *Ululodes* (Neuroptera: Ascalaphidae). Pan-Pacific Entomologist 38:186-188.
- MADDUX, D. E. 1954. A new species of dobsonfly from California (Megaloptera: Corydalidae). Pan-Pacific Entomologist 30:70-71.
- MAKARKIN, V. N. 1985. A Review of the Hemerobiidae (Neuroptera) of the USSR. I. The genera *Hemerobius* L., *Micromus* Ramb., and *Paramicromus* Nakah. [in Russian]. Entomologicheskoe Obozreniye 64:158-170. 49 figs. (English translation: 1985: Entomological Review 64(2):99-113.)
- MCCLENDON, J. F. 1901. A new species of *Chrysopa* from Texas. Psyche (Camb., Mass.) 9:215-216, 1 fig.
- . 1906a. Notes on the true Neuroptera. I. *Brachynemurus curriei*, a new ant-lion from Texas. Entomological News 17:93.
- . 1906b. Notes on the true Neuroptera. 3. A catalogue of Texas Neuroptera. Entomological News 17:169-173.
- McLACHLAN, R. [1867] 1868. New genera and species, &c., of neuropterous insects; and a revision of Mr. F. Walker's British Museum Catalogue of Neuroptera, part ii (1853), as far as the end of the genus *Myrmeleon*. Journal of the Linnean Society of London, Zoology 9:230-281.
- . 1868. A monograph of the British Neuroptera—Planipennia. Transactions of the Entomological Society of London 1868:145-224.
- . 1869. New species, &c., of Hemerobiina; with synonymic notes (first series). Entomologist's Monthly Magazine 6:21-27, 1 fig.
- . 1873a [1871]. An attempt towards a systematic classification of the family Ascalaphidae. Journal of the Linnean Society of London, Zoology 11:219-276.
- . 1873b. Notes sur les Myrméléonides décrits par M. le Dr. Rambur. Annales de la Société de Entomologie de Belgique 16:127-141.
- . 1881. A North American species of *Dilar*. Entomologist's Monthly Magazine 18:55.
- . 1891. Descriptions of new species of holophtalmous Ascalaphidae. Transactions of the Entomological Society of London 1891:509-515.
- MEINANDER, M. 1972. A revision of the family Coniopterygidae (Planipennia). Acta Zoologica Fennica 136:1-357, 223 figs.
- . 1974. Coniopterygidae from western North America (Neuroptera). Entomologica Scandinavica 5(3-4):217-232, 13 figs.
- . 1975. Coniopterygidae from North America (Neuroptera). Notulae Entomologicae, 55(1):28-32 4 figs.
- . 1990. The Coniopterygidae (Neuroptera, Planipennia). A check-list of the species of the world, descriptions of new species and other new data. Acta Zoologica Fennica 189:1-95, 29 figs.
- MERRITT, R. W. AND K. W. CUMMINS. 1978. An introduction to the aquatic insects of North America. Kendall/Hunt Publishing Company, Dubuque, Iowa. 441 pp.
- MONTGOMERY, B. E. AND A. W. TRIPPEL. 1933. A preliminary list of Indiana Neuroptera. Entomological News 44:258-261.
- MONSERRAT, V. J. 1993. New data on some species of the genus *Micromus* Rambur, 1842. Annali del Museo Civico di Storia Naturale "G. Doria" 89:477-516.
- . 1996. Revisión del Género *Hemerobius* de Latinoamerica (Neuroptera, Hemerobiidae). Fragmenta Entomologica, Roma, 27(2):399-523.
- MOTSCHULSKY, V. VON. 1853. Neuroptera, Pp. 211-212 in Etudes Entomologiques. Helsingfors.
- MUNROE, E. G. 1953. *Chauliodes disjunctus* Walker: a correction, with the descriptions of a new species and a new genus (Megaloptera: Corydalidae). Canadian Entomologist 85:190-192, 4 figs.
- NAKAHARA, W. 1960. Systematic studies on the Hemerobiidae (Neuroptera). Mushi 34:1-69, 16 pls.
- . 1966. Contributions to the knowledge of the Hemerobiidae of western North America (Neuroptera). Proceedings of the United States National Museum 116:205-222, 1 pl., 4 figs.

- NAVÁS, L. 1905. Notas zoológicas. VIII. Mis excursiones durante el verano de 1904. Boletín de la Sociedad Aragonesa de Ciencias Naturales, Zaragoza 4:107–131.
- . 1908. Neurópteros nuevos. Memorias de la real Academia de Ciencias y Artes de Barcelona (3)6:401–423, 29 figs.
- . 1909a. Monografía de la familia de los Diláridos (Ins. Neur.). Memorias de la real Academia de Ciencias y Artes de Barcelona (3)7:619–671.
- . 1909b. Sur deux Hémérobides (insectes névroptères) nouveaux. Annales de la Société Scientifique de Bruxelles 33:215–220.
- . 1911. Chrysopides nouveaux. Annales de la Société Scientifique de Bruxelles 35:266–282.
- . 1912a. Neurópteros nuevos de América. [1A]. Brotéria (Serie zoológica) 10:194–202.
- . 1912b. Crisópidos y Hemeróbidos (Ins. Neur.) nuevos ó críticos. Brotéria (Serie zoológica) 10:98–113.
- . 1913a. Les Chrysopides (Ins., Névr.) du Musée de Londres. [1A]. Annales de la Société scientifique de Bruxelles 37:292–330. [For the conclusion of this article, see Navás, 1914e.]
- . 1913b. Névroptères nouveaux de l'Amérique du Nord. [1]. Entomologische Zeitschrift, Frankfurt am Main 27:19–20.
- . 1913c. Crisópidos Sudamericanos. Brotéria (Serie zoológica) 11:73–104, 149–168.
- . 1913d. [Notas entomológicas, 2.a serie]. 5. Cuatro pequeñas colecciones de Neurópteros de la península Ibérica. Boletín de la Sociedad Aragonesa de Ciencias Naturales, Zaragoza 12:77–89, 2 figs.
- . 1913e. Neurópteros nuevos de América. [1b]. Brotéria (Serie zoológica) 11:45–53, 4 figs.
- . 1914a. Neurópteros sudamericanos. Primera serie. Brotéria (Serie zoológica) 12:45–56, 215–234.
- . 1914b. New Neuroptera from the United States. Bulletin of the Brooklyn Entomological Society 9:13–20, 5 figs.
- . 1914c. Some Neuroptera from the United States. Bulletin of the Brooklyn Entomological Society 9:60–62.
- . 1914d. Algunos Neurópteros del Museo de Oxford. I Serie. Boletín de la Sociedad Aragonesa de Ciencias Naturales, Zaragoza 13:61–68.
- . 1914e. Les Chrysopides (Ins., Névr.) du Musée de Londres. [1B]. Annales de la Société scientifique de Bruxelles 38:73–114. [For the beginning of this article, see Navás, 1913a.]
- . 1914f. Mantíspidos nuevos. (Segunda [II] serie). Memorias de la real Academia de Ciencias y Artes de Barcelona (3)11:83–103, 16 figs.
- . 1914g. Névroptères nouveaux de l'Amérique du Nord. IIème série. Entomologische Zeitschrift, Frankfurt am Main 28:18–20, 25–26.
- . 1915a. Some Neuroptera from the United States. Bulletin of the Brooklyn Entomological Society 10:50–54, 2 figs.
- . 1915b. Neurópteros nuevos ó poco conocidos. (Quinta [V] serie). Memorias de la real Academia de Ciencias y Artes de Barcelona (3)11:455–480.
- . 1915c. Neurópteros nuevos ó poco conocidos. (Sexta [VI] serie). Memorias de la real Academia de Ciencias y Artes de Barcelona (3)12:119–136. [Separate: (3)12(7):1–20.]
- . 1916a. Neurópteros sudamericanos. Tercera [III] serie. Neurópteros del Brasil recogidos por el R.P. Joaquín da Silva Tavares S.J. Brotéria (Serie zoológica) 14:14–35.
- . 1916b. Notas sobre el orden de los Rafidiópteros (Ins.). Memorias de la real Academia de Ciencias y Artes de Barcelona (3)12:507–513, 3 figs.
- . 1917a. Insecta nova. 1 and 2 series. Memorie dell' Accademia Pontifica dei Nuovi Lincei, Roma 3:1–22.
- . 1917b. Neue Neuropteren. Dritte Serie. Entomologische Mitteilungen (Berlin) 6(7/9): 274–282, 5 figs.
- . 1918a. Neurópteros nuevos ó poco conocidos. (Décima serie). Memorias de la real Academia de Ciencias y Artes de Barcelona (3)14:339–366, 20 figs.

- \_\_\_\_\_. 1918b. Insecta nova. IV Serie. Memorie dell' Accademia Pontificia dei Nuovi Lincei, Roma (2)4:13–23, 5 figs.
- \_\_\_\_\_. 1919a. Monografia de l'ordre dels Rafidiòpters (Ins). Arxius [Arxius] de l'Institute [d'Estudis Catalans, Seccio] de Ciencias, Barcelona. 90 pp., 42 figs.
- \_\_\_\_\_. 1919b. Algunos insectos Neurópteros de la República Argentina. Serie Tercera. Revista de la real Academia de Ciencias Exactas, Físicas y Naturales de Madrid 17:287–305, 6 figs.
- \_\_\_\_\_. 1920. Sur des Névroptères nouveaux ou critiques II. Annales de la Société Scientifique de Bruxelles 39:189–203, 5 figs.
- \_\_\_\_\_. 1921. Insectos americanos nuevos ó críticos. Brotéria (Serie zoológica) 19:113–124, 5 figs.
- \_\_\_\_\_. 1922a. Insectos exóticos. Brotéria (Serie zoológica) 20:49–63, 7 figs.
- \_\_\_\_\_. 1922b. Sur des Névroptères nouveaux ou critiques. Sixième [VI] série. Annales de la Société Scientifique de Bruxelles 42:166–173.
- \_\_\_\_\_. 1922c. Insectos nuevos ó poco conocidos. [I]. Memorias de la real Academia de Ciencias y Artes de Barcelona 17:383–400.
- \_\_\_\_\_. 1923. Estudis sobre Neuròpters (Insectes). Arxius [Arxius] de l'Institute [d'Estudis Catalans, Seccio] de Ciencias, Barcelona 7:179–203, 2 figs.
- \_\_\_\_\_. 1924a. Algunos insectos de Cuba, recogidos por don Fermín Z. Cervera. [I]. Revista de la real Academia de Ciencias Exactas Físicas y Naturales de Madrid (2)21:323–332.
- \_\_\_\_\_. 1924b. Algunos insectos de Cuba. Segunda [II] Serie. Revista de la real Academia de Ciencias Exactas, Físicas y Naturales de Madrid (2)21:333–340.
- \_\_\_\_\_. 1924c. Crisópidos (Ins. Neur.) neotrópicos. [I]. Revista Chilena de Historia Natural 27:110–116.
- \_\_\_\_\_. 1924d. Insectos de la Argentina y Chile. Estudios: Revista Mensual (Academia literaria del Plata, Buenos Aires) 22:358–368, 4 figs.
- \_\_\_\_\_. 1925a. Neuropteren aus Brasilien. Mitteilungen der Münchener entomologischen Gesellschaft 15:64–68, 2 figs.
- \_\_\_\_\_. [1925b] 1924. Comunicaciones Entomológicas. 7. Neurópteros del Museo de Berlín. Revista de la real Academia de Ciencias Exactas, Físico-Químicas y Naturales de Zaragoza (1)9:20–34, 8 figs.
- \_\_\_\_\_. 1925c. Insectos exóticos nuevos ó poco conocidos. Segunda [II] serie. Memorias de la real Academia de Ciencias y Artes de Barcelona (3)19:181–200, 11 figs.
- \_\_\_\_\_. 1925d. Crisópidos (Ins. Neur.) neotrópicos. Segunda [II] serie. Revista chilena de Historia natural 29:8–13, 2 figs.
- \_\_\_\_\_. 1925e. Entomologische Ergebnisse der schwedischen Kamtchatka-Expedition 1920–1922. 6. Neuroptera et Mecoptera. Arkiv för Zoologi 18B(2):1–4, 2 figs.
- \_\_\_\_\_. 1927a. Veinticinco formas nuevos de insectos. Boletín de la Sociedad ibérica de Ciencias naturales 26:48–75, 9 figs.
- \_\_\_\_\_. 1927b. Insecta Nova. Series XII. Memorie dell' Accademia Pontifica dei Nuovi Lincei, Roma (2)10:1–10, 4 figs.
- \_\_\_\_\_. 1928a. Insectos del Museo de Estocolmo. Revista de la real Academia de Ciencias Exactas, Físicas y Naturales de Madrid (2)24:28–39, 6 figs.
- \_\_\_\_\_. 1928b. Insectos neotrópicos. 3a serie. Revista chilena de Historia natural 31:316–328, 7 figs.
- \_\_\_\_\_. 1929a. Monografía de la Familia de los Berótidos (Insectos Neurópteros). Memorias de la Academia de Ciencias Exactas, Físico-Químicas y Naturales de Zaragoza 2:1–107, 44 figs.
- \_\_\_\_\_. 1929b. Insectos Neurópteros del Museo de Hamburgo. Memorias de la Real Sociedad Española de Historia Natural 15:315–322, 3 figs.
- \_\_\_\_\_. 1929c. Insecta Nova. Serie XIV. Memorie dell' Accademia pontifica dei Nuovi Lincei, Roma (2)12:24–32, 3 figs.
- \_\_\_\_\_. 1931. Decadas de Insectos Nuevos. Decada 3. Revista de la real Academia de Ciencias Exactas, Físicas y naturales de Madrid 26:79–86.

- . 1932. Alcuni insetti del museo di zoologia della real Università di Torino. Bollettino dei Musei di Zoologia e di Anatomia Comparata della R. Università di Torino (3)42(26):1–38, 32 figs.
- . 1933a. Insectos Suramericanos. Septima [VII] serie. Revista de la real Academia de Ciencias Exactas, Físicas y Naturales de Madrid 30:303–314, 7 figs.
- . 1933b. Neurotteri e Tricotteri del “Deutches Entomologisches Institut” di Berlino-Dahlem. Bollettino della Società Entomologica Italiana 65:105–113, 8 figs.
- . 1933c. Decadas de insectos nuevos. Década 24. Brotéria (Ciencias naturales) 2:101–110, 7 figs.
- . 1934. Insectos de Méjico. Boletín de la Sociedad Entomológica de España 16:68–75, 2 figs.
- . 1935. Monografía de la familia de los sisíridos (Insectos neurópteros). Memorias de la Academia de Ciencias Exactas, Físico-Químicas y Naturales de Zaragoza 4:1–87, 40 figs.
- NEEDHAM, J. G. 1901. Order Neuroptera, Pp. 540–560 *In* Needham and Betten, Aquatic Insects in the Adirondacks. New York State Museum Bulletin 47 (Entomology 13). 6 text figs., 1 pl.
- . 1905. Systematic notes on Hemerobiidae. Bulletin of the New York Museum 86 (Entomology 23):15–17, 2 pls.
- NEWMAN, E. 1838. Entomological Notes. Entomological Magazine 5:168–181, 372–402, 483–500.
- OHM, P. 1968. Vorläufige Beschreibung einer neuen europäischen *Aleuropteryx*-Art (Neuroptera, Coniopterygidae). Entomologisches Nachrichtenblatt, Wien 15:12–15, 4 figs.
- . 1973. Zwei neue Coniopterygiden-Arten aus Südwesteuropa (Planipennia, Coniopterygidae). Reichenbachia 14(30):237–243, 5 figs.
- OLIVIER, G. A. 1811. ‘Myrméléon, *Myrmeleon*.’ Pp. 115–128 and ‘Némoptère, *Nemoptera*.’ Pp. 175–179 *in* Encyclopédie Méthodique d’Histoire Naturelle, Insectes. Vol. 8. Paris.
- OSWALD, J. D. 1987. A new species of *Micromus* from the western United States (Neuroptera: Hemerobiidae). Journal of the New York Entomological Society 95(1):1–4, 11 figs.
- . 1988. A revision of the genus *Sympherobius* Banks (Neuroptera: Hemerobiidae) of America north of Mexico with a synominal list of the world species. Journal of the New York Entomological Society 96(4):390–451, 114 figs.
- . 1993. Revision and cladistic analysis of the world genera of the family Hemerobiidae (Insecta: Neuroptera). Journal of the New York Entomological Society 101:143–299, 222 figs, 10 tables, 5 appendices.
- OSWALD, J. D. AND PENNY, N. D. 1991. Genus-group names of the Neuroptera, Megaloptera and Raphidioptera of the World. Occasional Papers of the California Academy of Sciences 147:1–94.
- PARFIN, S. I. 1952. The Megaloptera and Neuroptera of Minnesota. American Midland Naturalist 47:421–434.
- . 1956. Taxonomic notes on *Kimminsia* (Neuroptera: Hemerobiidae). Proceedings of the Entomological Society of Washington 58:203–209, 15 figs.
- . 1958. Notes on the Bionomics of the Mantispidae. Entomological News 69:203–207.
- PARFIN, S. I. AND A. B. GURNEY. 1956. The spongillaflies, with special reference to those of the Western Hemisphere (Sisyridae, Neuroptera). Proceedings of the United States National Museum 105:421–529, 24 text figs., 3 pls., 24 figs.
- PENNY, N. D. [1982a] 1981. Review of the generic level classification of the New World Ascalaphidae (Neuroptera). Acta Amazônica 11:391–406, 11 figs.
- . 1982b. Review of the generic level classification of New World Mantispidae (Neuroptera). Acta Amazônica 12(1):209–223, 22 figs., 2 tables.
- . 1996. A remarkable new genus and species of Ithonidae from Honduras (Neuroptera). Journal of the Kansas Entomological Society 69(1):81–86.
- POIRRIER, M. A. 1969. Some fresh-water sponge hosts of Louisiana and Texas spongilla-flies with new locality records. American Midland Naturalist 81:573–575.
- POIRRIER, M. A. AND R. W. HOLZENTHAL. 1980. Records of spongilla-flies (Neuroptera: Sisyridae)

- from Mississippi. *Journal of the Mississippi Academy of Sciences* 25:1–2.
- POOLE, R. W. AND P. GENTILI. 1996. *Nomina Insecta Nearctica*, A Check List of the Insects of North America. Volume 2: Hymenoptera, Mecoptera, Megaloptera, Neuroptera, Raphidioptera, Trichoptera. Entomological Information Services.
- PROCTER, W. 1946. Biological survey of the Mount Desert Region incorporated. Part VII. The insect fauna. The Wistar Institute of Anatomy and Biology, Philadelphia. 566 pp.
- PROVANCHER, L. 1875. Description de plusieurs insectes nouveaux. *Le Naturaliste Canadien* 7:247–251.
- . 1877. Faune Canadienne (Petite Faune Entomologique du Canada). Les Névroptères. *Le Naturaliste Canadien* 9:38–43, 84–90, 118–123, 173–176, 201–205, 209–217, 241–244, 257–269.
- RAMBUR, J. P. 1842. Histoire naturelle des insectes, Névroptères. Librairie encyclopédique de Roret. Fain et Thunot, Paris. 534 pp.
- REDBORG, K. E. 1976. Species of Mantispidae (Neuroptera) at Dixon Springs Agricultural Center. Illinois Agricultural Experiment Station, Dixon Springs Agricultural Center Bulletin 4:17–18.
- REDBORG, K. E. AND MACLEOD, E. G. 1983. *Climaciella brunnea* (Neuroptera: Mantispidae): a mantispid that obligately boards spiders. *Journal of Natural History* 17:63–73, 1 fig., 6 tables.
- . 1985. The developmental ecology of *Mantispa uhleri* Banks (Neuroptera: Mantispidae). *Illinois Biological Monographs*. 53:1–130. 18 figs, 30 tables.
- REHN, J. W. H. 1939. Studies in North American Mantispidae (Neuroptera). *Transactions of the American Entomological Society* 65:237–264, 22 figs.
- REUTER, O. M. 1894. Neuroptera Fennica. Förteckning och Beskrifning öfver Finlands Neuropteror. *Acta Societatis pro Fauna et Flora fennica* 9(8):1–36.
- RICE, M. E. 1987. Morphological variation in *Plega dactylota* (Neuroptera: Mantispidae) and status of its subspecies. *Journal of the Kansas Entomological Society* 60:341–344.
- ROBLE, S. M. 1984. First records of *Limnephilus submonilifer* (Trichoptera: Limnephilidae) and *Neohermes concolor* (Megaloptera: Corydalidae) for Kansas. *Transactions of the Kansas Academy of Science* 87(1–2):69–70.
- ROSS, A. 1949. Les Mantispidés de la Province de Québec. *Le Naturaliste Canadien* 61:5–14.
- ROSS, H. H. 1937. Studies of Nearctic aquatic insects. I. Nearctic alderflies of the genus *Sialis* (Megaloptera, Sialidae). *Bulletin of the Illinois Natural History Survey* 21:57–78.
- SAY, T. 1823. Descriptions of insects belonging to the order Neuroptera Linn. Latr., collected by the expedition authorized by J. C. Calhoun, etc. under the command of Major S. H. Long. Godman's Western Quarterly Reporter of Medical Surgical, and Natural Science 2(2):160–165.
- . 1824. Order Neuroptera. Pp. 303–310 in Narrative of an expedition to the source of the St. Peter's River, Lake Winnepeek, Lake of the Woods, &c. performed in the year 1823, by order of the Hon. J.C. Calhoun, Secretary of War, under the command of Stephen H. Long, Major U.S.T.E., W.H. Keating, ed. Vol. 2. Carey and Lea, Philadelphia.
- . 1825. *Mantispa*. Plate 25 and unpaginated letterpress [7 pp.] in *American Entomology, or descriptions of the insects of North America*. Vol. 2. Philadelphia. 1 pl., 7 figs.
- . 1839. Descriptions of new North American neuropterous insects, and observations on some already described. *Journal of the Academy of Natural Sciences of Philadelphia* 8:9–46.
- SCHEIRING, J. F. 1979. New state record of fish fly *Nigronia serricornis* (Say) in Alabama (Megaloptera: Corydalidae). *Entomological News* 90:176.
- SCHIMMER, H. 1865. Description of the imago and larva of a new species of *Chrysopa*. *Proceedings of the Entomological Society of Philadelphia* 4:208–212.
- SCHNEIDER, W. G. 1851. *Symbolae ad monographiam generis Chrysopae*, Leach. Hirt, Vratislaviae. 178 pp., 60 pls.
- SHETLAR, D. 1977. The biosystematics of the Nearctic Ascalaphidae (Insecta: Neuropteroidea, Planipen-

- nia), with notes on biology and morphology. Unpublished Ph.D. Thesis, Pennsylvania State University. x + 233 pp., 47 figs. (University Microfilms International No. 78-3369.)
- SMITH J. B. [1900] 1899. Order Platyptera [= Megaloptera], Pp. 51-52, 2 figs. and Order Neuroptera, Pp. 53-57, 5 figs. in Insects of New Jersey. A list of species occurring in New Jersey, with notes on those of economic importance. Trenton, New Jersey. 755 pp.
- SMITH, R. C. 1922. The biology of the Chrysopidae. Memoirs of the Cornell University Agricultural Experiment Station 58:1287-1372, 10 text figs., 14 pls., 122 figs.
- . 1925. The Neuroptera and Mecoptera of Kansas. Bulletin of the Brooklyn Entomological Society 20:165-171.
- . 1931. The Neuroptera of Haiti, West Indies. Annals of the Entomological Society of America 24:798-823, 2 pls., 21 figs.
- . 1932. The Chrysopidae (Neuroptera) of Canada. Annals of the Entomological Society of America 25:579-601, 1 pl., 6 figs.
- SPENCER, G. J. 1942. A preliminary list of the Neuroptera of British Columbia. Proceedings of the Entomological Society of British Columbia 38:23-28.
- STANGE, L. A. 1961a. Lectotype designations in the New World Myrmeleontidae. Canadian Entomologist 93:674-677.
- . 1961b. New records for the genus *Nallachius* Navás (Neuroptera: Dilaridae). Pan-Pacific Entomologist 37:144.
- . 1970a. Revision of the ant-lion tribe Brachynemurini of North America (Neuroptera: Myrmeleontidae). University of California Publications in Entomology 25:1-192, 231 figs., 31 maps, 2 tables.
- . 1970b. A generic revision and catalog of the Western Hemisphere Glenurini with the description of a new genus and species from Brazil (Neuroptera: Myrmeleontidae). Los Angeles County Museum, Contributions in Science 186:1-28, 8 figs.
- . 1980. The ant-lions of Florida. I. Genera. Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Entomology Circular 215:4 unnumbered pages, 21 figs.
- . 1990. The alderflies of Florida (Megaloptera: Sialidae). Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Entomology Circular 329:2 unnumbered pages, 6 figs.
- . 1994. Reclassification of the New World antlion genera formerly included in the tribe Brachynemurini (Myrmeleontidae). Insecta Mundi 8(1):67-119.
- . 1995. Entomology Section: Insect Detection. Tri-ology 34(2):7.
- STANGE, L. A. AND R. B. MILLER. 1990. Classification of the Myrmeleontidae based on larvae (Insecta: Neuroptera). Pp. 151-169 in Advances in Neuropterology. M. Mansell and H. Aspöck, eds. Proceedings of the Third International Symposium on Neuropterology. Department of Agricultural Development, Republic of South Africa.
- STARK, B. P. AND P. K. LAGO. 1980. New records of Nearctic *Sialis* (Megaloptera: Sialidae), with emphasis on Mississippi fauna. Entomological News 91(4):117-121, 12 figs.
- . 1983. Studies of Mississippi fishflies (Megaloptera: Corydalidae: Chauliodinae). Journal of the Kansas Entomological Society 56(3):356-364, 17 figs.
- STEINMANN, H. 1964. The *Chrysopa* species (Neuroptera) of Hungary. Annales Historico-Naturales Musei Nationalis Hungarici 56:257-266.
- STEPHENS, J. F. 1829. A systematic catalogue of British insects, being an attempt to arrange all the hitherto discovered indigenous insects in accordance with their natural affinities; containing also references to every English writer on entomology, and to principal foreign authors; with all the published British genera to the present time. 2 vols. 416, 388 pp. London.
- . 1836. Illustrations of British entomology; or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food, economy, Mandibulata. Baldwin and Cradock, London. Vol. 6. 240 pp.
- STITZ, H. 1913. Mantispiden der Sammlung des Berliner Museums. Mitteilungen aus dem Zoologischen Museum in Berlin 7:1-49, 41 figs.

- STRÖM, H. 1788. Nogle Insect Larver med deres Forvandlinger. Ny Samling af det Kongelige Norske Videnskabers Selskabs Skrifter 2:375–400c 10 pls., 48 figs.
- TARTER, D. C., D. L. ASHLEY AND C. K. LILLY. 1976. New record of the alderfly *Sialis itasca* Ross, for West Virginia (Megaloptera: Sialidae). Entomological News 87(1/2):32.
- TARTER, D. C., W. D. WATKINS, D. L. ASHLEY AND J. T. GOODWIN. 1978. New state records and seasonal emergence patterns of alderflies east of the Rocky Mountains (Megaloptera: Sialidae). Entomological News 89(9–10):231–234.
- TARTER, D. C., W. D. WATKINS, M. L. LITTLE AND D. L. ASHLEY. 1977. New state record of the alderfly *Sialis concava* Banks from Cranberry Glades, West Virginia. Entomological News 88(3/4):104.
- TARTER, D. C., W. D. WATKINS, M. L. LITTLE AND J. T. GOODWIN. 1976. New state records of fishflies (Megaloptera: Corydalidae). Entomological News 87(7/8):223–228, 5 figs.
- TARTER, D. C. AND J. E. WOODRUM. 1973a. First record of the alderfly, *Sialis joppa* Ross (Megaloptera: Sialidae), in West Virginia. Proceedings of the West Virginia Academy of Sciences 45(2):165–167.
- . 1973b. Distribution and new record of the alderfly *Sialis* (Megaloptera: Sialidae) in West Virginia. Entomological News 84:147–148.
- TASCHENBERG, E. L. 1879. Die Arten der Gattung *Myrmecoleon* Br. und *Ascalaphus* des Zoologisches Museums der Universität Halle. Zeitschrift für die gesamte Naturwissenschaft 52:174–231.
- TAUBER, C. A. 1969. Taxonomy and biology of the lacewing genus *Meleoma* (Neuroptera: Chrysopidae). University of California Publications in Entomology 58:1–94, 67 figs., 7 pls.
- . 1975. Larval characteristics and taxonomic position of the lacewing genus *Suarius*. Annals of the Entomological Society of America 68(4):695–700, 4 figs.
- TAUBER, C. A. AND M. J. TAUBER. 1973. Diversification and secondary intergradation of two *Chrysopa carnea* strains (Neuroptera: Chrysopidae). Canadian Entomologist 105(9):1153–1167, 4 figs, 5 tables.
- TAYLOR, C. 1862. An orthopterian defense. Harper's New Monthly Magazine 24:490–496.
- TENNESSEN, K. J. 1968. Four new species records of *Sialis* (Megaloptera: Sialidae) for Wisconsin. Wisconsin Academy of Sciences 53:185–196.
- THRONE, A. L. 1971a. The Neuroptera - Suborder Planipennia of Wisconsin. Part I - Introduction and Chrysopidae. Michigan Entomologist 4:65–78, 15 figs., 3 tables.
- . 1971b. The Neuroptera - Suborder Planipennia of Wisconsin. Part II - Hemerobiidae, Polystoechotidae and Sisyridae. Michigan Entomologist 4:79–87, 18 figs.
- . 1972. The Neuroptera - Suborder Planipennia of Wisconsin. Part III - Mantispidae, Ascalaphidae, Myrmeleontidae and Coniopterygidae. The Great Lakes Entomologist 5(4):119–128, 17 figs, 1 table.
- TILLYARD, R. J. 1923. Descriptions of new species and varieties of lacewings (Order Neuroptera Planipennia) from New Zealand, belonging to the families Berothidae and Hemerobiidae. Transactions of the New Zealand Institute, Wellington 54:217–225, 5 figs.
- TJEDER, B. 1960. Neuroptera from Newfoundland, Miquelon, and Labrador. Opuscula Entomologica 25:146–149.
- . 1967. Kullabergs nätringar, näbbsländor, nattsländor och harkrankar. Kullabergs Natur 12:1–21, 1 fig.
- TOWNSEND, L. H. 1939. A new species of *Sialis* (Megaloptera, Sialidae) from Kentucky. Proceedings of the Entomological Society of Washington 41:224–226, 6 figs.
- TSUKAGUCHI, S. 1995. Chrysopidae of Japan (Insecta, Neuroptera). Osaka, Japan. 223 pp. (Privately published.)
- TULLGREN, A. 1906. Zur Kenntnis schwedischer Coniopterygiden. Arkiv för Zoologi 3(5):1–15, 9 figs.
- VAN DYKE, E. C. 1945. A new dobsonfly (neuroptera) from California. Pan-Pacific Entomologist 20:110.
- VOGTSBERGER, R. C. 1990. First United States record of *Ascalobyas* (Neuroptera: Ascalaphidae), a range

- extension from northeastern Mexico to Texas. *Entomological News* 101(3):158–160.
- WALKER, F. 1853. Catalogue of the specimens of neuropterous insects in the collection of the British Museum, Part II (pp. 193–476). (*Sialides -Nemopterides*). British Museum (Natural History), London.
- . 1860. Characters of undescribed Neuroptera in the collection of W. W. Saunders. *Transactions of the Entomological Society of London* (N. S.) 5:176–199.
- . 1866. Order Neuroptera. P. 334 in *The Naturalist in Vancouver Island and British Columbia*, J. K. Lord, ed. Vol. 2. Richard Bently, London.
- WALLENGREN, H. D. J. 1870. *Anteckningar i Entomologi. Ofversigt af Kongl. Vetenskaps-Akademiens Förhandlinger*, Stockholm 27:145–182.
- . 1871. Skandinaviens Neuroptera. *Kongl. svenska Vetenskaps-Akademiens Handlingar* (N. F.) 9(8):1–76.
- WEELE, H. W. VANDER. [1909a] 1908. Ascalaphiden. Monographisch Bearbeitet. Collections zoologiques du Baron Edmond de Selys Longchamps. Catalogue Systématique et Descriptif. Fasc. VIII. 326 pp., 254 figs., 2 pls.
- . 1909b. New genera and species of Megaloptera Latr. Notes from the Leyden Museum 30:249–264.
- WELCH, J. L. AND B. C. KONDRATIEFF. 1991. The Mantispidae (Neuroptera) of Colorado. *Journal of the Kansas Entomological Society* 64(1):69–76, 16 figs.
- WELCH, P. S. 1914. The early stages of the life history of *Polystoechotes punctatus* Fabr. *Bulletin of the Brooklyn Entomological Society* 9:1–6.
- WESMAEL, C. 1841. Notice sur les Hémérobides de Belgique. *Bulletin de l'Académie Royale des Sciences et Belles-Lettres de Bruxelles* 8(1):203–221.
- WESTWOOD, J. O. 1842. Description of some insects which inhabit the tissue of *Spongilla fluviatilis*. *Transactions of the Royal Entomological Society of London* (1)3:105–108. 1 pl, 12 figs.
- WHEELER, A. G., JR. 1980. First United States record of *Aleuropteryx simillima*, a predator of scale insects on ornamental juniper (Neuroptera: Coniopterygidae). *The Southwestern Entomologist* 5(1):51–52.
- WHITING, M. F. 1991a. New species of *Sialis* from southern California (Megaloptera: Sialidae). *Great Basin Naturalist* 51(4):411–413, 3 figs.
- . 1991b. A distributional study of *Sialis* (Megaloptera: Sialidae) in North America. *Entomological News* 102(1):50–56.
- WISE, K. A. J. 1973. New records in the New Zealand Neuroptera: Hemerobiidae. *New Zealand Entomologist* 5:181–185.
- WOGLUM, R. S. AND E. A. MACGREGOR. 1964. A new snakefly from southern California (Neuroptera - Raphidiidae). *Bulletin of the Southern California Academy of Sciences* 63:201–203, 1 fig.
- ZETTERSTEDT, J. W. 1840. Ordo VII. Neuroptera. Columns 1037–1074 in *Insecta Lapponica*. Voss, Lipsiae. 1140 columns. This work printed in 1140 individually numbered, double columns.



1997. "Species catalog of the Neuroptera, Megaloptera, and Raphidioptera of America north of Mexico." *Proceedings of the California Academy of Sciences, 4th series* 50, 39–114.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/53426>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/51792>

**Holding Institution**

MBLWHOI Library

**Sponsored by**

MBLWHOI Library

**Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: California Academy of Sciences

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.