neck is a duller, more cinnamomeous, hue. It does not, however, approach in intensity of coloration even the palest and dullest colored specimens of *F. moniliger* from Guatemala, Honduras, and other parts of the latter's range. Its measurements are as follows: Length, (stretched skin), about 8.00; wing, 3.55; tail, 2.23; exposed culmen, 0.82; tarsus, 1.20; middle toe, 0.80.

While there can be no doubt that this form is merely a pallid local race of *F. moniliger*, and should therefore be known as *Formicarius moniliger pallidus*, we have not yet the proof of such fact; and, in order to preserve uniformity of nomenclature in this paper, I have given it a binomial appellation, as I have done with forms which undoubtedly are conspecific with *F. analis*.

11. *Formicarius rufipectus* Salvin.

Formicarius rufipectus SALV., P. Z. S., 1866, 73, 74, pl. VIII (Veragua); 1867, 145 (do.).—SCL. and SALV., Nom. Neotr., 1873, 75 (Veragua).—SCL., Cat. B. Brit. Mus., XV, 1890, 306 (Veragua; Baisa, Ecuador?).—SALV. and GODM., Biol. Centr.-Am., Aves, II, pt. 30, 1892, 235 (Santiago de Veragua; Baisa, Ecuador?).
? *Formicarius thoracicus* "STOLZM. MS.," TACZAN. and BERL., P. Z. S., 1885, 101 (Machay, e. Ecuador).

HAB.—Veragua; eastern Ecuador?

SP. CHAR.—Above brownish black, the rump dusky rufous, the pileum tinged with rufous; lores and throat black; breast, crissum, and middle of the belly chestnut-rufous; sides of the body sooty; bill black; feet brown.

Total length, 7 inches; wing, 3.40; tail, 2.25; tarsi, 1.50; bill to the rictus, 1.10.

Similar to *F. analis* (D'ORB. and LAFR.), but at first sight distinguished by the chestnut-rufous breast. [Translation of the original description.]

According to Messrs. Salvin and Godman (Biol. Centr.-Am., Aves, II, pt. 30, pp. 235, 236), "this well-marked species comes next to *F. analis* [*i. e.*, *F. nigricapillus* Cherrie] in many of its characters, such as the absence of the white spot on the lores and the wholly black ear coverts. Its rufous breast, however, renders it readily distinguishable as well from *F. analis* as from all its congeners."

Never having seen a specimen of this very distinct species, I am unable to give a more detailed description of it.

(12 ?) *Formicarius thoracicus* Taczanowski and von Berlepsch.

Formicarius thoracicus "STOLZM. MS." TACZAN. and BERL., P. Z. S., 1885, 101 (Machay, e. Ecuador).—SCL., Cat. B. Brit. Mus., XV, 1890, 301 footnote.—SALV. and GODM., Biol. Centr.-Am., Aves, II, pt. 30, 1890, 236, sub *F. rufipectus*.

HAB.—Eastern Ecuador (Machay, altitude 5,000 feet).

SP. CHAR.—Above dusky olive-brown; entire head, with throat, black; breast dark rufous; abdomen olive-brown; under tail coverts rufous; wings blackish, the upper coverts and the outer webs of the

remiges the same color as the back; the under wing coverts varied with ochraceous and black; tail black.

Adult male.—The black occupies the whole of the head, including the throat; the upper parts are brownish olive, much darker on the rump, and changing into rusty on the upper tail coverts; fore neck and breast very dark rust-red; abdomen sooty olive, much lighter than the back; under tail coverts dark rusty. Wings blackish, the upper coverts and outer webs of the remiges the same color as the back, the under wing coverts bright ochraceous with two large black cross-bands, the inner webs of the remiges russet at the base. Tail blackish. Bill horn-black; feet deep brownish gray; iris deep brown.

Female.—Resembles the male in all particulars, and is only distinguished by the less intense rusty on the breast, extended to the middle of the abdomen and continued as a wide stripe of russet-ocher to the under tail coverts.

Male.—Length of wing 89, tail 59, bill 27, tarsus 39 millimeters.

Female.—Total length 218, spread of wing 310, wing 89, tail 60, bill 27, tarsus 38 millimeters.

The above is a free translation of the original description, cited above.

Although Messrs. Salvin and Godman (*l. c.*) consider this bird as being probably the same as *F. thoracicus* Salv., there are features in the description which seem to indicate its distinctness. For example, the following points of disagreement in the descriptions may be cited:

	<i>F. rufipectus.</i>	<i>F. thoracicus.</i>
Upper parts	Brownish black	Dusky olive-brown.
Top of head and hind neck.....	Brownish black, tipped with rufous..	Black.

It may be, however, that the two birds are really the same species, and that their apparent differences may disappear in a larger series; but until such is proven to be the case I prefer to keep them separate.

Ridgway, Robert. 1893. "Description of a new storm petrel from the coast of western Mexico." *Proceedings of the United States National Museum* 16(962), 687–688. <https://doi.org/10.5479/si.00963801.962.687>.

View This Item Online: <https://www.biodiversitylibrary.org/item/53716>

DOI: <https://doi.org/10.5479/si.00963801.962.687>

Permalink: <https://www.biodiversitylibrary.org/partpdf/53070>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.