The islands are covered with dense scrub, mostly composed of shrubs 4 to 8 feet high. No large trees except mangroves now exist, and small plants are remarkably scarce. There are no ferns or orchids, but considerable quantities of *Orchilla* moss are gathered. Formerly "porché" and "bois rose" grew to considerable size, judging from the decaying stumps and logs. A few cocoanut trees exist, mostly upon Île Michel.

Aldabra is not permanently inhabited, but there are usually a few fishermen from Seychelles living there. The whole Aldabra group, including Aldabra, Astove, Assumption, and Cosmoledo, belongs to the British colony of Seychelles, and is leased by the Government for the turtle and tortoise-shell fishery.

ASSUMPTION ISLAND.

Assumption lies 20 miles southeast of Aldabra and is about 5 miles long by $1\frac{1}{4}$ miles wide. In its physical features it is much like Aldabra, but its surface is smoother and it is not so densely covered with scrub. A considerable part of the surface is covered with "champignon," as the rugged fossil coral rock of Aldabra is called. There are two large sand dunes upon the eastern shore, about 70 feet high, which are visible from a considerable distance. No fresh water exists, unless just after a rain, when a little collects in hollows in the rock. The animal life and vegetation is similar to that of Aldabra. The little flightless rail (*Rougetius abbotti*, Ridgway)* abounds, as well as most of the other land-birds found upon the latter island. Numbers of goats run wild, having been introduced many years since from Europa Island (in Mozambique Channel).

GLORIOSO ISLAND.

Glorioso Island lies about 90 miles west-northwest of the north point of Madagascar (Cape Amber), and 120 southeast of Aldabra. It is

-
3. *Lycana telicanus*, Hüb. South Europe through East Africa to the Cape of Good Hope.

Dragonfly:

1. *Pantala flavescens*, Fab. A common East African species.

Myrmeleon. *Myrmeleon*, sp.

Mantid:

1. *Polyspilota variegata*, Oliv. An East African species.

Beetles:

1. *Oxythyrea amabilis*, Schönh.; var. Smaller than the continental forms.
2. Small scarabæid—undetermined.
3. Small longicorn—undetermined.

Wasps:

1. *Sphex*, sp.
2. *Monedula*, sp.

Other diptera:

Odontomyia, sp.

F. W. T.

* A different species from that of Aldabra, described in *The Auk*, for January, 1894.—R. R.

situated upon the south end of the Glorioso bank. The bank is about 8 miles long by 2 in width. Île du Lise lies at the north end of the bank.

Glorioso Island contains about 700 acres, being $1\frac{1}{2}$ miles long by 1 mile wide. It is partly covered with sand hills 50 to 60 feet high. Formerly it was almost entirely covered with a growth of "porché," "bois rose," "fouche" and other large trees, but at present many have been cut down. The soil is unusually fertile for an oceanic island, having been manured for ages by thousands of sea-birds. "Champignon," or fossil—coral rock, such as composes Aldabra, Cosmoledo, etc., exists in only a few spots, and the soil or sand is of fair depth. Large quantities of maize are grown. Water from wells is of poor quality. There are five species of land-birds, three of which, a sun bird,* *Zosterops*,† and a bulbul‡ are probably peculiar.

Common fowls run wild in the jungle in considerable numbers. They are very shy and not easy to shoot. Among sea-birds there is a booby, which seems to be peculiar to the island.§ They breed in large numbers upon the "fouche" trees, in company with frigates and common boobies.

Upon the neighboring small islet of Lise vast numbers of "Wide-awake" terns (*Sterna fuliginosa*) breed, together with "General" and "Capucin" boobies (*Sula cyanops* and *S. piscator*). A gecko (*Hemidactylus mabouia*) and two other lizards, (*Ablepharus gloriosus* Stejneger, *Zonosaurus madagascariensis*) are plentiful. Numbers of wild cats range the jungle, so that birds are far less numerous, individually, than in Aldabra.

* *Cinnyris souimanga*.

† *Z. madagascariensis*.

‡ *Ixocincla madagascariensis*?

§ Two species of boobies were collected by Dr. Abbott, *Sula cyanops* and *S. piscator*, but both of these are of very wide distribution.—R. R.

REMARKS ON JAPANESE QUAILS.

BY

LEONHARD STEJNEGER.

When writing my remarks on the Japanese quails recently sent me by Dr. Ijima (Proc. U. S. Nat. Mus., XVI, 1893, p. 623) I had not yet seen Mr. Oglivie Grant's "Notes on the Genus *Coturnix*" (Ann. Mag. Nat. Hist. (6) x, 1892, pp. 166-173), in which he advances the theory, or rather announces as a demonstrated fact, that there are two species occurring in Japan (and other portions of eastern Asia) viz: *C. coturnix*, the typical European species, and *C. japonica*, which, in their purity, may be distinguished as follows:

- a*¹ Feathers on throat and chin short and rounded.
 - b*¹ A black band down the middle of the throat *C. coturnix* ♂
 - b*² No black band down the middle of the throat.
 - c*¹ Chin and throat white *C. coturnix* ♀
 - c*² Chin and throat dark vinaceous-cinnamon [dull brick-red, O. G.]
C. japonica ♂ ad.
- a*² Feathers on throat and chin elongate and lanceolate.
 - b*¹ Entire throat white *C. japonica* ♀
 - b*² Middle of throat suffused with dark cinnamon-rufous *C. japonica* ♂ juv.

The multitude of specimens which do not fall within the limits I have here drawn, he disposes of by the following remark: "The intermediate forms are, as I shall presently show, undoubtedly the results of interbreeding." But I am sorry to say that he does not show this, for there is no discussion of the material upon which he bases his remarks, nor are we furnished with a list of his specimens with the accompanying data upon which we might be enabled to base an opinion as to the correctness of his conclusions. All he gives us is a bare assertion to the above effect, the essential part of which is as follows: "In Japan and China the migratory Quail (*C. coturnix*), as already pointed out, inhabits the same tract of country during the breeding season as *C. japonica*, and there can not be the slightest doubt that the two species frequently interbreed, with the result that all sorts of intermediate hybrids are produced. These intermediate plumages are most noticeable among the male hybrids. For instance, some have the dull brick-red throat of *C. japonica* and the black anchor-shaped mark of *C. coturnix*, others have only the upper two-thirds of the throat dull red and the lower third white, while again a third lot have, in addition, a black band down the center of the red part, and all kinds of intermediate stages between these three examples may be found."

It is evidently in order to meet the objection that rufous-throated males are often found in Europe that he makes the following remark: "Equally also, though of secondary importance, *C. coturnix* interbreeds freely with the red-throated resident race (*C. capensis*)* in South Africa and the islands surrounding the coast, and the results are seen in the many male birds from South Africa and Southern Europe, etc., in which the white parts on the sides of the head and throat are more or less suffused with the bright rufous chestnut of the resident bird."

But this is hardly more than a postulate, and it is, in fact, somewhat difficult to see how such a hybridization can take place between a resident species and a subspecies (and he calls them only "races"). the results to be found both among the residents and the migrants, The facts are that these so-called intermediates between *C. capensis* and *C. coturnix* are not only found in South Africa and Southern Europe, but that they are quite common in Central Europe, as evidenced by the detailed description of the throat color and markings by Naumann (Naturg. Vög. Deutschl., VI, 1833, pp. 578, 579, and particularly pp. 580-581). From his remarks it will be seen that the male quails in Germany vary as much and almost in the same way as the Japanese and Chinese birds described by Mr. Ogilvie-Grant, and by him asserted to be hybrids.

Looking over my material I find nothing in it to contradict the supposition that the color and markings of the throat of the male Japanese bird is subject to as much individual variation as in the German bird, and I can see no reason for regarding these various plumages as "intermediate stages" or "hybrids." I think such a view also effectually disposes of the somewhat curious peculiarity that "these intermediate plumages are most noticeable among the male hybrids."

Mr. Ogilvie-Grant does not mention any specimens in which the supposed hybridism is expressed in an intermediate state of the elongated throat feathers. On the other hand, in the males he regards the presence of these specialized feathers as the sign of youth, in support of which he mentions the case of "a rather more mature male" in which "one side of the throat has lost the immature elongate feathers like those of the female and assumed the short, rounded, dull rufous feathers characteristic of the male adult," but all other data which would make it profitable to discuss the case are wanting.

I now turn to the material before me.

(1) U. S. Nat. Mus., No. 95980; ♂ ad.; collected by Blakiston at Sapporo, Yezo, May, 11, 1877. In coloration this specimen is exactly like the front figure of Fauna Jap. Av., pl. lxi, with the exception that the posterior half of the superciliary stripe is white and the anterior half spotted with white; the flank feathers are less marked with blackish; throat feathers, both in the middle and on the sides, short and rounded.

* I would suggest that the proper name of this subspecies is *Coturnix coturnix africana* (Schlegel) (see Fauna Jap. Aves, p. 103). There is no reference to this name in Mr. Ogilvie-Grant's synonymy.

(2) Imp. Mus., Tokyo, No. 2168; Province of Owari, Hondo; Mr. Ota coll.; no date. Coloration almost identical with the foregoing specimen, though with a faint blackish wash on the middle of the throat. Otherwise the similarity of the two specimens is so complete that I have no hesitation in pronouncing it of the same sex and age as the foregoing. Throat feathers in the middle short and rounded; on the sides perceptibly longer and narrower, though not pointed.

(3) Imp. Mus., Tokyo, No. 2170; Province of Owari, Hondo; no date. Coloration like 1 and 2, but chin, middle of throat, and first lateral branch of the throat patch black; flanks as in figure quoted above; throat feathers much as in 2.

The above three specimens thus appear to be fully adult summer males.

(4) U. S. Nat. Mus., No. 109409; ♂; Shimosa, Hondo; December 22, 1885. General coloration much as the above, but all light markings, including superciliary stripe, more strongly washed with buff; breast deeper ferruginous; chest feathers with a large chestnut spot in either web, but no black spots; feathers on middle of throat dull cinnamon-rufous, with broad white terminal margins; those on the sides of throat and on cheeks cinnamon-rufous with a white shaft streak, with terminal black spots on the cheeks. Middle throat feathers rounded; lateral ones elongated and pointed; a few of the latter still in their sheaths.

The richness of the coloration, especially that of the breast, leads me to believe that this is a fully adult male in, at least, its second winter.

(5) U. S. Nat. Mus., No. 91582; ♂; Yokohama, Hondo; April 4, 1883; P. L. Jouy coll. Considerably paler than any of the foregoing; throat and upper fore neck white, with a narrow dusky band down the middle of the throat, united below with a semicircular dusky line descending from the ear; lateral throat feathers edged with cinnamon-rufous; middle ones more or less suffused with the same color and tipped with whitish; a few blackish spots on the chest; all the throat feathers elongate and pointed.

(6) U. S. Nat. Mus., No. 95983; sex not given; A. Owston coll.; no date. Practically identical with foregoing, except that only lateral throat feathers are pointed, the median ones being short and rounded.

(7) U. S. Nat. Mus., No. 95982; ♂; Nagasaki, Kiusiu; January 1, 1877; F. Ringer coll. Like the foregoing, but throat band twice as broad, occupying the whole middle part and continued backward beyond the first semicircular line, though not reaching the second; throat feathers all strongly elongated and pointed.

(8) U. S. Nat. Mus., No. 114127; ♂; Fusan, Korea; November 21, 1885; P. L. Jouy coll. Like the foregoing, but throat and upper fore neck pure white, with a faint indication of a dusky band down the middle of the throat, caused by the dark bases of the feathers shining through the white tips; two semicircular black lines, the upper one imperfect on the median line; median throat feathers rounded, lateral ones moderately elongated, pointed.

Stejneger, Leonhard. 1894. "Remarks on Japanese quails." *Proceedings of the United States National Museum* 16(974), 765–769.

<https://doi.org/10.5479/si.00963801.974.765>.

View This Item Online: <https://www.biodiversitylibrary.org/item/53716>

DOI: <https://doi.org/10.5479/si.00963801.974.765>

Permalink: <https://www.biodiversitylibrary.org/partpdf/53082>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.