When using this piece of apparatus, the light should be opposite to the small aperture (*a*), and in a position at right angles to the body of the microscope. When the lamp is used alone, an image of its flame will be seen upon the object; but the whole field of view may be illuminated, or unilateral light may be obtained, by placing a small condenser in different positions before the lamp.

Under this method of illumination, which is available with the highest powers, the appearances presented by objects are very remarkable, and they vary exceedingly according to the character and condition of the specimen. It is in all cases best to have the object uncovered. The subject is of great importance and interest, and requires thorough investigation; but this, from the moderate cost of the piece of apparatus (10s. 6d.), is within the reach of every microscopist.

Observations on some Lepidosirens (Protopterus annectens, Owen) which have lived in the Menagerie of Reptiles in Paris, and formed their Cocoon there. By A. DUMÉRIL.

M. A. Duméril has observed the formation of the cocoon by two specimens of *Lepidosiren* living in the Menagerie at the Jardin des Plantes. About the 20th February last, these two animals showed indications of a desire to shelter themselves in the soft soil at the bottom of the aquarium; and as previous specimens had died, owing to the necessary change of conditions not being realized for them, the water was nearly all drawn off from the aquarium, so as to leave the mud at its bottom as nearly as possible in the same condition as the rice-fields inhabited by the *Lepidosiren* at the approach of the dry season. In three weeks the ground gradually hardened, forming a mass with fissures in several parts; no trace of the animals was to be detected.

The fissures in the mass of dried mud enabled portions of it to be removed; and the two *Lepidosirens* were found, about eleven weeks after their disappearance, enveloped in regular cocoons. The block of dried mud, which is preserved in the museum, exhibits a cavity moulded upon the cocoon, with the walls perfectly smooth and lined with a strongly adherent portion of the cocoon.

This cocoon is produced by a mucous secretion. Its brown colour might lead to the supposition that it is formed of dried leaves; but when examined by Professor Decaisne, it presented no trace of vegetable structure; and when burnt, it diffused the characteristic odour of animal substances.

The *Lepidosirens*, when desirous of burying themselves, emit an abundant mucosity from the surface of the body. This coats and agglutinates the portions of the soil which they traverse, so that the walls of the subterranean canal made by each animal and remaining open after desiccation are smooth and appear polished; then when the animal stops, this exudation acquires the consistency of a membranous envelope. The *Lepidosirens* when examined by M. Duméril were still alive, as was proved by their slight movements when touched; they have since died.—*Comptes Rendus*, January 8, 1866, p. 97.

Duméril, Auguste Henri André. 1866. "Observations on some Lepidosirens (*Protopterus annectens*, Owen) which have lived in the Menagerie of reptiles in Paris, and formed their cocoon there." *The Annals and magazine of natural history; zoology, botany, and geology* 17, 160–160.

View This Item Online: <https://www.biodiversitylibrary.org/item/54464>

Permalink: <https://www.biodiversitylibrary.org/partpdf/53714>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.