

**KEY TO THE SUBFAMILIES AND TRIBES OF THE NEW WORLD
COREIDAE (HEMIPTERA), WITH A CHECKLIST OF
PUBLISHED KEYS TO GENERA AND SPECIES**

RICHARD J. PACKAUSKAS

Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, Connecticut 06269-3043.

Abstract. — Keys are presented to the three subfamilies and 15 tribes of Coreidae which occur in the New World. A list of published keys to New World genera and species is also presented.

Key Words: Hemiptera, Heteroptera, Coreidae, New World, tribal keys, subfamily keys

Scientific work must be accessible. In taxonomy, this places upon specialists the responsibility to provide keys and catalogs, in addition to descriptive and phylogenetic research. The publishing of keys and catalogs opens the literature to others, and often stimulates further work. Studies on the New World Coreidae have languished for half a century. Neglect of these often large and abundant and occasionally economically important bugs has been due, I believe, to the lack of means to identify them. Most literature treating the New World Coreidae has been restricted to the North American fauna, except for the recent efforts of Brailovsky (1983 to present). Torre-Bueno (1941) provided synoptic keys to species of North America and Blatchley (1926) covered the family for eastern North America. More recently, Froeschner's (1988) catalog to the North American Coreidae and Baranowski and Slater's Coreidae of Florida (1986) have contributed greatly to our understanding of this family. However, except for the outdated and untranslated key of Stål (1867), no key exists to the subfamilies and tribes of the New World Coreidae.

The present classification of New World Coreidae is based primarily on Stål's (1867,

1870) work which is the only comprehensive treatment for the identification of subfamilies, tribes, and genera. But in the century since, there have been many changes in the higher classification, including descriptions of new tribes (Blatchley 1926, O'Shea and Schaefer 1978, Brailovsky 1988a), and the placement of three New World genera in the tribe Hydarini, until recently thought to occur only in the Old World (Brailovsky 1988b). The New World component of this family presently contains about 136 genera and 838 species (Packauskas, unpublished).


This paper provides an updated summary of the classification and a key to the subfamilies and tribes of the New World Coreidae with references to published keys to genera and species. All groups keyed here are exclusively New World except for the Anisoscellini (mostly New World with the exception of one species) and the Pseudophloeinae, Coreini, and Hydarini. It is my hope that this information will stimulate further research on the family, and aid collectors and curators in sorting unidentified material to make it more readily available for study.

It is important to note that "Anisosceli-


dini,” “Leptoscelidini,” and “Meropachydinae” have been spelled that way for many years and by many people. Nevertheless, the Greek words from which these family-group names are formed, -scelis and -pachys, do not take d in their combining roots. As a result, the correct spelling of these groups is “Anisoscelini,” “Leptoscelini,” and “Meropachyinae.”

KEY TO THE SUBFAMILIES AND TRIBES
OF NEW WORLD COREIDAE


- 1. Tibiae not sulcate; metathoracic scent gland orifices with a single large or two nearly fused discs (auricles) (Fig. 1); meso- and metasterna sulcate; males with a bilobed genital capsule opening filled by the clearly visible parameres (Fig. 2); all species less than 12 mm long Pseudophloeinae Stål, 1867
- Tibiae sulcate (sometimes shallowly) or, if appearing nonsulcate, without the above combination of characters; length variable 2
- 2. Hind tibiae with an apical tooth or spine (may be obscured by setae, Fig. 3) Meropachyinae Stål, 1867 . . . 3
- Hind tibiae unarmed at apex Coreinae Leach, 1815 . . . 5
- 3. Scutellum elongate, extending beyond clavus; posterior acetabula projecting laterally, visible from above (Fig. 4); posterior tibiae broadly curved distally Meropachyini Stål, 1867
- Scutellum triangular, about as long as broad, shorter than clavus; posterior acetabula not or only slightly projecting laterally, not visible from above (Fig. 5); posterior tibiae never curved 4
- 4. Posterior femora longer than abdomen, narrow and cylindrical basally, swollen apically; body small, robust, thick, somewhat compressed laterally; scutellum more or less elevated, terminating in a slightly elevated knob Merocorini Stål, 1870
- Posterior femora shorter than abdomen, swollen for entire length; body larger, more elongate, not compressed laterally; scutellum more or less flat, if elevated basally, never terminating in a raised knob Spathophorini Kormilev, 1954
- 5. Tylus projecting conspicuously beyond juga as a distinct knob or strongly compressed plate (Figs. 6, 7); juga strongly deflexed Acanthocephalini Stål, 1870
- Tylus not or only vaguely projecting beyond juga; juga variable 6
- 6. With the combination of hind femora spinose or tuberculate, strongly incrassate in males; head not over half as long as pronotum and not extending past antenniferous tubercles or very slightly so; antenniferous tubercles occupying most of anterior head width, space between them rarely wider than one tubercle (Figs. 8, 9) 7
- Not with this combination of characters 8
- 7. Metathoracic scent gland with a single fused auricle (sometimes bilobed, Fig. 11); ocellar tubercles large; usually with a spine present on antenniferous tubercles (Fig. 8); parameres membranous Acanthocerini Bergroth, 1913
- Metathoracic scent gland with two completely separated auricles (area between them depressed, Fig. 10); ocellar tubercles small; antenniferous tubercles never armed (Fig. 9); parameres sclerotized Nematopodini Amyot & Serville, 1843
- 8. Hind tibiae expanded, expansion slight in some taxa (Fig. 13) Anisoscelini Amyot & Serville, 1843
- Hind tibiae never expanded 9
- 9. Juga extended and exceeding or equal to tylus in length (Fig. 15) 10
- Juga deflexed or at least not exceeding nor equal to tylus in length 11
- 10. Head shorter than pronotum, antennae terete; brachypterous. Barreratalpini Brailovsky, 1988a
- Head equal in length to pronotum; antennae triangular in cross section; macropterous Chelinideini Blatchley, 1926
- 11. All femora spinose below, profemoral spines sometimes minute; bucculae never reaching midlength of head; head porrect; juga and tylus extended forward; rostrum reaching or extending onto abdomen, first segment reaching or extending past base of head Leptoscelini Stål, 1867
- Not with this combination of characters. 12
- 12. Distance between hind coxae equal to distance from coxa to lateral margin; antenniferous tubercles prominent, subcontiguous above deflexed tylus and juga Spartocerini Amyot & Serville, 1843
- Distance between hind coxae much narrower than distance from coxa to lateral margin, if distances nearly equal then head elevated between antenniferous tubercles; tylus and juga variable 13
- 13. Third antennal segment expanded, other segments terete (Fig. 12); tylus and juga deflexed, not filling space between antenniferous tubercles; all femora armed distally Chariesterini Stål, 1867
- Third antennal segment never expanded, if appearing expanded, then other segments also



Figs. 1-5. 1. Metathoracic scent gland opening, *Coriomeris occidentalis* Dolling & Yonke (Pseudophloeinae). 2. Genital capsule, *C. occidentalis*. 3. Hind femur and tibia, *Merocoris distinctus* Dallas (Meropachyinae). 4. Dorsal view, *Hirulcus alternatus* (Dallas) (Meropachyini). 5. Dorsal view, *Diariptus* sp. (Spathophorini). act = acetabulum, clv = clavus, cx = coxa, par = paramere, sct = scutellum.



Figs. 6–11. Heads of coreids: 6. *Lucullia flavovittata* Stål (Acanthocephalini). 7. *Nyttum punctatum* (Dallas) (Acanthocephalini). 8. *Crinocerus sanctus* (Fabricius) (Acanthocerini). 9. *Mozena* sp. (Nematopodini). Meta-thoracic scent gland openings: 10. *Mozena* sp. (Nematopodini). 11. *Sagotylus confluens* (Say) (Acanthocerini). atu = antenniferous tubercle, aur = auricle, cx = coxa, otu = ocellar tubercle, sp = spine.



Figs. 12–16. 12. Antenna, *Chariesterus antennator* (F.) (Chariesterini). 13. Hind tibia, *Tarpeius brevicornis* Stål (Anisoscelini). 14. Antenna, *Madura perfida* Stål (Hydarini). 15. Head, *Cheliniidea vittiger* Uhler (Chelini-deini). 16. Abdomen (Discogastrini). jug = jugum, spr = spiracle, tyl = tylus.

- appearing expanded; tylus and juga usually extended anteriorly between antenniferous tubercles; femora rarely armed 14
14. Spiracles situated twice as far from posterior border of middle abdominal segments (III–VI) as from anterior border (Fig. 16) Discogastrini Stål, 1867
- Spiracles equidistant from anterior and posterior borders of middle abdominal segments (III–VI) 15
15. Fourth antennal segment flattened; third antennal segment over twice as long as second (Fig. 14) Hydarini Stål, 1873¹
- Fourth antennal segment not flattened; third antennal segment usually shorter than second, never twice as long as second Coreini Leach, 1815

LIST OF KEYS TO NEW WORLD GENERA AND SPECIES OF COREIDAE

Key to North American species (Torre-Bueno 1941)

Pseudophloeinae (3 genera)

Key to genera (Dolling 1977)

Key to species of *Ceraleptus* Costa (Froeschner 1963)

Key to species of *Vilga* Stål (Dolling 1977)

Key to North American species of *Coriomeris* Westwood (Dolling and Yonke 1976)

Meropachyinae (15 genera)

Key to tribes and genera of Meropachyinae (Kormilev 1954) (With the exception of *Accocopus* Stål, 1864, *Allopeza* Bergroth, 1912, *Diariptus* Stål, 1859, *Eretmophora* Stein, 1860, and *Hoplophthonia* Schmidt, 1911.)

Coreinae

Acanthocephalini (15 genera)

Key to species of *Acanthocephala* Laporte of North and Central

America (Gibson and Holdridge 1918a)

Key to species of *Nyttum* Spinola (Dolling and Casini 1988)

Key to species of *Petalops* Amyot & Serville (Brailovsky 1991)

Key to species of *Stenametapodus* Breddin (Brailovsky 1984a)

Acanthocerini (16 genera)

Key to genera of Acanthocerini (O'Shea 1980a) (also see Brailovsky 1988c)

Key to species of *Moronopelios* Brailovsky (Brailovsky 1988c)

Key to species of *Thlastocoris* Mayr (Brailovsky 1990a)

Anisoscelini (9 genera, or more, see Osuna 1984)

Key to genera of Anisoscelini (Osuna 1984)

Key to species of *Leptoglossus* Guérin-Meneville (Allen 1969)

Key to species of *Narnia* Stål (Gibson and Holdridge 1918b)

Barreratalpini (1 genus) (see Brailovsky 1988a)

Chariesterini (4 genera)

Key to genera of Chariesterini (Yonke 1972)

Key to species of *Chariesterus* Laporte (Ruckes 1955)

Key to species of *Ruckesius* Yonke (Yonke 1972)

Chelinideini (1 genus)

Key to species of *Chelinidea* Uhler (Herring 1980)

Coreini (30 genera)

Key to species of *Acidomeria* Stål (Brailovsky 1983a)

Key to species of *Althos* Kirkaldy (Brailovsky 1990d)

Key to species of *Anasa* Amyot & Serville (Brailovsky 1985)

Key to species of *Catorhintha* Stål (Brailovsky and Garcia 1987)

Key to Mexican species of *Cebrenis* Stål (Brailovsky 1984b)

Key to species of *Collatia* Stål (Brailovsky 1990b)

¹ Brailovsky (1988b) placed three New World genera in this mostly Old World tribe. Ahmad (1970) raised Hydarini to subfamily status, but this interpretation has not been universally accepted.

- Key to species of *Daphnasa* Brailovsky (Brailovsky 1984c)
- Key to species of *Himella* Dallas (Brailovsky and Barrera 1986a)
- Key to species of *Hypselonotus* Hahn (Whitehead 1974)
- Key to species of *Nirovecus* Stål (Brailovsky 1984d, 1990b)
- Key to species of *Paryphes* Burmeister (Brailovsky 1986a)
- Key to species of *Scolopocerus* Uhler (Brailovsky 1989a)
- Key to species of *Sethenira* Spinola (Brailovsky 1988d)
- Key to species of *Sphictyrtus* Stål (Brailovsky and Meléndez 1989)
- Key to species of *Stenoprasia* Horvath (Brailovsky 1986b)
- Key to species of *Sundarus* Amyot & Serville (Brailovsky 1988e)
- Key to species of *Vazqueticoris* Brailovsky (Brailovsky 1990d)
- Discogastrini (7 genera)
 - Key to genera of Discogastrini (Brailovsky 1984e)
 - Key to species of *Cnemomis* Stål (Brailovsky and Barrera 1986b)
 - Key to Mexican species of *Savius* Stål (Brailovsky 1986c)
 - Key to species of *Scamurius* Stål (Brailovsky 1986d)
- Hydarini (3 genera)
 - Key to genera and species of Hydarini (Brailovsky 1988b)
- Leptoscelini (4–9 genera)
 - Key to species of *Dalmatomammurius* Brailovsky (Brailovsky 1990c) (tentative tribal placement)
 - Key to species of *Plunentis* Stål (Brailovsky 1989b) (tentative tribal placement)
- Nematopodini (15 genera)
 - Key to genera of Nematopodini (O'Shea 1980b)
 - Key to species of *Cnemyrtus* Stål (Brailovsky 1989b)

- Key to species of *Curtius* Stål (Brailovsky 1986e)
- Key to species of *Mozena* Amyot & Serville (Hossain 1970)
- Key to Mexican species of *Nematus* Berthold (Brailovsky and Zurbia 1983)
- Key to Mexican species of *Piezogaster* Amyot & Serville (Brailovsky and Barrera 1984)
- Key to species of *Quintius* Stål (Brailovsky & Berrera 1986c)
- Key to species of *Zoreva* Amyot & Serville (Brailovsky & Barrera 1982)
- Spartocerini (4 genera)
 - Key to species of *Sephina* Amyot & Serville (Brailovsky 1983b)
 - incertae sedis (3 genera)
 - Molchina* Amyot & Serville 1843
 - Mammurius* Stål 1862
 - Rochrosoma* Reed 1901

ACKNOWLEDGMENTS

I am grateful to Ms Mary Jane Spring for help with the illustrations, T. A. Suits, Department of Modern and Classical Languages, University of Connecticut, for advice on the spelling of Anisoscelini, Leptoscelini, and Meropachyinae, and Carl W. Schaefer for help in preparing this paper, and also a special thank you to the anonymous reviewer and Thomas J. Henry, both of whose comments helped tighten up the key.

LITERATURE CITED

- Ahmad, I. 1970. Some aspects of the female genitalia of *Hygia* Uhler 1861 (Coreidae: Colpurinae) and their bearing on classification. *Pakistan Journal of Zoology* 2: 235–243.
- Allen, R. C. 1969. A revision of the genus *Leptoglossus* Guérin (Hemiptera: Coreidae). *Entomologia Americana* 45: 35–140.
- Amyot, C. J. B. and A. Serville. 1843. *Histoire Naturelle des Insectes—Hémiptères*. Librairie Encyclopédique de Roret, Paris. 675 pp.
- Baranowski, R. M. and J. A. Slater. 1986. Coreidae

- of Florida. Arthropods of Florida and Neighboring Land Areas. Vol. 12. Gainesville, Florida. 82 pp.
- Bergroth, E. 1912. Notes on Coreidae and Neididae. *Annales de la Société Entomologique de Belgique* 56: 76-93.
- . 1913. Supplementum Catalogi Heteropterorum Bruxellensis II. Coreidae, Pyrrhocoridae, Colobathristidae, Neididae. *Mémoires de la Société Entomologique de Belgique* 22: 126-183.
- Blatchley, W. S. 1926. Heteroptera or true bugs of eastern North America. The Nature Publishing Co., Indianapolis, Indiana. 1116 pp.
- Brailovsky, H. 1983a. El género *Acidomeria* Stål con descripción de una nueva especie (Hemiptera-Heteroptera-Coreidae-Coreini). *Folia Entomológica Mexicana* 56: 3-20.
- . 1983b. El género *Sephina* Amyot y Serville en Venezuela con descripción de una nueva especie (Hemiptera-Heteroptera-Coreidae-Spartocerini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 53: 277-283.
- . 1984a. Una nueva especie del género *Stenometapodus* Breddin y algunas notas acerca de *Empedocles tenuicornis* (Westwood) (Hemiptera-Heteroptera-Coreidae-Acanthocephalini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 55: 63-68.
- . 1984b. Hémiptera-Hétéroptera de México XXXII. El género *Cebrenis* Stål y descripción de una especie nueva (Coreidae-Coreini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 55: 123-132.
- . 1984c. Una nuevo género y tres nuevas especies de la tribu Coreini (Hemiptera-Heteroptera-Coreidae) para el continente Americano. *Anales del Instituto de Biología Universidad Nacional Autónoma México* 55: 161-176.
- . 1984d. El género *Nirovecus* Stål con descripción de una nueva especie (Hemiptera-Heteroptera-Coreidae-Coreini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 54: 97-104.
- . 1984e. Descripción de un nuevo género y una nueva especie de la tribu Discogastrini (Hemiptera; Heteroptera; Coreidae). *Folia Entomológica Mexicana* 62: 17-26.
- . 1985. Revision del género *Anasa* Amyot-Serville (Hemiptera-Heteroptera-Coreidae-Coreinae-Coreini). *Monografías del Instituto de Biología Universidad Nacional Autónoma México*, No. 2: 1-266.
- . 1986a. Nuevos dato distribucionales y descripción de tres nueva especies del género *Parryphes* Burmeister (Hemiptera-Heteroptera-Coreidae-Coreini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 56: 101-128.
- . 1986b. Revision del género *Stenoprasia* Horvath y descripción de dos especies nuevas (Hemiptera-Heteroptera-Coreidae-Coreini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 56: 453-464.
- . 1986c. Hemiptera-Heteroptera de Mexico XXXV. Revision de la familia Coreidae Leach. Parte I. Tribu Discogastrini Stål. *Anales del Instituto de Biología Universidad Nacional Autónoma México* 56: 401-422.
- . 1986d. Revision del género *Scamurius* Stål (Hemiptera-Heteroptera-Coreidae-Discogastrini). *Folia Entomológica Mexicana* 70: 25-52.
- . 1986e. Revision del género *Curtius* Stål, con descripción de tres nuevas especies y su ubicación dentro de la tribu Nematopodini (Hemiptera-Heteroptera-Coreidae). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 57: 269-280.
- . 1988a. Hemiptera-Heteroptera de Mexico XXXIX. Descripción de una tribu nueva de Coreidos recolectados en bambu (*Bambusa* sp.) (Coreidae-Coreinae). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 58: 155-164.
- . 1988b. La tribu Hydarini Stål, en el continente Americano con descripción de dos nuevos géneros, una nueva especie y una nueva subespecie (Hemiptera-Heteroptera-Coreidae). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 58: 623-650.
- . 1988c. Un nuevo género y dos nuevas especies de coreidos neotropicales (Hemiptera-Heteroptera-Coreidae-Acanthocerini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 58: 171-178.
- . 1988d. Dos nuevos especies del género *Sethenira* Spinola y nuevos arreglos nomenclatoriales dentro de *Acidomeria* Stål (Hemiptera-Heteroptera-Coreidae-Coreini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 58: 179-198.
- . 1988e. Revision del género *Sundarus* Amyot-Serville (Hemiptera-Coreidae-Coreini) para el continente Americano. *Anales del Instituto de Biología Universidad Nacional Autónoma México* 58: 561-622.
- . 1989a. Revision del género *Scolopocerus* Uhler con descripción de una especie nueva (Hemiptera-Heteroptera-Coreidae-Coreini) de NorteAmerica. *Anales del Instituto de Biología Universidad Nacional Autónoma México* 59: 65-76.
- . 1989b. Nuevos arreglos tribales dentro de la familia Coreidae y descripción de dos especies nuevas SudAmericanos (Hemiptera-Heteroptera). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 59: 159-180.

- . 1990a. Generos nuevos y especies nuevas de coreidos neotropicales (Hemiptera-Heteroptera-Coreidae: Acanthocerini, Leptoscelidini y Anisoscelidini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 61: 107–123.
- . 1990b. Un genero nuevo y seis especies nuevas de coreidos neotropicales (Hemiptera-Heteroptera-Coreidae-Coreinae). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 61: 257–278.
- . 1990c. Nuevos arreglos nomenclatoriales y descripción de dos especies nuevas del genero *Dalmatomammurius* Brailovsky (Hemiptera-Heteroptera-Coreidae-Leptoscelidini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 61: 343–355.
- . 1990d. Revision del complejo *Althos* con descripción de generos nuevos y especies nuevas (Hemiptera-Heteroptera-Coreidae-Coreini). *Publicaciones Especiales del Instituto de Biología Universidad Nacional Autónoma de México*. 156 pp.
- . 1991. El género *Petalops* con descripción de cinco especies nuevas (Hemiptera-Heteroptera: Coreidae: Acanthocephalini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 62: 453–468.
- Brailovsky, H. and E. Barrera. 1982. Revision del genero *Zoreva* Amyot y Serville y descripción de cinco nuevas especies (Hemiptera-Heteroptera-Coreidae-Mictini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 52: 301–345.
- . 1984. Hemiptera-Heteróptera de México XXXIII. El Género *Piezogaster* Amyot y Serville con descripción de dos nuevas especies (Coreidae-Nematopodini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 55: 133–154.
- . 1986a. El genero *Himella* Dallas con descripción de dos especies nuevas (Hemiptera-Heteroptera-Coreidae-Nematopodini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 56: 423–436.
- . 1986b. El género *Cnemomis* Stål y descripción de cuatro nuevas especies (Hemiptera-Heteroptera-Coreidae-Discogastrini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 57: 119–136.
- . 1986c. El genero *Quintiis* Stål con descripción de un subgenero nuevo y tres especies nuevas (Hemiptera-Heteroptera-Coreidae-Nematopodini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 56: 437–452.
- Brailovsky, H. and M. Garcia. 1987. Revision del genero *Catorhintha* Stål (Hemiptera-Heteroptera-Coreidae-Coreinae-Coreini). *Monografías del Instituto de Biología Universidad Nacional Autónoma México* No. 4: 1–148.
- Brailovsky, H. and V. Meléndez. 1989. Revisión del género *Sphictyrtus* Stål (Hemiptera-Heteróptera-Coreidae-Coreinae-Coreini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 60: 1–76.
- Brailovsky, H. and R. Zurbia. 1983. Hemiptera-Heteroptera de México XXVIII. Una nueva especie del género *Nematopus* Berthold (Coreidae-Nematopodini). *Anales del Instituto de Biología Universidad Nacional Autónoma México* 53: 213–218.
- Dolling, W. R. 1977. A revision of the neotropical genus *Vilga* Stål (Hemiptera: Coreidae). *Systematic Entomology* 2: 27–44.
- Dolling, W. R. and C. E. Casini. 1988. Revision of the neotropical genus *Nyttum* (Hemiptera: Coreidae). *Systematic Entomology* 13: 143–156.
- Dolling, W. R. and T. R. Yonke. 1976. The genus *Coriomeris* in North America. *Annals of the Entomological Society of America* 69: 1147–1152.
- Froeschner, R. C. 1963. The genus *Ceraleptus* Costa in the Western Hemisphere (Hemiptera: Coreidae). *Journal of the Kansas Entomological Society* 36: 109–113.
- . 1988. Family Coreidae Leach, 1815, pp. 69–92. In Henry, T. J. and R. C. Froeschner, eds., *Catalog of the Heteroptera, or True Bugs of Canada and the Continental United States*. E. J. Brill, New York. 958 pp.
- Gibson, E. H. and A. Holdridge. 1918a. Notes on North and Central American species of *Acanthocephala* Lap. (Fam. Coreidae: Heteroptera). *Canadian Entomologist* 50: 237–241.
- . 1918b. The genus *Narnia* Stål and a key to the genera of Anisoscelini Amyot and Serville (Coreidae: Heteroptera). *Psyche* 25: 1–4.
- Herring, J. L. 1980. A review of the cactus bugs of the genus *Chelinidea* with the description of a new species (Hemiptera: Coreidae). *Proceedings of the Entomological Society of Washington* 82: 237–251.
- Hossain, M. A. 1970. A revision of the genus *Mozena* Amyot and Serville (Hemiptera: Coreidae). Unpublished Master's Thesis, Texas A&M University, College Station.
- Kormilev, N. A. 1954. Notas sobre Coreidae neotropicales II; (Hemiptera) Merocorinae de la Argentina y países limítrofes. *Revista Ecuatoriana de Entomología y Parasitología* 2: 153–186.
- Leach, W. E. 1815. Hemiptera, p. 123. In Brewster's *Edinburgh Encyclopedia*. Volume 9. Edinburgh, Scotland. Pp. 57–192.
- O'Shea, R. 1980a. A generic revision of the Acanthocerini (Hemiptera: Coreidae: Coreinae). *Studies on Neotropical Fauna and Environment* 15: 57–80.

- . 1980b. A generic revision of the Nemato-podini (Heteroptera: Coreidae: Coreinae). *Studies on Neotropical Fauna and Environment* 15: 197–225.
- O'Shea, R. and C. W. Schaefer. 1978. The Mictini are not monophyletic (Hemiptera: Coreidae: Coreinae). *Annals of the Entomological Society of America* 71: 776–784.
- Osuna, E. 1984. Monografía de la tribu Anisosceldini (Hemiptera, Heteroptera, Coreidae) I. Revisión Genérica. *Boletín Entomológica Venezuela* 3: 77–148.
- Reed, E. C. 1898–1902. Sinopsis de los Hemipteros de Chile. Valparaíso. 107 pp.
- Ruckes, H. 1955. The genus *Chariesterus* de Laporte (Heteroptera, Coreidae). *American Museum Novitates* 1721: 1–16.
- Schmidt, E. 1911. *Deutsche Entomologische Zeitschrift* 1911: 565–571.
- Stål, C. 1859. Hemiptera. Species novas descripsit. *Kongliga Svenska Fregattens Eugenies Resa Omkring Jorden, III (Zoologi, Insekter)*: 219–298.
- . 1862. Synopsis Coreidum et Lygaeidum Suecia. Öfversigt af Kongliga Svenska Vetenskaps-Akadamiens Förhandlingar 19: 203–225.
- . 1864. Hemiptera nonnulla nova vel minus cognita. *Annales de la Société Entomologique de France, Series 4, 4*: 47–68.
- . 1867. Bidrag till Hemipterernas systematik. Öfversigt af Kongliga Svenska Vetenskaps-Akadamiens Förhandlingar 28: 491–560.
- . 1870. Enumeratio Hemipterorum. Bidrag till en företeckning öfver alla hittills kända Hemiptera, jemte systematiska meddelanden. Pt. 1. *Svenska Vetenskaps-Akadamiens Handlingar* 9(1): 1–232.
- . 1873. Enumeratio Hemipterorum. Bidrag till en företeckning öfver alla hittills kända Hemiptera, jemte systematiska meddelanden. Pt. 3. *Svenska Vetenskaps-Akadamiens Handlingar* 11(2): 1–163.
- Stein, J. P. E. F. 1860. Ueber einige Coreiden-Gattungen. *Berliner Entomologische Zeitschrift* 4: 246–256.
- Torre-Bueno, J. R. de la. 1941. A synopsis of the Hemiptera-Heteroptera of America north of Mexico. Part II. Families Coreidae, Alydidae, Corizidae, Neididae, Pyrrhocoridae and Thaumastotheriidae. *Entomologica Americana (New Series)* 21: 41–122.
- Whitehead, D. R. 1974. Variation and synonymy in *Hypsilonotus* (Heteroptera: Coreidae). *Journal of the Washington Academy of Sciences* 64: 223–233.
- Yonke, T. R. 1972. A new genus and two new species of neotropical Chariesterini (Hemiptera: Coreidae). *Proceedings of the Entomological Society of Washington* 74: 283–287.



Packauskas, Richard J. 1994. "Key to the subfamilies and tribes of the new world Coreidae (Hemiptera), with a checklist of published keys to genera and species." *Proceedings of the Entomological Society of Washington* 96, 44–53.

View This Item Online: <https://www.biodiversitylibrary.org/item/54711>

Permalink: <https://www.biodiversitylibrary.org/partpdf/54638>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Entomological Society of Washington

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.