
VIII. *An Illustration of the Species of Lycium which grow wild at the Cape of Good Hope. By Sir Charles Peter Thunberg, Knight of the Order of Wasa, Professor of Botany at Upsal, F.M.L.S.*

Read June 17, 1806.

LYCIUM ad illas plantas, quæ latius territorium occupant, quæ diversa climata ferunt, quæ in utroque hemisphærio telluris inventæ sunt, referendum erit. In Europa boreali et australi, in Siberia, in Africæ variis angulis inque America meridionali species hujus variæ læte crescunt et florent. Facile quidem genus hocce a congeneribus Pentandris distingui videtur corolla imprimis tubulosa fauce intus barbata, atque fructu baccato. Sed variare non raro solet staminibus quatuor, corolla et calyce quadrifidis. Præterea etiam *Serissa* ab hoc ipso genere divulsa nimis consimilis est, sic ut cum *Lycio* affinitate summa jungatur.

Cum vero mihi, sub meis in australi Africa itineribus plures *Lycii* species detegere, et ante detectas ulterius examinare licuerit, in animum induxi, hasce species fusius descriptas et novarum specierum icones pictas illustri Societati Linnæanæ offerre ejusque examini subjicere.

Serissa fætida imprimis fructu infero dispermo a *Lycio* distincta describitur; hic vero character valde dubius a me in *Flora Japonica* reputatus fuit, quare ego *Lycium japonicum* appellavi, Linnè junior *Lycium fætidum*, et Retzius *Lycium indicum*. Hæc tamen nota illust. Heretier satis sufficiens fuit

ad

ad novum genus, *Buchozia coprosmoides* dictum et D^{no} Loureiro ad *Dysodam* suam *fasciculatam* inde formandum, licet varie variant et staminum numerus et aliæ floris partes. Sunt itaque *Lycii* species septem, quæ Caput Bonæ Spei patriam suam agnoscunt, licet duæ earum, scilicet *Lycium afrum* et *barbarum* alibi quoque sponte crescant. *Lycii* species sunt vel spinescentes, vel inermes.

LYCIUM.

* *Spinosa.*

1. *L. barbarum* : foliis lanceolato-ellipticis, ramis laxis.
2. *L. afrum* : foliis lineari-ellipticis, ramis strictis, pedunculis foliis longioribus.
3. *L. rigidum* : foliis linearibus, ramis strictis, pedunculis foliis brevioribus.
4. *L. tetrandrum* : foliis ovatis obtusis, ramis strictis angulatis, floribus subsessilibus, corollis quadrifidis.
5. *L. cinereum* : foliis lanceolatis, ramis teretibus divaricatis, floribus subsessilibus.
6. *L. horridum* : foliis obovatis carnosissimis, ramis strictis, floribus subsessilibus.

** *inermis.*

7. *L. barbatum* : foliis ovatis, floribus subpaniculatis.

LYCIUM BARBARUM.

Lycium barbarum. Linn. *Syst. Veg.* xiv. p. 228. per Gmel.
Tom. i. P. 2. p. 1059.

Crescit in Carro inter Roggefelt et Bockefeld, florens Novembri, Decembri.

Caulis teres, cinereus, glaber, laxis, erectus.

Directions for placing the Plates of the Ninth Volume.

TAB. 1. Apion	- - - - -	to face page	80
2. Cancer floridus, &c.	}		
3. Cancer subterraneus, &c.			
4. Cancer Locusta, &c.			
5. Cancer rubricatus, &c.		- - - - -	114
6. Bulla Hydatis, &c.			
7. Doris longicornis, &c.			
8. Amphitrite Infundibulum			
9. Ursus indicus	- - - - -	-	116
10. Variolaria multipuncta & V. globulifera	- - - - -	-	137
11. Lecidea aromatica & L. atro-flava	- - - - -	-	140
12. Parmelia velata & P. carneo-lutea	- - - - -	-	143
13. Parmelia Clementi & P. Borreri	- - - - -	-	147
14. Lycium rigidum	- - - - -	-	153
15. Lycium tetrandrum	}		
16. Lycium cinereum		- - - - -	154
17. Lycium horridum			
18. Insect that destroys the Wheat	- - - - -	-	159
19. Didelphis cynocephala & D. ursina	- - - - -	-	174
20. Dimorpha grandiflora	- - - - -	-	180
21. Piper quadrangulare & P. bracteatum	- - - - -	-	202
22. Nauclea Gambir	- - - - -	-	218
23. Hookeria	- - - - -	-	277
24. } Notoclea	- - - - -	-	
25. }	- - - - -	-	294
26. Edwardsia chrysophylla, Splachnum squarrosum, &c.	}		
27. Neckera sphærocarpa, &c.		- - - - -	322
28. Bryum heterophyllum, &c.		- - - - -	

END OF THE NINTH VOLUME.

Directions for placing the Plates of the Ninth Volume.

TAB. 1.	Apion	-	-	-	-	-	to face page	80
2.	Cancer floridus, &c.	}						
3.	Cancer subterraneus, &c.							
4.	Cancer Locusta, &c.							
5.	Cancer rubricatus, &c.			-	-	-	-	114
6.	Bulla Hydatis, &c.							
7.	Doris longicornis, &c.							
8.	Amphitrite Infundibulum							
9.	Ursus indicus	-	-	-	-	-	-	116
10.	Variolaria multipuncta & V. globulifera	-	-	-	-	-	-	137
11.	Lecidea aromatica & L. atro-flava	-	-	-	-	-	-	140
12.	Parmelia velata & P. carneo-lutea	-	-	-	-	-	-	143
13.	Parmelia Clementi & P. Borreri	-	-	-	-	-	-	147
14.	Lycium rigidum	-	-	-	-	-	-	153
15.	Lycium tetrandrum	}						
16.	Lycium cinereum							154
17.	Lycium horridum							
18.	Insect that destroys the Wheat	-	-	-	-	-	-	159
19.	Didelphis cynocephala & D. ursina	-	-	-	-	-	-	174
20.	Dimorpha grandiflora	-	-	-	-	-	-	180
21.	Piper quadrangulare & P. bracteatum	-	-	-	-	-	-	202
22.	Nauclea Gambir	-	-	-	-	-	-	218
23.	Hookeria	-	-	-	-	-	-	277
24.	} Notoclea							
25.		-	-	-	-	-	-	294
26.	Edwardsia chrysophylla, Splachnum squarrosum, &c.	}						
27.	Neckera sphærocarpa, &c.							322
28.	Bryum heterophyllum, &c.							

END OF THE NINTH VOLUME.



Thunberg, Carl Peter. 1808. "An Illustration of the Species of Lycium which grow wild at the Cape of Good Hope." *Transactions of the Linnean Society of London* 9, 151–155. <https://doi.org/10.1111/j.1096-3642.1818.tb00333.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/13719>

DOI: <https://doi.org/10.1111/j.1096-3642.1818.tb00333.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/5579>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.