

EGG MORPHOLOGY OF *DRACONIA RUSINA* DRUCE FROM HONDURAS (LEPIDOPTERA: THYRIDIDAE)

STEVEN PASSOA AND BRUCE A. STEINLY

(SP) USDA/APHIS/PPQ, 8995 East Main Street, Bldg. 3, rm. 109, Reynoldsburg, Ohio 43068; (BAS) Illinois Natural History Survey, Center for Economic Entomology, 607 East Peabody Drive, Champaign, Illinois 61820.

Abstract.—The egg and some first instar morphological features of *Draconia rusina* are given based on specimens collected in Honduras. This is the first record of *D. rusina* from Honduras and the second illustration of a Neotropical thyridid egg.


Key Words: *Draconia rusina*, Thyrididae, egg, first instar, Honduras

The systematic position of the Thyrididae is unclear. Munroe (1972) and Whalley (1976) tentatively united thyridids with other Pyraloidea. Common (1990) and Minet (1983) elevated this group to its own superfamily named Thyridoidea. Passoa (1985) suggested pupal pilifers as a synapomorphy of Pyralidae, Hyblaeidae, Pterophoridae, and Thyrididae. Although the presence of pilifers is apomorphic (by virtue of the outgroup comparison method where *Monotrysis* is a sister group to *Ditrysis*), this character is homoplastic because pupal pilifers also occur in the unrelated Papilionoidea (Mosher 1916). Very little is known about immature thyridids (Whalley 1976). Therefore, the egg and some first instar larval characteristics of *Draconia rusina* Druce are described here to aid in comparing the morphology of pyraloid families for systematic and identification purposes.

Draconia rusina is a large moth previously recorded from Guatemala, Panama and Venezuela (Gaede 1936). Several specimens of *D. rusina* were collected in Comayagua, Honduras (Department of Comayagua) at a blacklight during April, 1979, and again in March and May, 1980. Approximately 20 eggs were obtained in 1979

(parental female number 78, S. Passoa coll.) using cheesecloth as an oviposition substrate. They were either dried or preserved in 80% ethanol. After several years, a few dried eggs were mounted on aluminum stubs with Elmer's glue and sputter coated twice at different angles with gold-palladium for further study. These were examined with an information Scientific instrument DS-130 scanning electron microscope and photographed with Polaroid type 55 positive/negative film. Structural measurements, expressed in microns or millimeters, were made directly from the scale line on the photographs after dividing by the magnification. Egg terminology follows Downey and Allyn (1981).

The egg (Figs. 1-4) is cream-colored and subcylindrical with nearly parallel truncate ends (length 1.29-1.34 mm, mean = 1.30 mm, $n = 4$; width 0.61-0.68 mm, mean = 0.65 mm, $n = 3$). The chorion sculpturing is composed of approximately 21 ridges parallel to the long axis of the egg (Fig. 1) that converge and terminate at the micropylar shoulder (Fig. 3). Ridges across the long axis are usually straight and separated, although a few are crossed or bifurcate (Figs. 1, 2; inter-ridge distance 17.02-29.79 μm ,



Figs. 1-4. Egg of *Draconia rusina* Druce. 1, lateral view (59 \times). 2, chorion ridges of the egg (235 \times). 3, oblique view of the micropyle (122 \times). 4, aeropyle apertures of chorion (580 \times). Scale lines marked in microns under each figure.

mean = 22.33 μm , $n = 10$). Aeropyle apertures occur on the chorion ridges (Figs. 2, 4) with one aperture per 21.28 to 28.40 μm (mean = 24.01 μm , $n = 55$). These apertures (Fig. 4) are irregularly shaped, with a width (opening parallel to the egg's long axis) of 6.38–19.15 μm (mean = 10.55 μm , $n = 12$) and a height of 10.34–20.69 μm (mean = 15.17 μm , $n = 5$). Micropylar end (Fig. 3) consists of a circular depression (diameter = 180.3 μm , $n = 1$) with a central raised tubercle (diameter = 77.87 μm , $n = 1$).

A review of published thyridid egg descriptions revealed some differences between taxa. Aniello (1980) mentioned a

hexagonal pattern of cells on the chorion of *Dysodia sica* Druce but did not include longitudinal ridges in her illustration. In contrast, *Thyris fenestrella* (Scopoli) (Sarlet 1964), *Aglaopus pyrrhata* (Walker) (Common 1990) and *Draconia rusina* eggs all have chorionic ridges. All four genera (*Aglaopus*, *Draconia*, *Dysodia* and *Thyris*) share a rounded, truncate apex which may be a general characteristic of thyridid eggs. Common (1990), Aniello (1980) and Sarlet (1964) mention that thyridids have upright eggs. The location of the micropyle in *D. rusina* is consistent with this type of egg.

Although our *Draconia rusina* eggs never

hatched, dissection revealed a fully developed larva within the egg shell. First instar larva of *D. rusina* have a short front, a bisetose SV group and apparently only one L seta on the thoracic segments, uniordinal crochets in a circle, and long setae on the anal shield. The latter character is shared with mature *Thyridopyralis* larvae which also have long setae on the anal shield. Aniello (1980) stated the crochets of first instar *Dysodia sica* larvae are in a biordinal circle, but this arrangement was not found in *Draconia rusina*.

ACKNOWLEDGMENTS

We thank the Center for Electron Microscopy at the University of Illinois for advice and use of facilities. The comments of T. Harrison, Dr. M. Berenbaum (University of Illinois) and Dr. G. Godfrey (Illinois Natural History Survey) on various drafts of this manuscript were greatly appreciated. J. Apperson helped with the figures. Dr. D. Habeck (University of Florida) allowed us access to his larval collection. This study could not have been completed without the identification of the parental female by Dr. D. Ferguson (USDA, Systematic Entomology Laboratory).

LITERATURE CITED

- Aniello, A. 1980. Life history of *Dysodia sica* (Lepidoptera: Thyrididae) in Panama. *Psyche* 87: 121-130.
- Common, I. F. B. 1990. *Moths of Australia*. E. J. Brill, Netherlands. 535 pp.
- Downey, J. C. and A. C. Allyn. 1981. Chorionic sculpturing in eggs of the Lycaenidae. Part 1. *Bulletin of the Allyn Museum* 61: 1-29.
- Gaede, M. 1936. Thyrididae. In Seitz, A. (ed.). *Macrolepidoptera of the World. American Bombyces and Sphinges*. A. Kernen, Stuttgart. 1327 pp.
- Minet, J. 1983. Étude morphologique et phylogénétique des organes tympaniques des Pyraloidea. I. Généralités et homologues (Lep. Glossata). *Annales de la Société Entomologique de France* 19: 175-207.
- Mosher, E. 1916. A classification of the Lepidoptera based on characters of the pupa. *Bulletin Illinois State Laboratory of Natural History* 12: 1-159.
- Munroe, E. 1972. Pyraloidea. Pyralidae (part). In Dominick, R. B., et al. (eds.). *The Moths of America North of Mexico*. Fascicle 13.1A. E. W. Classey, London. 134 pp.
- Passoa, S. 1985. Taxonomy of the larvae and pupae of economically important Pyralidae in Honduras. M.S. thesis. University of Florida, Gainesville. 486 pp.
- Sarlet, L. 1964. Iconographie des oeufs de Lépidoptères belges (Rhopalocera-Heterocera). Deuxième partie. Heterocera: Bombycides-Sphingides. *Mémoires de la Société Royale d'Entomologie de Belgique* 29: 1-172.
- Whalley, P. E. S. 1976. Tropical leaf moths. A monograph of the subfamily Striglininae (Lepidoptera: Thyrididae). *Bulletin of the British Museum of Natural History*. 194 pp.



Passoa, S and Steinly, B A. 1992. "Egg morphology of *Draconia rusina* Druce from Honduras (Lepidoptera: Thyrididae)." *Proceedings of the Entomological Society of Washington* 94, 556–558.

View This Item Online: <https://www.biodiversitylibrary.org/item/54937>

Permalink: <https://www.biodiversitylibrary.org/partpdf/55984>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Entomological Society of Washington

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.