XVI. *New species of Staphylinidae from Singapore.*
 Part IV (Conclusion). By MALCOLM CAMERON,
 M.B., R.N., F.E.S.

[Continued from Trans. Ent. Soc. Lond. 1920, p. 212.]

[Read November 17th, 1919.]

TABLE OF THE SUB-FAMILIES.*

1. Head in front of the eyes with a raised or thickened border, under which the antennae are inserted.	2.
— Head in front of the eyes without a raised or thickened border, the antennae either freely inserted on the front, or on or beneath the simple side margin of the head in front of the eyes.	4.
2. Posterior coxae transverse, not prominent.	3.
— Posterior coxae conical, prominent .	PAEDERINAE.
3. Gular sutures usually fused together for the greater part, often indistinct, at least fused at one point	OXYTELINAE.
— Gular sutures completely separated .	MEGALOPSINAE.
4. Antennae 12-jointed	ADINOPSINAE.
— Antennae 10- or 11-jointed.	5.
5. 1st joint of maxillary palpi elongate.	6.
— 1st joint of maxillary palpi short.	7.
6. Tarsal formula 5, 5, 5	STENINAE.
— Tarsal formula 4, 4, 4, or 5, 4, 4	EVAESTHETINAE.
7. Antennae inserted in a cup-shaped depression on the underside of the head .	PYGOSTENINAE.
— Antennae not so inserted.	8.
8. Antennae inserted on the front margin of the head	STAPHYLININAE.
— Antennae otherwise inserted.	9.

* The characters given in the tables do not necessarily apply to species not found in Singapore.

9. Antennae inserted freely on the lateral borders of the front; the elytral epipleurae not separated from the rest of the surface by a raised line; the elytra not extending beyond the metathorax **ALEOCHARINAE.**
- Antennae inserted under the simple (not raised or thickened) lateral borders of the front; the elytral epipleurae usually separated by a raised line from the rest of the surface; the elytra extending beyond the metathorax . . . **TACHYPORINAE.**

TABLE OF THE TRIBES.

I. Sub-family **OXYTELINAE.**

1. Anterior coxae globose, not prominent; the base of the abdomen keeled below **PIESTINI.**
- Anterior coxae conical, prominent; the base of the abdomen not keeled below. **2.**
2. Head with two ocelli **OMALIINI.**
- Head without ocelli. **3.**
3. Abdomen margined **OXYTELINI.**
- Abdomen not margined **OSORIINI.**

II. Sub-family **MEGALOPSINAE.**

1. Eyes large and prominent; 1st joint of maxillary palpi short; thorax sub-cylindrical; scutellum distinct; abdomen bordered. Tarsal formula 5, 5, 5 **MEGALOPSINI.**

III. Sub-family **STENINAE.**

1. Eyes large and prominent; maxillary palpi with the 1st joint elongate. Tarsal formula 5, 5, 5 **STENINI.**

IV. Sub-family **EVAESTHETINAE.**

1. Tarsal formula 4, 4, 5 **STENAESTHETINI.**
- Tarsal formula 4, 4, 4 **EVAESTHETINI.**

V. Sub-family **PAEDERINAE.**

1. 4th joint of maxillary palpi large **PINOPHILINI.**
- 4th joint of maxillary palpi small **PAEDERINI.**

VI. Sub-family **STAPHYLININAE.**

1. Anterior portion of the prosternum separated by a suture from the posterior portion. Antennae at the base usually nearer to each other than to the eyes, at least not farther apart. **XANTHOLININI.**
- Anterior portion of the prosternum not separated by a suture from the posterior portion. **2.**
2. Anterior angles of the thorax extended considerably beyond the anterior angles of the prosternum. Under side of the head with a longitudinal raised line at least behind . . . **QUEDIINI.**
- Anterior angles of the thorax not or scarcely extended beyond the anterior angles of the prosternum. Under side of the head without longitudinal raised line . . . **STAPHYLININI.**

VII. Sub-family **PYGOSTENINAE.**

One genus : **Delibius** Fam. (*q. v.*).

VIII. Sub-family **TACHYPORINAE.**

1. Sides of the thorax explanate; sculpture of the head and thorax coarse and rugose . . . **MEGARTHROPSINI.**
- Sides of the thorax not explanate; sculpture of the head and thorax fine or very fine, not rugose . . . **TACHYPORINI.**

IX. Sub-family **ADINOPSINAE.**

One genus : **Adinopsis**, n. (*q. v.*).

X. Sub-family **ALEOCHARINAE.**

1. Head more or less produced in front. **2.**
- Head not produced in front. **4.**
2. Tarsal formula 4, 4, 4 . . . **DIGLOTTINI.**
- Tarsal formula 4, 5, 5, or 4, 4, 5. **3.**
3. Tarsal formula 4, 5, 5 . . . **PRONOMAEINI.**
- Tarsal formula 4, 4, 5 . . . **MYLLAENINI.**
4. Antennae 10-jointed, tarsi 4-jointed . **OLIGOTINI.**
- Antennae 11-jointed. **5.**
5. Tarsal formula 5, 5, 5 . . . **ALEOCHARINI.**

- Tarsal formula otherwise 5.
- 6. Tarsal formula 4, 4, 5 BOLITOCARINI.
- Tarsal formula, 4, 5, 5, 4, 4, 4, or 3, 4, 4. MYRMEDONINI.

TABLE OF THE GENERA.

PIESTINI.

- 1. Abdomen bordered **Eleusis** Cast.
- Abdomen not bordered. 2.
- 2. Anterior tibiae serrated externally 3.
- Anterior tibiae not serrated. 5.
- 3. Anterior coxae separated **Leptochirus** Germ.
- Anterior coxae contiguous 4.
- 4. Mandibles much produced, their length about twice that of the head; 3rd joint of the maxillary palpi elongate, as long, or nearly as long, as the last **Borolinus** Bernh.
- Mandibles not much produced, their length at most a little longer than the head; 3rd joint of maxillary palpi broader than long, much shorter than the last **Priochirus** Shp.
- 5. Anterior tibiae with two small spines near the apex on the external border **Ancaeus** Fauv.
- Anterior tibiae without spines on the external border. 6.
- 6. Abdomen obliquely striolate; posterior angles of the thorax somewhat prominent **Holusus** Motsch.
- Abdomen not striolate, normally punctured; posterior angles of the thorax not prominent **Lispinus** Er.

OMALIINI.

- 1. Labrum emarginate anteriorly; mesosternum not keeled; last joint of the maxillary palpi slender, distinctly smaller than the 3rd **Phloeonomus** Heer.

OXYTELINI.

- 1. Anterior and middle tibiae spinose externally. 2.
- Anterior and middle tibiae not spinose externally. 3.

2. Anterior tibiae with a double row of spines; species cylindrical with strongly geniculate antennae . . . **Bledius** Mannh.
- Anterior tibiae with a single row of spines; species rather depressed, antennae not or scarcely geniculate. Thorax more or less tri-sulcate (except in *thoracicus*); intermediate coxae approximate . . . **Oxytelus** Er.
3. Scutellum visible. Elytra without epipleurae, the postero-internal angles separately rounded so that a small triangular space is apparent at the suture . . . **Thinobius** Kiesw.
- Scutellum concealed. Elytra with distinct epipleurae, the postero-internal angles not separately rounded . . . **Trogophloeus** Mannh.

OSORIINI.

1. Tibiae spinose. 2.
- Tibiae not spinose. 3.
2. Antennae geniculate; anterior tibiae dentate-spinose; last joint of the tarsi slender, not tumid . . . **Osorius** Latr.
- Antennae not geniculate; anterior tibiae simply spinose; last joint of the tarsi tumid . . . **Mimogonus** Fauv.
3. Thorax strongly contracted at the base; 4th joint of maxillary palpi subulate . . . **Paragonus** Fauv.
- Thorax not or scarcely contracted at the base; 4th joint of maxillary palpi not subulate . . . **Holotrochus** Er.

MEGALOPSINI.

1. Antennae 11-jointed; tarsal formula 5, 5, 5. Form oblong, stout. Tibiae simple . . . **Megalops** Er.

STENINI.

1. Eyes very large, occupying nearly the whole of the side of the head; 4th tarsal joint simple or bilobed; apex of abdomen without styles . . . **Stenus** Latr.

STENAESTHETINI.

1. Antennae very slender; abdomen except for the 1st (visible) and 5th segments immarginate. Tarsal formula 4, 4, 5. Antepenultimate joint simple; sculpture umbilicate on head and thorax . **Stenaesthetus** Shp.

EVAESTHETINI.

1. Tarsi 4-jointed; head deeply impressed on either side of the front; abdomen distinctly margined. Head and thorax very smooth and shining. . **Edaphus** J. Lec.

PINOPHILINI.

1. Abdomen bordered.	2.	
— Abdomen not bordered.	3.	
2. Labrum simple; last joint of the maxillary palpi narrow and elongate . . .		Pinophilus Gr.
— Labrum bilobed; last joint of the maxillary palpi securiform . . .		Neopinophilus Cam.
3. Sculpture of abdomen strongly imbricate; last joint of maxillary palpi securiform; terminal joint of antennae of normal length		Palaminus Er.
— Sculpture of abdomen not imbricate; last joint of maxillary palpi slender, fusiform; terminal joint of antennae very elongate, forming nearly half the length of the organ		Eucirrus Fauv.

PAEDERINI.

1. Antennae not geniculate.	2.
— Antennae strongly geniculate.	11.
2. 4th tarsal joint dilated, the distal margin more or less emarginate, the 5th joint articulating on its dorsal surface near the base, giving the appearance when viewed from above of the 4th joint being bilobed.*	3.

* All authors speak of this joint being bilobed. This, however, is incorrect. When examined as a microscopical preparation the structure is found to be as above described.

- 4th tarsal joint simple, not presenting a
bilobed appearance. 7.
- 3. 4th joint of maxillary palpi very short,
broad and obtuse; anterior tarsi dilated **Paederus** F.
- 4th joint of maxillary palpi minute,
subulate. 4.
- 4. Labrum bi-dentate or slightly emarginate
in the middle of the anterior border. 5.
- Labrum with 5 or 6 distinct teeth; head
with simple puncturation; elytra
strongly punctured, more or less in
rows **Psilotrachelus** Kr.
- 5. Labrum bidentate. 6.
- Labrum emarginate **Acanthoglossa** Kr.
- 6. Thorax elongate; abdomen parallel;
anal styles distinct **Astenus** Steph.
- Thorax shorter and broader; abdomen
somewhat contracted at the base; anal
styles wanting. **Stilicopsis** Sachse.
- 7. Antennae with the first two joints stout,
the following slender **Thinocharis** Kr.
- Antennae of normal build. 8.
- 8. Neck slender or very slender. 9.
- Neck broad **Medon**, Steph.
- 9. Labrum without teeth; 1st joint of the
antennae sulcate from apex nearly to
the base **Parascopaeus** Cam.
- Labrum toothed; 1st joint of the
antennae not sulcate. 10.
- 10. 1st joint of the posterior tarsi elongate,
distinctly longer than the last; tongue
bifid **Stilicus** Latr.
- 1st joint of the posterior tarsi short, not
longer than the last; tongue trifid . **Scopaeus** Er.
- 11. Tibiae spinose; 4th joint of the maxillary
palpi distinct, conical; tongue bi-
lobed **Cryptobium** Mannh.
- Tibiae setose; 4th joint of the maxillary
palpi very small, obtuse, scarcely
visible; tongue simple **Calliderma** Motsch.

XANTHOLININI.

- 1. Tibiae not spinose **Somoleptus** Shp.
- Tibiae spinose. 2.

2. Antennae geniculate.	3.
— Antennae not geniculate	Diochus Er.
3. 3rd joint of the maxillary palpi longer than the 2nd.	4.
— 3rd joint of the maxillary palpi not longer than the 2nd.	5.
4. Gular sutures obsolete; intermediate coxae narrowly separated	Oligolinus Cas.
— Gular sutures distinct; intermediate coxae widely separated	Leptacinus Er.
5. Labrum broadly emarginate	Eulissus Mannh.
— Labrum with several short blunt teeth	Thyrecephalus Guer.

STAPHYLININI.

1. Tarsal formula 4, 4, 5; small depressed species	Holisomimus Cam.
— Tarsal formula 5, 5, 5.	2.
2. Anterior and posterior femora furnished below with two rows of fine spines	Belonuchus Nordm.
— Anterior and posterior femora not furnished below with two rows of spines, at most (in some species of Philonthus) with a few spines towards the apex.	3.
3. 2nd joint of the antennae thickened, much thicker than the 3rd	Actobius Fauv.
— 2nd joint of the antennae not thickened, not or scarcely thicker than the 3rd.	4.
4. Lateral setae of the thorax placed on the side margin itself or very near thereto.	5.
— Lateral setae of the thorax distant from the side margin.	6.
5. Anterior tarsi dilated in both sexes; mesosternum without a transverse impressed line	Orthidus Rey.
— Anterior tarsi at most dilated in the ♂; mesosternum with a transverse impressed line	Philonthus Curt.
6. Last joint of the maxillary palpi nearly twice as long as the preceding; mesosternum broadly rounded behind	Hesperus Fauv.
— Last joint of the maxillary palpi scarcely longer than the preceding; mesosternum pointed	Cafius Steph.

QUEDIINI.

1. Antennae strongly geniculate . **Acylophorus Nordm.**

PYGOSTENINI.

1. Form navicular. Tarsal formula 4, 4, 5 :
legs long; intermediate coxae widely
separated; antennae longer than the
head and thorax; scutellum very large **Delibius Fauv.**

MEGARTHROPSINI.

1. Sides of the thorax explanate; sculpture
of head and thorax coarse and rugose **Megarthritis Cam.**

TACHYPORINI.

1. Tarsal formula 5, 4, 4 . . . **Atanygnathus Jacobson.**
— Tarsal formula 5, 5, 5. 2.
2. Last joint of the maxillary palpi small,
subulate; abdomen not or scarcely
margined . . . **Conosoma Kr.**
— Last joint of the maxillary palpi not
subulate, often longer than the 3rd. 3.
3. Species very convex, strongly contractile
and retractile; posterior angles of the
thorax effaced; tibiae not spinose . **Mimocyptus Cam.**
— Species much less convex, not contractile;
posterior angles of the thorax distinct;
tibiae more or less spinose. 4.
4. 1st joint of the posterior tarsi as long as
the three following together . **Leucoparyphus Kr.**
— 1st joint of posterior tarsi shorter than
the three following together. 5.
5. 1st visible dorsal segment at the base on
either side of the middle line with a
short transversely extended tomentose
patch. Species larger . . . **Tachinomorphus Kr.**
— 1st visible dorsal segment without tomen-
tose patches. Species small or very
small . . . **Coproporus Kr.**

GYMNUSINI.

1. Head deflexed, concealed; the sides and
anterior margin of the thorax forming
a semi-circle, posterior angles promin-
ent, pointed. Legs slender, all the

tarsi 5-jointed, the 1st joint of the posterior pair as long as the three following together **Leucocraspedum** Kr.

MYLLAENINI.

1. Tarsal formula 4, 4, 5. Labial palpi slender, styliform, obscurely 3-jointed; maxillary palpi very long and slender **Myllaena** Er.

PRONOMAEINI.

1. Tarsal formula 4, 5, 5. Labial palpi very long, styliform, not jointed. Maxillary palpi very long and slender, the 4th joint very short, subulate . . . **Pronomaea** Er.

DIGLOTTINI.

1. Tarsal formula 4, 4, 4. Labial palpi very long and slender, obscurely 3-jointed. Maxillary palpi very long and slender, the last joint very small, subulate . **Diglotta** Champ.

OLIGOTINI.

1. Tarsal formula 4, 4, 4. Antennae 10-jointed. Labial palpi obscurely 3-jointed **Oligota** Mannh.

BOLITOCCHARINI

1. Mesosternal process narrow and pointed, the intermediate coxae contiguous or but little separated.	2.
— Mesosternal process broader, apex rounded, the intermediate coxae distant.	14.
2. Labial palpi 2-jointed.	3.
— Labial palpi 3-jointed, the 2nd joint much shorter than the 1st and 3rd. Right mandible with a distinct tooth at the middle of the inner border. Temples bordered below. (See also Heterota , 10.)	Pseudatheta , Cam.
3. Middle and posterior tibiae with at least one long seta.	4.
— Middle and posterior tibiae without long setae, at most with a weak seta.	5.

4. Shining convex species with strongly pointed abdomen. Labial palpi elongate, the 1st joint not constricted at the inner border, and twice as long as the 2nd. Mandibles simple. Facies somewhat resembling **Tachyporus** . **Hetairotermes**, n. n.*
- Rather depressed, dull parallel species, the 1st joint of the labial palpi constricted at the inner border. Right mandible with a tooth . . . **Homalota** Mannerh.
5. Temples not bordered below. 6.
- Temples bordered below. 8.
6. Elytra sinuate. Tongue narrow, elongate, bifid. 1st joint of the labial palpi constricted at the inner border beyond the middle; the 2nd elongate, shorter than the 1st. Head narrowed and rounded behind the eyes . . . **Neomalota** Cam.
- Elytra not sinuate. Tongue bifid or emarginate. 1st joint of labial palpi not constricted at the inner border. Head quadrate. 7.
7. Tongue elongate, bifid. Terminal joint of the tarsi not dilated; 8th dorsal segment of the abdomen toothed at the posterior border. Habitat under bark **Thectura** Thoms.
- Tongue obviate, emarginate anteriorly. Terminal joint of the tarsi dilated; 8th dorsal segment of the abdomen not toothed at the posterior border. Habitat maritime . . . **Paractocharis** Cam.
8. Head and thorax very finely, very sparingly and obsoletely punctured. Depressed, shining, parallel species . **Lampromalota** Cam.
- Head and thorax distinctly and closely punctured. 9.
9. Tongue simple. 10.
- Tongue elongate, more or less divided or emarginate. 11.
10. Tongue short and broad. 1st joint of labial palpi not constricted at inner border. Elytra not sinuate . . **Placusa** Er.

* *Termophila* Lea, nom. praeoc.

- Tongue elongate. Labial palpi obscurely 3-jointed **Heterota** Rey.
- 11. 1st joint of labial palpi not constricted at the inner border. 12.
- 1st joint of labial palpi constricted at the inner border **Chledophila** Cam.
- 12. 2nd joint of the labial palpi distinctly shorter than the 1st; tongue narrowed at the base, widened towards the apex. Facies of **Homalota** **Mimomalota** Cam.
- 2nd joint of the labial palpi as long or longer than the 1st. 13.
- 13. Tongue very narrow, elongate, parallel. Facies of **Placusa** **Pseudoplacusa** Cam.
- Tongue broader, narrowed at the base, widened towards the apex. Facies of **Neosilusa** **Deralia** Cam.
- 14. Mesosternum finely carinate. 15.
- Mesosternum not carinate. 17.
- 15. Elytra distinctly sinuate, the sides with 3 long and strong setae. Tongue broad with rounded sides, narrowed at the base, nearly bilobed. Labial palpi 3-jointed, the 3rd joint minute, subulate **Adelarthra** Cam.
- Elytra distinctly sinuate, the sides without long setae. Labial palpi 2-jointed. 16.
- 16. Labial palpi not styliform, the 2nd joint as long as, but narrower than the 1st. Tongue bifid nearly to the base. Right mandible with a small tooth. Facies of **Pseudoligota** **Sternotropa** Cam.
- Labial palpi styliform, the 2nd joint longer than the 1st. Tongue narrow, bifid for half its length. Right mandible with a small tooth. Facies of **Neosilusa** **Prosilusa** Cam.
- 17. Tongue simple. 18.
- Tongue bifid. 20.
- 18. Tongue short and broad, halberd-shaped, labial palpi 2-jointed, the 1st joint short and broad, the antero-external angle prominent and with a strong seta, the inner border constricted before

the apex; 2nd joint as long as, but much narrower than the 1st. Facies

of **Acrotona** **Pelekoglossa** Cam.

— Tongue long or moderately long, not halberd-shaped.

19.

19. Labial palpi with the 2nd joint as long as the 1st, styliform

Neosilusa Cam.

— Labial palpi with the 2nd joint much shorter than the 1st, not styliform .

Gyrophaena Mannh.

20. Labial palpi distinctly 2-jointed, the 1st joint without constriction at the inner border; tongue small and triangular, split at the apex; inner lobe of the maxilla truncate and finely pectinate at apex; temples not bordered below.

Facies of **Oligota**

Pseudoligota Cam.

— Labial palpi obscurely 3-jointed, the 1st joint having a constriction at the inner margin from which a more or less apparent oblique suture passes forwards and outwards; tongue elongate and bifid; inner lobe of maxilla pointed and narrow, the inner margin strongly pectinate; temples bordered below.

21.

21. Last joint of the labial palpi as long as the 1st (true) joint; outer lobe of maxilla simply ciliate at apex. Facies of **Neosilusa**

Ousilusa Cam.

— Last joint of the labial palpi shorter than the 1st (true) joint; outer lobe of maxilla ciliate plumose at apex.

22.

22. Elytra distinctly sinuated at the postero-external angle; middle and posterior tibiae without long setae. Facies somewhat resembling **Gyrophaena** .

Pseudophaena Cam.

— Elytra not sinuated at the postero-external angle; middle and posterior tibiae with a long seta. Facies somewhat resembling **Homalota** .

Caenonica Kr.

MYRMEDONIINI.

1. Tarsal formula 4, 5, 5.

2.

— Tarsal formula 4, 4, 4, or 3, 4, 4.

13.

2. Maxillary socket wide and deep, extending to the level of the posterior border of the eye or beyond it. Mesosternum broadly rounded or truncate behind, extending but little between the intermediate coxae; metasternum produced, bordered, not meeting the mesosternum. 3.
- Maxillary socket neither wide nor deep, not extending to the level of the posterior border of the eye. 4.
3. 1st joint of the posterior tarsi twice as long as the 2nd **Zyras** Steph.
- 1st joint of the posterior tarsi but little longer than the 2nd **Myrmedonota** Cam.
4. Labial palpi distinctly 2-jointed. 5.
- Labial palpi 3-jointed, sometimes (**Paratheta** and **Fenyesia**) obscurely so. 8.
5. Sculpture coarse and rugose **Schistogenia** Kr.
- Sculpture fine, not rugose. 6.
6. Labial palpi almost styliform, the 2nd joint half as long as the 1st. Tongue narrow and elongate, a little widened anteriorly, the apex with a small emargination **Exatheta** Cam.
- Labial palpi normal. Tongue split into two lobes. 7.
7. Tongue rather long, split nearly to the base into two narrow diverging lobes. Mesosternal process gradually narrowed, the apex bluntly pointed, the intermediate coxae moderately separated. **Mimatheta** Cam.
- Tongue rather short and broad, split to the middle into two diverging teat-shaped lobes. Mesosternal process narrow, sharply pointed, the intermediate coxae narrowly separated . . . **Mimacrotona** Cam.
8. Head with very narrow, distinctly exposed neck; the base of the head not at all concealed by the thorax. 9.
- Head with broad neck; the base of the head more or less concealed by the thorax. 10.

9. 1st joint of the posterior tarsi a little longer than the 2nd; thorax obtusely angled before the middle, the sides strongly contracted and sinuate posteriorly **Amaurodera** Fauv.
- 1st joint of the posterior tarsi as long as the three following together; thorax not obtusely angled before the middle . **Falagria** Mann.
10. Labial palpi distinctly 3-jointed. 11.
- Labial palpi obscurely 3-jointed. 12.
11. Tongue short and broad, broadest at the base, emarginate in front **Pelioptera** Kr.
- Tongue longer, narrow at the base, more or less bifid **Atheta** Thoms.
12. 1st joint of the posterior tarsi elongate, about twice as long as the 2nd; thorax strongly transverse, convex, the posterior angles acute and produced. Mesosternal process narrow and pointed, the intermediate coxae very narrowly separated **Fenyesia** Cam.
- 1st joint of the posterior tarsi short, sub-equal to the 2nd; thorax much less transverse, the posterior angles not acute or produced. Mesosternal process broad, truncate posteriorly, keeled longitudinally in the middle line, the intermediate coxae widely separated **Paratheta** Cam.
13. Tarsal formula 4, 4, 4. Labial palpi 2-jointed **Termitochus** Silvestri.
- Tarsal formula 3, 4, 4. Labial palpi 3-jointed: facies approaching **Falagria** **Eusteniamorpha** Cam.

ALEOCHARINI.

1. Maxillary palpi 5-, labial palpi 4-jointed. 2.
- Maxillary palpi 4-, labial palpi 2-jointed. **Myrmedonella** Cam.
2. Anterior and middle tibiae spinose . **Aleochara**, Gr.
- Anterior and middle tibiae not spinose. 3.
3. Elytra not sinuate at the postero-external angle. Tongue moderately broad, split to the middle into two narrow lobes. 4.

- Elytra strongly sinuate at the postero-external angle. Tongue narrow and elongate, the apex only bifid . . . **Hoplandria** Kr.*
- 4. Thorax with four large punctures placed quadrately on the disc. Species robust, oblong **Tetrasticta** Kr.
- Thorax without four quadrately placed punctures on the disc. Species more elongate **Paraleochara** Cam.

TABLES OF THE SPECIES.

Eleusis Cast.

- 1. Species black, the elytra testaceous with apical margin narrowly black . . . *humilis* Er.
- Species in great part testaceous or reddish-testaceous. 2.
- 2. Elytra very narrowly infusate posteriorly. 3.
- Elytra broadly infusate posteriorly . . . *lunigera* Fauv.
- 3. Species smaller (1.6 mm.); head more or less pitchy *fusciceps* Kr.
- Species larger (3 mm.); head reddish testaceous *kraatzi* Fauv.

Leptochirus Germ.

- 1. Front of head without impressed line; prosternal process much widened behind (Sub-gen. *Strongylochirus* Bernh.). Clypeus declivous, separated from the front by a transverse line and from the sides by a curved impressed line *laevis* cast.

Borolinus Bernh.

- 1. Red, the elytra and apical part of the abdomen more or less black. Length 8 to 10 mm. (*minutus* Cast.) v. *cruentus* Fauv.

* Stated by Kraatz (Linn. Ent. 1857, p. 4) to have the anterior tarsi 4-jointed; this is incorrect. The minute accessory joint of the maxillary and labial palpi was also overlooked by this author.

Priochirus Sharp.

1. Front of the head with a tooth in the middle (Sub-gen. *Triacanthus*) and one on either side, all of about equal length; sides of the thorax uniformly punctured; femora pitchy-black. Length 7 to 8 mm. . (*tridens* Motsch.) v. *insularis* Bernh.
- Front of the head with a deep excision in the middle line (Sub-gen. *Cephalomerus* Bernh.). 2.
2. Frontal excision deeper; lateral teeth separated by a broader and deeper excision from the central ones; thorax much more transverse *hoplites* Fauv.
- Frontal excision less deep: lateral teeth separated by a smaller and shallower excision from the central ones; thorax less transverse *pygmaeus* Kr.

Ancaeus Fauv.

1. Head, thorax and elytra with very indistinct ground sculpture; species testaceous, larger, more shining; thorax much more transverse . . . *exiguus* Er.
- Head, thorax and elytra with very distinct longitudinally strigose ground sculpture; species usually pitchy-brown, smaller, less shining; thorax much less transverse (1) *singularis* Cam.

Holosus Motsch.

1. Elytra without elevated lines or keels; facies of *Tachyporus* . . . *tachyporiformis* Motsch.
- Elytra each with two elevated lines or keels *plicatus* Bernh.

Lispinus Er.

1. Posterior angles of the thorax scarcely impressed. 2.
- Posterior angles of the thorax distinctly impressed. 3.
2. Size larger (3.5 mm.); abdomen distinctly but sparingly punctured; antennae ferruginous *coarcticollis* Kr.

- Size smaller (1.7 mm.); abdomen impunctate, antennae testaceous . (3) *minutus* Cam.
- 3. Disc of each elytron with two rows of large setiferous punctures . . . (2) *setosus* Cam.
- Disc of the elytra without rows of large setiferous punctures. 4.
- 4. Species shining; ground sculpture of the fore-parts very indistinct; disc of thorax distinctly and not sparingly punctured (2a) *sharpi* Cam.
- Species with greasy lustre only; ground sculpture of the fore-parts very distinct, coriaceous; disc of thorax sparingly punctured. 5.
- 5. Lateral impression of the thorax deeper, impunctate; antennae shorter, the penultimate joints more transverse . *impressicollis* Motsch.
- Lateral impressions of the thorax shallower, punctured; antennae longer, the penultimate joints less transverse *tenuicornis* Kr.

Phloeonomus Heer.

- 1. Head and thorax opaque, elytra black or pitchy-black, scarcely shining . *obscurus* Kr.
- Head and thorax shining, elytra with the disc testaceous, shining . . . *discalis* Cam.

Trogophloeus Mannerh.

- 1. 5th to the 7th joints of the antennae longer than broad (*Trogophloeus* s.str.). 2.
- 5th to the 7th joints of the antennae not longer than broad (Sub-gen. *Taenosoma*). 3.
- 2. Eyes very large, occupying nearly the whole of the side of the head; abdomen very thickly covered with fine grey pubescence; species duller, thorax much less strongly contracted at the base (4) *orientalis* Cam.
- Eyes moderate, temples longer; abdomen much less thickly pubescent; species more shining, thorax strongly contracted at the base (5) *silvestris* Cam.

- 3. Species in great part reddish-testaceous. 4.
- Species entirely or in great part dark. 5.
- 4. Size larger 1.75 mm., shining, the fore-parts punctured, not shagreened; the diameter of the eyes equal to the length of the temples, and moderately prominent (8) *littoralis* Cam.
- Size smaller 1.4 mm., with greasy lustre only, the fore-parts shagreened, not punctured; the eyes small and flat, their diameter much less than the length of the temples (9) *rufotestaceus* Cam.
- 5. Species smaller, black, nearly opaque, head and thorax shagreened, not punctured (6) *halophiloides* Cam.
- Species larger, castaneous, shining, the fore-parts distinctly punctured, not shagreened (7) *lucens* Cam.

Aploderus Steph.

- 1. Rufo-testaceous, shining, finely and sparingly punctured (10) *testaceus* Cam.

Oxytelus Grav.

- 1. 1st joint of the antennae elongate, constricted before the apex; eyes large, occupying nearly the whole side of the head (Sub-gen. *Caccoporus* Thoms.). 2.
- 1st joint of the antennae only moderately long, gradually thickened and not constricted before the apex. 4.
- 2. Head black, nearly opaque, not or very obsoletely punctured, densely coriaceous; size larger (4.5 to 5 mm.). 3.
- Head reddish-testaceous, shining, distinctly punctured posteriorly in front, coriaceous; size smaller (2.75 mm.).
- ♂: 7th ventral segment broadly and feebly emarginate, on each side with a small spine; 6th ventral segment with a small tubercle at posterior margin on each side of the middle line *ferrugineus* Kr.

3. Head, thorax and elytra impunctate, densely coriaceous; species less shining. ♂: 7th ventral segment with a deep narrow excision on either side, bounding a central quadrate lobe, this furnished with a tubercle in the middle and with the posterior border a little produced and elevated in the middle line; 6th ventral segment a little produced in the middle and truncate *bengalensis* Er.
- Head, thorax and elytra distinctly punctured, thorax and elytra not coriaceous; species more shining. ♂: characters of the preceding, but with the posterior border of the central lobe of the 7th segment broadly emarginate on either side, and the central produced point smaller and sharper *nigriceps* Kr.
4. Species black or blackish with dark antennae at most, lighter at the base. 5.
- Species in great part reddish-testaceous or ferruginous with reddish antennae. 8.
5. Head entirely shining, with large and distinct punctures posteriorly; thorax shining, strongly rugose and strigose; size larger (2.6 mm.) *exasperatus* Kr.
- Head impunctate, thorax strigose, not rugose; size smaller (1.5–2 mm.). 6.
6. Sides of the head with a distinct raised line internal to the eyes, extending from the base and continuous with the frontal margin anteriorly; species entirely opaque except the abdomen . *latiusculus* Kr.
- Sides of the head without raised line internal to the eyes 7.
7. Thoracic ridges and front of the head shining, the latter scarcely perceptibly strigose; species somewhat shining. ♂: 7th ventral segment with a small tubercle on either side near the middle and in front of the posterior margin; 6th ventral segment with a moderately

- broad, shallow emargination of the posterior border *pygmaeus* Kr.
- Thoracic ridges completely dull, front of the head scarcely shining, distinctly longitudinally strigose; species opaque (13) *obscurus* Cam.
8. Elytra simply punctured, not strigose, size smaller (1.4 mm.) (12) *frugicola* Cam.
- Elytra distinctly strigose, size larger. 9.
9. Head (except clypeus) and thorax opaque, species dark ferruginous *raffrayi* Fauv.
- Head and thorax shining. 10.
10. Thorax distinctly tri-sulcate. 11.
- Thorax with feeble median sulcus only, the lateral wanting. Head and thorax very finely and sparingly punctured, not rugose *thoracicus* Motsch.
11. Head dark brown or black *nitidifrons* Woll.
- Head testaceous or ferruginous. 12.
12. Penultimate joints of the antennae distinctly transverse. Head in part with distinct coriaceous ground-sculpture. ♂: 7th ventral segment truncate . *kraatzii* (*pulcher* Kr.).
- Penultimate joints of the antennae scarcely transverse. Head scarcely at all coriaceous. ♂: 7th ventral segment slightly emarginate on either side . *granadillae* Cam.

Bledius Mannerh.

1. Labrum not emarginate; thorax in the ♂ with a long spine directed forwards (*Bledius* s.str.); species shining, black; antennae and legs testaceous *bellicosus* Fauv.

Thinobius Kies.

1. Head smaller than the thorax, eyes moderate (*Thinobius* s.str.). Black, elytra, antennae and legs testaceous . *marinus* Cam.

Paragonus Fauv.

1. Pitchy-black or pitchy-brown; thorax obtusely angulate behind the middle, the sides from thence anteriorly strongly and obliquely contracted, posteriorly strongly arcuate and contracted. Length 2.75 to 3.3 mm. . *heteroceros* Fauv.

Mimogonus Fauv.

1. Cylindrical, black, shining; elytra castaneous; fore-parts with rather large superficial punctures. Length 2.75 mm. *fumator* Fauv.

Holotrochus Er.

1. Cylindrical, black, shining; head and thorax distinctly, elytra obsoletely punctured. Length 3-3.5 mm. . (14) *nitidus* Cam.

Osorius Latr.

1. Black, shining; head closely and densely longitudinally strigose between the eyes, the sides in front with fine asperate punctures; thorax finely and not very closely punctured. Length 7 mm. . *rugifrons* Er.

Stenus Latr.

1. 4th tarsal joint bilobed; abdomen not bordered (*Hypostenus* Rey). Black, shining, each elytron with a small round orange spot; antennae elongate, the first two joints testaceous, the following reddish, the 9th to the 11th black; palpi and legs testaceous. Length 5 mm. ? *bivulneratus* Motsch.
- Elytra without orange spot. 2.
2. Species in great part brown; head with smooth, elevated, longitudinal impunctate space in the middle; antennae elongate, the first two joints testaceous; legs testaceous, the apex of the femora and base of the tibiae infusate (16) *castaneus* Cam.
- Species black. 3.
3. Head between the eyes flat; species smaller, less shining, less coarsely punctured, antennae shorter, with the 1st joint pitchy. Length 3.3 mm. . ? *monomeros* Fauv.
- Head between the eyes concave; species larger, shining, more coarsely punctured, antennae longer, with the 1st joint testaceous. Length 3.75 mm. (15) *fortepunctatus* Cam.

Stenaesthetus Sharp.

1. Brown, sub-opaque, fore-parts strongly and closely punctured, abdomen very finely and closely punctured; antennae and legs testaceous *sunioides* Sharp.

Edaphus Le Conte.

1. Rufous; head deeply sulcate between and before the eyes on each side; 2nd abdominal segment carinate, the 3rd bi-fossulate in the middle, 4th and 5th more broadly impressed, the former with a small triangular elevation. Length 1.5 mm. *dentiventris* Fauv.

Pinophilus Grav.

1. Head with smooth, triangular, shining impunctate space in front; abdomen uniformly grey, pubescent, the centre of the segments not more shining than the sides. Length 5.75-6.5 mm. (16a) *orientalis* Cam.
- Head without smooth impunctate space in front; abdomen with the sides of the segments, and especially the 6th, clothed with long golden pubescence, the centre of the segments much more shining than the sides. Length 14 mm. *borneensis* Fauv.

Neopinophilus Cam.

1. Rufous, shining; head very finely and sparingly punctured; elytra one-third shorter than the thorax. Length 7.8 mm. (17) *notabilis* Cam.

Eucirrus Fauv.

1. Rufo-ferruginous, shining; head elongate with coarse umbilicate puncturation; thorax elongate; subserially punctured; elytra transverse, shorter than the thorax, strongly and roughly punctured. Length 6.5 mm. *miricornis* Fauv.

Palaminus Er.

1. Size larger (3.4 mm.); antennae longer, the 4th to the 6th joints fully three times longer than broad; elytra longer (18) *bryanti* Cam.
- Size smaller (2.75 mm.); antennae shorter, the 4th to the 6th joints not more than twice as long as broad; elytra shorter (18) *parvus* Cam.

Paederus F.

1. Blue-black, thorax and first four abdominal segments red; antennae, palpi and legs (including the coxae) black . . . *tamulus* Er.

Astenus Steph.

1. Species reddish-testaceous, each elytron with an oblong black spot in the middle; base of the 6th abdominal segment black . . . *gracilentus* Fauv. (*gracilis* Kr.)
- Species darker, elytra immaculate; 6th abdominal segment concolorous. 2.
2. Sides of thorax and elytra with strong setae; elytra with moderately large superficial puncturation; abdomen finely punctured (19) *orientalis* Cam.
- Sides of thorax and elytra with weak setae; elytra with large deep puncturation; abdomen at the bases of the segments rather coarsely and deeply punctured (19a) *castaneus* Cam.

Stilicopsis Sachse.

1. Sides of the elytra with 3 or 4 long and strong setae; disc of thorax without median longitudinal impression. 2.
- Sides of the elytra without long setae; disc of the thorax with median longitudinal impression (21) *persimilis* Cam.
2. Antennae shorter, the 9th and 10th joints distinctly transverse; thorax broader, species smaller. ♂: 7th ventral segment with a deep acutely triangular excision; 6th with a broad

shallow emargination; 5th with a moderately broad, nearly semi-circular excision in the middle of the posterior border *breviceps* Fauv.

- Antennae longer, the 9th and 10th joints not transverse; thorax narrower, species larger. ♂: 7th ventral segment with a deep obtusely pointed excision, 6th with a small obtuse excision . . . (20) *obliqua* Cam.

Stilicis Lat.

1. Ferruginous red, dull, elytra testaceous, shining, with a small brown spot on the reflexed margin at the middle, and a larger one on the disc posteriorly on either side of the suture; abdomen pitchy with copper reflex, sericeous . . . *ocularis* Fauv.

Psilotrachelus Kr.

1. Black, head shining, moderately finely and closely punctured; thorax dull, closely granulate, with median shining keel; elytra with large punctures more or less in rows, the interspaces finely granulate. Antennae and legs reddish-brown. Length 4.5 mm. *crassus* Kr.

Thinocharis Kr.

1. Head transverse, subquadrate. 2.
— Head not transverse, subovate *pygmaea* Kr.
2. Species of darker colour; elytra more closely and distinctly punctured . . . (22) *nigricans* Cam.
— Species of lighter colour; elytra more sparingly and less distinctly punctured *carinicollis* Kr.

Acanthoglossa Kr.

1. Reddish-brown, clothed with long erect yellow pubescence; head and thorax closely punctured *hirta* Kr.

Medon Steph.

1. Base of the abdomen keeled below. Eyes moderate or small. 2.
— Base of the abdomen not keeled below. Eyes large. 4.

2. Prothoracic epimera present. Labrum more or less emarginate anteriorly in the middle, with the angles often dentiform. 6.
- Prothoracic epimera wanting. Labrum with a strong tooth in the middle of the emargination which projects a little beyond the anterior border (Sub-gen. *Charichirus*). 3.
3. Antennae entirely reddish-testaceous; legs testaceous; posterior part of the elytra more or less broadly and distinctly rufo-testaceous *chinensis* Boh.
- Antennae black, the last 3 or 4 joints reddish-testaceous; legs pitchy; posterior part of the elytra obscurely dull reddish (26a) *terminalis* Cam.
4. Mandibles 4-dentate (Sub-gen. *Isocheilus*). Species larger (7 mm.), blackish, elytra obscure testaceous, more or less extensively infusate on the disc . . . *staphylinoides* Kr.
- Mandibles with the right 4-dentate, the left 3-dentate (Sub-gen. *Arthocharis*). Species smaller. 5.
5. Head and thorax with distinct smooth median line; species brighter and more shining; abdomen much less thickly punctured and pubescent . . *ochracea* Grav.
- Head and thorax without median smooth shining line; species of duller colour, less shining; abdomen much more thickly punctured and pubescent . . *uvula* Kr.
6. Gular sutures fused or very narrowly separated (Sub-gen. *Medon* s.str.). 7
- Gular sutures not fused, widely separated. 9.
7. Elytra shining testaceous, the base broadly infusate; head and thorax bright reddish-testaceous, shining . (23) *rubicundus* Cam.
- Elytra otherwise coloured. 8.
8. Antennae slender, the 5th joint distinctly longer than broad; species smaller (3 mm.); vertex of head simply punctured, a little shining; elytra reddish, largely infusate posteriorly . (22a) *orientalis* Cam.

- Antennae short, the 5th joint scarcely longer than broad; species larger (3.75 mm.), vertex of head granular, not shining; elytra reddish, less infusate posteriorly *opacellus* Fauv.
- 9. Sculpture of head and thorax granular; species rufo-testaceous, the elytra more or less infusate posteriorly (26) *granulatus* Cam.
- Sculpture of head and thorax not granular. 10.
- 10. Head and thorax shining, without visible ground sculpture. 11.
- Head and thorax scarcely shining, with distinct coriaceous ground sculpture and superficial umbilicate puncturation *debilicornis* Woll.
- 11. Elytra testaceous, with broad pitchy transverse fascia; puncturation of thorax fine, not umbilicate (24) *fasciatus* Cam.
- Elytra uniform reddish-testaceous; puncturation of thorax moderately coarse, umbilicate (25) *lucens* Cam.

Parascopaeus Cam.

- 1. Shining pitchy-brown; head rather coarsely, thorax finely, elytra indistinctly punctured; antennae, mouth-parts and legs testaceous (27) *nitidus* Cam.

Scopaeus Er.

- 1. Antennae dark, the last four or five joints testaceous; species black; length 4 mm. (28) *niger* Cam.
- Antennae entirely reddish-testaceous; species at least partly testaceous or reddish-testaceous. 2.
- 2. Size larger (3.6 mm.); 2nd joint of the antennae distinctly shorter than the 3rd *testaceus* Motsch.
- Size smaller (2—2.5 mm.); 2nd joint of the antennae not shorter than the 3rd. 3.
- 3. 3rd joint of the antennae moniliform; under surface of the head deeply

- punctured; species red, shining, the elytra testaceous, with transverse indeterminate dark fascia nearer to the posterior margin than to the base. Length 2 mm. *puncticeps* Kr.
- 3rd joint of the antennae not moniliform; under surface of the head not deeply punctured. 4.
4. Species reddish-testaceous, shining, the elytra in great part infusate. 5.
- Species pale testaceous, but slightly shining; the elytra entirely pale, exceedingly finely and obsoletely punctured *pallidulus* Kr.
5. Elytra infusate, the posterior margin testaceous; 4th and 5th joints of the antennae a little longer than broad; species larger (2.5 mm.) and more robust *limbatus* Kr.
- Elytra testaceous with ill-defined dark fascia situated nearer the posterior border than the base; 4th and 5th joints of the antennae moniliform; species smaller (2 mm.). Very similar to *puncticeps* *micros* Kr.

Calliderma Motsch.

1. Thorax (except for parts of the elevated lines) entirely opaque, densely punctured (29) *rufum* Cam.
- Thorax entirely shining. 2.
2. Thorax red, impunctate (30) *nitens* Cam.
- Thorax red, with transverse dark fascia; in part coarsely and rugosely punctured (31) *rugicolle* Cam.

Cryptobium Mannerh.

1. Eyes very small; form narrow and elongate; legs reddish; size smaller (7 mm.). 7th dorsal segment with substrigose sculpture *filum* Kr.
- Eyes moderate; form broader; legs pale testaceous; size larger (8.5 mm.). 7th dorsal segment simply punctured (32) *foveatum* Cam.

Oligolinus Cas.

1. Black, shining; antennae, mouth-parts and legs reddish-testaceous, the middle and posterior tibiae more or less infusate (33) *parvus* Cam.

Leptacinus Er.

1. Black, shining; thorax reddish or pitchy, with a dorsal series of 5 or 6 large punctures; elytra pale testaceous, the base and apex somewhat infusate. Length 4 mm. *tricolor* Kr.

Somoleptus Sharp.

1. Shining, head and abdomen black, thorax and elytra pitchy-brown, the latter finely and sparingly punctured. Length 3 mm. (34) *linearis* Cam.

Eulissus Mannerh.

1. Black, shining; elytra and abdomen pitchy, the disc of the former obscure, the side margins of the latter clear testaceous (35) *lateralis* Cam.

Thyreocephalus Guer.

1. Shining bronze-green or coppery-bronze, the elytra and 3rd and 4th (visible) segments of the abdomen bright reddish; the last six joints of the antennae and legs testaceous. Length 10.5–15 mm. *annularis* Fauv.

Diochus Er.

1. Pitchy-black, shining; thorax entirely, apex of the elytra broadly, reddish-testaceous; abdomen pitchy-red, the apex testaceous. Antennae, mouth-parts and legs testaceous. Length 3 mm. (36) *pulchellus* Cam.

Holisomimus Cam.

1. Abdomen unicolorous pitchy-brown. Length 2–3 mm. (37) *parvus* Cam.
- Abdomen in part reddish-testaceous. Length 1.75 mm. (38) *cingulatus* Cam.

Actobius Fauv.

1. Black, shining; antennae and legs fuscous, the first two joints of the former and the femora testaceous. Length 4 mm. (39) *laticeps* Cam.

Philonthus Curtis.

1. Last joint of the labial palpi not longer than the preceding; head oblong (Subgen. *Gabrius*). 13.
- Last joint of the labial palpi longer than the preceding. 2.
2. Thorax on either side of the middle line with a row of three punctures,* elytra with a double series of large punctures, two sutural and three or four subhumeral. Length 6–9 mm. *notabilis* Kr.
- Thorax on either side of the middle line with a row of more than three punctures. 3.
3. Thorax on either side of the middle line with a row of four punctures; head suborbiculate; 1st joint of the antennae, coxae and legs testaceous, the tibiae often infusate; abdomen slightly iridescent. Length 6–8.5 mm. *delicatulus* Boh.
- Thorax on either side of the middle line with a row of five punctures. 4.
4. Antennae entirely rufo-testaceous; thorax and elytra castaneous-red; abdomen pitchy; legs testaceous. Length 6 mm. (41) *castaneipennis* Cam.
- Antennae dark, at most with the base and more or less of the apex lighter. 5.
5. Antennae with the base and at least the terminal joint reddish-testaceous, the penultimate joints strongly transverse. 6.
- Antennae with the base at most lighter. 7.
6. Antennae with the base and terminal joint rufo-testaceous; elytra entirely black. Length 3.75 mm. *crassicornis* Fauv.

* Erichson's notation.

- Antennae with the base and last four or five joints reddish-testaceous; elytra with the base, suture and apical margin red. Length 6 mm. *circumductus* Fauv.
- 7. Head small, narrow, oval; thorax narrowed in front. Length 6.5–7.5 mm. *longiceps* Fauv.
- Head subquadrate or suborbicular. 8.
- 8. Penultimate joints of the antennae distinctly transverse. 9.
- Penultimate joints of the antennae not or scarcely transverse. 10.
- 9. Elytra bronze-green, the suture narrowly reddish; 1st joint of the antennae pitchy-testaceous; elytra and abdomen more sparingly punctured. Length 8 mm. *aeneipennis* Boh.
- Elytra black, the suture and apical margin narrowly reddish; first two joints of the antennae clear reddish-testaceous; elytra and abdomen much more closely punctured. Length 5.75 mm. *flavocinctus* Motsch.
- 10. Head sub-quadrate. Species black, the sides of the elytra and posterior margins of the dorsal abdominal segments concolorous. Length 8–8.5 mm. (42) *belonuchoides* Cam.
- Head suborbicular. 11.
- 11. Base of the first three visible dorsal segments of the abdomen with a large puncture on either side of the middle line; elytra and abdomen black, concolorous; size smaller. Length 6 mm. *gemellus* Kr.
- Base of the first three visible dorsal segments of the abdomen without large puncture on either side. Size larger (8.5 mm.). 12.
- 12. Front of the head between the antennal tubercles with a short, deep, longitudinal sulcus in the middle line; diameter of the eyes viewed from above rather less than the length of the temples; 1st joint of the posterior tarsi scarcely longer than the last, Length 6.5–7 mm. (40) *sulcatus* Cam.

- Front of head without sulcus; the diameter of the eyes viewed from above much greater than the length of the temples; 1st joint of the posterior tarsi distinctly longer than the last *geminus* Kr.
- 13. Elytra pitchy, the apex and suture reddish-testaceous; penultimate joints of the antennae scarcely transverse . . . *pulchellus* Kr.
- Elytra uniformly fusco-testaceous; penultimate joints of the antennae distinctly transverse *maritimus* Motsch.

Orthidus Muls and Rey.

1. Shining brassy-bronze; elytra copper-bronze; antennae, mouth-parts and legs ferruginous. Length 10 mm. (43) *cupreipennis* Cam.

Cafius Steph.

1. Thorax with a narrow, shining, impunctate, median line; the rest of the surface closely and uniformly punctured; size larger (8 mm.) . . . *nauticus* Fairm.
- Thorax, with broader shining, impunctate median area, on either side with a row of 14 or 15 punctures, the sides more or less closely punctured; size smaller (5.5 mm.) *corallicola* Fairm.

Hesperus Fauv.

1. Black, shining; thorax, base of the elytra, 3rd and 4th visible abdominal segments and first three joints of the antennae, red; apical border of the elytra and of the 5th visible abdominal segment, last three or four joints of the antennae and the legs, pale testaceous. Length 8-8.5 mm. *laevigatus* Fauv.

Belonuchus Nordm.

1. Species entirely black *mutator* Fauv.
- Species not entirely black, the elytra with bronze-green reflex; antennae with the first three and last two joints testaceous *aeneipennis* Fauv.

Acylophorus Nordm.

1. Black, shining; abdomen iridescent; thorax with the sides strongly rounded, the disc with a single puncture on either side of the middle line. Length 6-7 mm. (44) *rotundicollis* Cam.

Delibius Fauv.

1. Obscure rufo-testaceous, rather shining; head posteriorly, disc of the elytra and middle of the abdomen more or less infusate. Length 3 mm. *longicornis* Fauv.

Megarhropsis Cam.

1. Elongate, fusiform, moderately shining, black, the sides and posterior half of the thorax, abdomen and legs reddish-testaceous; antennae long and slender, obscure testaceous, the middle joints infusate. Length 4 mm. (45) *decorata* Cam.

Atanygnathus Jacobson.

1. Black, shining, the thorax, posterior margins of the elytra and of the abdominal segments red; base and apex of the antennae and legs testaceous *terminalis* Er.

Conosoma Kr.

1. Sides of the elytra without setae.	2.
— Sides of the elytra with long setae.	11.
2. Species entirely or in great part black or reddish-brown.	3.
— Species in great part bright reddish-testaceous.	10.
3. Elytra with a more or less distinct macula at the base of each.	4.
— Elytra immaculate at the base.	5.
4. Each elytron with a reddish, ill-defined spot at the middle of the base, not extending to the lateral margin; posterior margin and postero-external angles of the thorax rufescent	46 (a) <i>malayanum</i> Cam.

- Each elytron with a rounded, well-defined yellow spot at the middle of the base; posterior angles of the thorax yellow (48) *flavoguttatum* Cam.
- 5. Antennae very long and slender, the joints not appreciably compressed . . . (50) *championi* Cam.
- Antennae shorter, the joints distinctly compressed. 6.
- 6. Abdomen shining; species shining, much less thickly punctured and pubescent . (49) *abdominale* Cam.
- Abdomen dull; species more opaque, much more thickly punctured and pubescent. 7.
- 7. Species larger and more robust. Length 5 mm. (46) *robustum* Cam.
- Species smaller and less robust. Length 3-3.5 mm. 8.
- 8. Pectinations of the anterior tibiae on the outer border testaceous; posterior third of the elytra obscurely reddish (51) *walkeri* Cam.
- Pectinations of the anterior tibiae on the outer border, black; elytra unicolorous. 9.
- 9. Species larger and broader; black. Length 3.5 mm. *ceylanense* Kr.
- Species smaller and narrower; reddish-brown (47) *rufobrunneum* Cam.
- 10. Base of the thorax with a black, subtriangular spot on either side of the middle line (sometimes united); base of the elytra broadly, and apex narrowly, rufo-testaceous; 4th visible abdominal segment (except the posterior border) black *suave* Fauv.
- Base of the thorax immaculate; elytra obscurely darker posteriorly; abdomen concolorous (52) *perplexum* Cam.
- 11. Base of the thorax with a black spot on either side of the middle line; elytra with a black fascia extending from the lateral margin nearly to the suture; 6th to the 10th joints of the antennae black (53) *nigromaculatum* Cam.

- Base of the thorax immaculate, elytra without dark fascia; 6th to the 10th joints of the antennae scarcely infusate (54) *rufotestaceum* Cam.

Tachinomorphus Kr.

- 1. Last joint of the antennae testaceous, the penultimate joints much less transverse *fulvipes* Er.
- Last joint of the antennae black, the penultimate joints strongly transverse *ceylonicus* Bernh.

Coproporus Kr.

- 1. Antennae entirely testaceous. 2.
- Antennae at least in part dark. 3.
- 2. 4th joint of antennae longer than broad, the penultimate joints scarcely transverse; head and thorax entirely impunctate; species smaller (1.75 mm.), rufo-testaceous (57) *parvulus* Cam.
- 4th joint of the antennae distinctly transverse, the penultimate joints distinctly transverse; head and thorax exceedingly finely punctured; species larger (2.2 mm.), black, pitchy or more or less rufo-testaceous (56a) *varians* Cam.
- 3. 5th joint of the antennae strongly transverse; elytra finely but distinctly punctured; species minute. Length 1 mm. *atomus* Kr.
- 5th joint of the antennae not transverse; species larger. 4.
- 4. 5th joint of the antennae distinctly longer than broad. 5
- 5th joint of the antennae as long as broad. 9.
- 5. Head clear reddish-testaceous; species in great part reddish-testaceous. 6.
- Head black or pitchy-red. 7.
- 6. Elytra testaceous-yellow without dark markings *secretus* Bernh.
- Elytra testaceous-yellow at the base, with a large black spot not extending to the suture or the lateral margin *fasciipennis* Kr.

7. Elytra testaceous (56) *flavipennis* Cam.
 — Elytra dark. 8.
8. Head, thorax and elytra without trace of puncturation; head and abdomen reddish; species larger and more convex. Length 4.5 mm. (55) *rufiventris* Cam.
- Head, thorax and elytra finely but distinctly punctured; head and abdomen black; species smaller and more depressed. Length 2.75 mm. *subdepressus* Kr.
9. Thorax before the base with a large puncture on either side of the middle line, otherwise completely impunctate; elytra distinctly punctured; size larger and more convex. Length 3 mm. *brunneicollis* Motsch.
- Thorax without large punctures before the base, scarcely perceptibly punctured; elytra exceedingly finely punctured; size smaller and more depressed. Length 1.75 mm. *minimus* Motsch.

Leucoparyphus Kr.

1. Black, shining, the margins of the thorax, base, shoulders, postero-external angles and apical margins of the elytra testaceous. Length 2.75–3 mm. *silphoides* L.

Mimocyptus Cam.

1. Convex, shining ferruginous-red, the foreparts impunctate, the abdomen scarcely perceptibly punctured; antennae with the first four joints and apex of the last, mouth-parts and legs testaceous. Length 1.2 mm. (in well-extended examples) (58) *globulus* Cam.

Adinopsis Cam.

1. Minute, obscure reddish-brown, densely and finely punctured and pubescent; antennae, mouth-parts and legs testaceous. Length 1.2 mm. (59) *rufobrunnea* Cam.

Leucocraspedum Kr.

1. Black, convex, shining, acuminate posteriorly; antennae stout, testaceous; the last two joints infusate; legs pitchy-testaceous. Length scarcely 3 mm. (60) *nigrum* Cam.

Myllaena Er.

1. Narrow, elongate, acuminate, blackish; the thorax, elytra, posterior margins of the first four visible and the whole of the last two abdominal segments reddish-testaceous; antennae, mouth-parts and legs clear testaceous. Length 3·4 mm. (61) *faberensis* Cam.

Pronomaea Er.

1. Chestnut brown, rather shining, the fore-parts finely and closely punctured; antennae fuscous, the first two joints, palpi and legs testaceous. Length 3 mm. (62) *leontopolitana* Cam.

Diglotta Champ.

1. Linear, pitchy; abdomen black, scarcely shining, densely and finely pubescent; antennae, elytra, legs and last abdominal segment testaceous. Length 1·5 mm. (63) *testaceipennis* Cam.

Oligota Mannerh.

1. Black, elytra pitchy-brown, apex of abdomen rufescent. Length scarcely ·75 mm. (64) *moultoni* Cam.
— Entirely testaceous. Length ·75 mm. (65) *forticornis* Cam.

Pseudoligota Cam.

1. Last seven joints of the antennae infusate; species narrow and less robust; length 1 mm. ♂: sutural margin of the elytra posteriorly with four or five minute tubercles on either side; spine of the 8th abdominal segment furnished with yellow setae (66) *varians* Cam.

- Last seven joints of the antennae black; species broader and more robust; length 1.1 mm. ♂: sutural margin of the elytra posteriorly with a small tubercle on either side and sometimes with traces of two others in front; spine of 8th abdominal segment without setae (61) *robusta* Cam.

Gyrophæna Mannerh.

1. Species larger (2.5–3 mm.), thorax and greater part of the abdomen bright reddish-testaceous. ♂: sides of 4th (visible) dorsal segment with a narrow plate pointed at apex; 7th with a row of six small tubercles placed transversely across the middle; 8th trifold, the lateral lobes broader than the median *appendiculata* Motsch.
- Species smaller (1.5–2.5 mm.), coloration more obscure, pitchy or metallic, with elytra and base of the abdomen often more or less obscure testaceous. ♂ without lateral appendage to the 4th dorsal segment. 2.
2. Head strongly transverse, eyes very prominent (*Gyrophæna* s.str.). 3.
- Head much less transverse, eyes less prominent (Sub-gen. *Phaenogyra* Rey). Species with copper-bronze metallic reflex on the fore-parts. ♂: 7th dorsal segment with obsolete tubercle on either side of the middle line in front of the posterior border; 8th dorsal segment with a rather slender, slightly incurved spine on either side, the border between with two minute teeth separated by a feeble emargination and much nearer to the lateral spines than to each other (76) *metallica* Cam.
3. Thorax with a row of two or more larger punctures on either side of the disc. 4.
- Thorax without distinct row of larger punctures on either side of the disc,

which is either irregularly or scarcely at all punctured.

7.

4. Antennae entirely testaceous; species small .5–1.75 mm.

5.

— Antennae with the last joints black, average size larger (1.5–2.5 mm.). ♂: 8th dorsal segment with a median stout spine and on either side a slightly curved, pointed tooth projecting a little beyond the level of the apex of the median spine and separated from it by a semicircular excision . (68) *tridentata* Cam.

5. 5th joint of the antennae not transverse, size larger (1.75 mm.). ♂: 7th dorsal segment with a very obsolete row of six tubercles; 8th with a large flat central tubercle at the base, the posterior margin on either side with a feeble emargination, so that it presents three rounded crenulations, the central one being the largest and most prominent (70) *crenulata* Cam.

— 5th joint of the antennae transverse, size smaller (.5–1.2 mm.).

6.

6. Head and thorax with fine transverse, strigose ground-sculpture. Length 1.2 mm. ♂: 7th dorsal segment with a curved, transverse row of six small tubercles, of which the central pair are considerably larger, the lateral sometimes more or less obsolete; 8th narrowed and slightly emarginate on either side, so as to form three short processes, the central one bluntly rounded, wider and more produced than the lateral, which are triangular . (69) *granulosa* Cam.

— Head and thorax without ground-sculpture. Length .6 mm. ♂: postero-external angles of the elytra with a strong, raised oblique crest; 8th dorsal segment narrowed and rounded (71) *cristata* Cam.

7. Thorax exceedingly finely and sparingly punctured, ground-sculpture distinct, transversely strigose. ♂: 8th dorsal

- segment produced on either side into a rather stout, slightly incurved spine, the posterior margin between these bisinuate (72) *bidens* Cam.
- Thorax distinctly punctured; ground-sculpture absent or very indistinct. 8.
8. Elytra exceedingly finely and exceedingly sparingly punctured; middle of the disc of thorax impunctate . . . (73) *dubia* Cam.
- Elytra finely, but not exceedingly sparingly punctured; middle of the disc of thorax punctured. 9.
9. Thorax finely and uniformly punctured.
 ♂: dorsal segment with a flat, semi-circular tubercle in front of the posterior margin in the middle; 8th with a small triangular excision on either side of the middle of the posterior border, so that a short, blunt, triangular process is formed in the middle and the lateral margins project as sharp triangular teeth (75) *moultoni* Cam.
- Thorax with unequal, irregular punctures on the disc, the sides impunctate.
 ♂: 8th dorsal segment with a short, stout, blunt, slightly incurved tooth on either side (74) *irregularis* Cam.

Sternotropa Cam.

1. Species black, the elytra often chestnut-brown. ♂: 8th dorsal segment with a pointed tooth on either side, separated by a nearly semi-circular emargination (77) *nigra* Cam.
- Species black, the thorax, base and apex of the abdomen reddish-testaceous. ♂: suture of the elytra with a row of three obsolete tubercles towards the posterior part; 7th dorsal segment with a minute tubercle on either side of the middle line in front of the posterior margin; 8th with a moderately long, pointed, slightly incurved tooth on either side (78) *ruficollis* Cam.

Adelarthra Cam.

1. Shining dark pitchy-red; elytra pitchy-black; 3rd, 4th and 8th abdominal segments reddish-testaceous. Length 1.2 mm. (79) *barbara* Cam.

Hetairotermes, n.n.

1. Disc of the elytra glabrous, the sides and postero-external angles finely and moderately closely punctured . . . (80) *agilis* Cam.
- Disc of the elytra uniformly but sparingly punctured (81) *piceus* Cam.

Pseudatheta Cam.

1. Rufo-testaceous, the elytra (except the base), posterior half of the 5th and whole of the 6th abdominal segments black. Length 1.75 mm. (82) *elegans* Cam.

Pelekoglossa Cam.

1. Pitchy, moderately shining, the thorax, base and apex of the abdomen obscure reddish-testaceous; first three joints of the antennae fusco-testaceous . . . (83) *cingulata* Cam.

Placusa Er.

1. 5th joint of the antennae as broad as long; species larger (2 mm.), pitchy, elytra testaceous, infusate at the scutellum. ♂: 8th dorsal segment of the abdomen finely crenulate; 6th ventral segment narrowed and produced (84) *conura* Cam.
- 5th joint of the antennae distinctly transverse, species smaller (1 to 1.6 mm.). 2.
2. First three joints of the antennae clear testaceous; species larger (1.6 mm.) and more robust; thorax pitchy, elytra obscure testaceous. ♂: 8th dorsal segment deeply excised on either side, the lateral margin forming a long, sharp, incurved spine, the central portion forming a lobe with rounded apex furnished with a broad, flat tubercle on the disc (85) *bispina* Cam.

- First three joints of the antennae pitchy-testaceous; species smaller (1—1.2 mm.) and narrower. 3.
3. Thorax black, more than half as broad again as long, species less finely punctured. ♂: 8th dorsal segment as in *bispina*, but the central lobe without tubercle (86) *lobata* Cam.
- Thorax pitchy, one-third as broad again as long; species more finely punctured. ♂: 8th dorsal segment with three equal and closely placed teeth at the middle of the posterior border, which is deeply emarginate on either side, the emargination bounded externally by a long, sharp, incurved spine; 3rd dorsal segment variable, either (1) the posterior border triangularly produced in the middle into a lobe with rounded apex, broadly emarginate on either side, the emargination bounded externally by a stout spine extending nearly to the level of the posterior border of the 4th segment, or (2) the posterior margin only slightly produced into a broader lobe emarginate posteriorly, otherwise as in (1) or (3), the posterior margin not at all produced and the lateral spines shorter and broader . . . (87) *notabilis* Cam.

Pseudoplacusa Cam.

1. Rufo-testaceous, moderately shining, head and elytra pitchy-black; antennae reddish-testaceous. Length 2.1 mm. (88) *rufiventris* Cam.

Chledophila Cam.

1. Narrow, elongate, scarcely shining, rufo-testaceous, the head pitchy-red; elytra, 6th abdominal segment and last seven joints of the antennae fuscous. Length 2.1 mm. (89) *annularis* Cam.

Neosilusa Cam.

1. Head and thorax less shining, the puncturation finer but more rugose . . . *ceylonica* Kr.
- Head and thorax more shining, the puncturation coarser but less rugose . . . (90) *moultoni* Cam.

Ousilusa Cam.

1. Head coarsely and closely punctured, sides of the thorax exceedingly finely punctured, the disc (except posteriorly) with moderately fine umbilicate puncturation; elytra very coarsely and deeply punctured . . . (91) *myrmecobia* Cam.
- Head finely, superficially and closely punctured; thorax finely and uniformly punctured; elytra superficially sculptured with larger and smaller punctures . . . (92) *castanea* Cam.

Prosilusa Cam. .

1. Rufo-castaneous, shining, finely punctured; the elytra in great part, and the base of the 6th abdominal segment, pitchy-black. Antennae, mouth-parts and legs reddish-testaceous . . . (93) *rufa* Cam.

Deralia Cam.

1. Pitchy-black, moderately shining; thorax and abdomen (except the 5th and 6th segments, which are pitchy) rufo-testaceous; antennae fuscous, the first three joints and apex of the last mouth-parts and legs testaceous . . . (94) *fuscipennis* Cam.

Pseudophaena Cam.

1. Rufo-castaneous, shining; elytra strongly punctured; first three joints of the antennae, mouth-parts and legs reddish-testaceous, the anterior and apex of the intermediate and posterior tibiae pitchy. Length 1.8 mm. . . (95) *castanea* Cam.

Coenonica Kr.

1. Thorax pitchy-red, the sculpture consisting of granules . . . , (97) *stricticollis* Cam.

— Thorax black, the sculpture consisting of punctures.

2.

2. Head between the antennal tubercles smooth and shining; thorax broader, more shining, less closely punctured . . . *puncticollis* Kr.

— Head between the antennal tubercles coarsely punctured; thorax narrower, less shining, more closely punctured . . . (96) *angusticollis* Cam.

Mimomalota Cam.

1. Species dark; size larger and more robust . . . (98) *bispina* Cam.

— Species in great part testaceous; size smaller and less robust . . . (99) *testacea* Cam.

Neomalota Cam.

1. Pitchy or reddish-brown, the elytra black, the abdomen reddish-testaceous, with the 6th segment blackish . . . (100) *cingulata* Cam.

Lampromalota Cam.

1. Depressed, shining; head and thorax very finely and sparingly punctured; elytra and abdomen fusco-testaceous (101) *brunneicollis* Cam.

Homalota Mannerh.

1. Antennae in great part black or infusate.	2.
— Antennae clear reddish-testaceous.	7.
2. Elytra more or less testaceous.	3.
— Elytra dark.	5.
3. Thorax reddish-brown; size larger (2 mm.); 6th and 7th abdominal segments much more sparingly punctured than the preceding	<i>platygaster</i> Kr.

— Thorax black; size smaller; 6th and 7th abdominal segments similarly punctured to those preceding.

4.

4. Head and thorax dull, densely coriaceous, the puncturation confused with the ground-sculpture; elytra fusco-testaceous *tuberculicollis* Kr.

— Head and thorax a little shining, less coriaceous, the puncturation more distinct from the ground-sculpture; elytra clear testaceous (102) *nitescens* Cam.

5. 4th joint of the antennae longer than broad, the penultimate joints scarcely transverse; size larger (2 mm.) . . . (105) *fuscipennis* Cam.
- 4th joint of the antennae distinctly transverse, the penultimate joints strongly transverse; size smaller. 6.
6. Abdomen reddish-testaceous, the 6th segment pitchy; posterior angles of the thorax forming a minute tooth . . (104) *cingulata* Cam.
- Abdomen black, the posterior half of the 7th and the 8th segment reddish-testaceous; posterior angles of the thorax not forming a tooth . . . (103) *bidens* Cam.
7. 4th joint of the antennae square, size smaller (1.75 mm.) *variventris* Kr.
- 4th joint of the antennae distinctly transverse; size larger (2–2.4 mm.). 8.
8. Larger and more robust; fore-parts more coarsely punctured; length 2.4 mm. ♂: 8th dorsal segment emarginate on either side, the emargination bounded externally by a rather long, sharp tooth, the posterior border between the emarginations with eight sharp teeth (107) *denticulata* Cam.
- Smaller and less robust; fore-parts more finely sculptured; length 2 mm. ♂: 8th dorsal segment with a feeble emargination on either side, bounded externally by a small tooth, the posterior border between the emarginations finely serrate (106) *serrata* Cam.

Thectura Thoms.

1. Rather shining; head black, thorax pitchy-brown, elytra and 5th and 6th abdominal segments pitchy, the rest of the latter reddish-testaceous. Length 1.6 mm. (108) *brunneicollis* Cam.

Heterota Rey.

1. Black, with greasy lustre; elytra with indeterminate orange spot occupying the sutural region towards the apex;

antennae, mouth-parts and legs reddish-testaceous (109) *arenaria* Cam.

Paractocharis Cam.

1. Very narrow, elongate, fragile, depressed; obscure brown, head and abdomen black; antennae and legs testaceous. Length 1.4 mm. *fucicola* Cam.

Falagria Mannerh.

1. Thorax cordiform; the sides and base not bordered; the elytra with a distinct impression internal to the shoulders (Sub-gen. *cardiola* Muls and Rey); species ferruginous, pubescence erect *vestita* Boh.
- Thorax not cordiform; the sides and base finely bordered; the elytra without impression internal to the shoulders. 2.
2. Scutellum on either side with a raised line following the lateral border and united at the apex, the base more or less keeled; species larger (2.8–3 mm.). 3.
- Scutellum without raised line at the lateral borders, the base not keeled; species smaller (1.75 mm.). 5.
3. Head and thorax shining. 4.
- Head and thorax dull; antennae very slender, all the joints much longer than broad; postero-external angles of the thorax acute, prominent (110) *tenuicornis* Cam.
4. Postero-external angles of the thorax prominent, acute; antennae stouter, the 10th joint scarcely as long as broad (111) *brevicornis* Cam.
- Postero-external angles of the thorax not prominent, obtuse; antennae much more slender, the 10th joint much longer than broad *dimidiata* Motsch.
5. Pitchy-black; vertex of head not sulcate; antennae brown, the base reddish-testaceous; elytra fusco-testaceous *pygmaea* Kr.
- Black; vertex of head with deep longitudinal sulcus; antennae entirely dark; elytra testaceous with base infusate (112) *flavipennis* Cam.

Amaurodera Fauv.

1. Head and elytra shining, brown, very finely and sparingly punctured, thorax reddish-brown, opaque, strongly shagreened; 2nd to 4th segments of the abdomen testaceous, the rest black *veluticollis* Motsch.

Eusteniamorpha Cam.

1. Rufo-castaneous, moderately shining, the 6th abdominal segment pitchy-black; thorax and abdomen strongly constricted at their base (113) *rufa* Cam.

Pelioptera Kr.

1. Antennae with the first three joints reddish-testaceous. Species shining; size larger (2-2.75 mm.) *micans* Kr.
- Antennae entirely dark. Species with greasy lustre only; size smaller (1.75 mm.) *opaca* Kr.

Atheta Thoms.

1. Abdomen more or less pointed posteriorly. 12.
- Abdomen parallel-sided. 2.
2. Antennae with the penultimate joints not or scarcely transverse. 3.
- Antennae with the penultimate joints distinctly transverse. 5.
3. Abdomen glabrous; Species bright reddish-testaceous, the 5th, 6th and anterior part of the 7th abdominal segments black (121) *miriventris* Cam.
- Abdomen finely and uniformly punctured. 4.
4. Species dark; elytra uniformly pitchy-brown or pitchy-black (115) *alophila* Cam.
- Species reddish; elytra pitchy, the base and apical margin testaceous; 5th, 6th and anterior part of the 7th abdominal segments black (114) *moultoni* Cam.
5. 3rd joint of the antennae scarcely shorter than the 2nd; species larger (2.5 mm). *dilutipennis* Motsch.

- 3rd joint of the antennae distinctly shorter than the 2nd; species smaller (1.3–1.75 mm.). 6.
- 6. Head and thorax with metallic copper-bronze reflex (120) *purpurascens* Cam.
- Head and thorax without metallic reflex. 7.
- 7. Species in great part testaceous, the head and 5th and 6th abdominal segments black *putridula* Kr.
- Species obscurely coloured, black, pitchy or brown. 8.
- 8. Thorax slightly transverse (116) *picea* Cam.
- Thorax distinctly transverse. 9.
- 9. Antennae lighter at the base. 10.
- Antennae entirely dark. 11.
- 10. Species shining, more depressed; 4th joint of the antennae but slightly broader than long. ♂: 8th dorsal segment of the abdomen truncate; 6th ventral segment produced, narrowed and rounded at the apex (118) *malayana* Cam.
- Species with greasy lustre only, less depressed; 4th joint of the antennae distinctly transverse. ♂: 8th dorsal segment of the abdomen truncate on either side with a small tooth . *inutilis* Kr.
- 11. Intermediate tibiae with a distinct seta near the middle. ♂: 8th dorsal segment of the abdomen truncate; 6th ventral segment a little produced, narrowed and rounded (119) *vulgaris* Cam.
- Intermediate tibiae without distinct seta; ♂: 8th dorsal segment of the abdomen with deep and broad semicircular emargination of the posterior borders. (117) *melata* Cam.
- 12. Sides of the thorax uniformly rounded, the lateral setae feeble or absent, the epipleurae not visible when viewed laterally; abdomen in some thickly punctured and pubescent throughout. 17.
- Sides of the thorax contracted behind, the lateral setae distinct; the epipleurae visible when viewed laterally;

- abdomen always more sparingly
punctured behind. 13.
13. 3rd joint of the antennae distinctly
shorter than the 2nd; species smaller
(1-1.2 mm.) Sub-gen. *Datomicra* Rey. 14.
- 3rd joint of the antennae not shorter than
the 2nd; species larger (1.8-2.5 mm.).
Sub-gen. *Dimetrota* Rey. 15.
14. Antennae entirely dark; thorax closely
and asperately punctured; species
uniformly dark (125) *onthophila* Cam.
- Antennae with the first two joints tes-
taceous; thorax finely and not asper-
ately punctured; thorax, base of the
elytra, base and apex of the abdomen
more or less pitchy-red (126) *mycetophila* Cam.
15. 7th joint of the antennae longer than
broad (124) *mycetophaga* Cam.
- 7th joint of the antennae distinctly
transverse. 16.
16. Abdomen in great part reddish-tes-
taceous; thorax pitchy-red . . . (122) *carpophila* Cam.
- Abdomen and thorax black . . . (123) *xylophila* Cam.
17. Abdomen thickly punctured and pu-
bescent throughout (Sub-gen. *Colpo-
dota* Rey). Pitchy, greasy-lustrous,
elytra testaceous, infusate at scutel-
lum; penultimate joints of antennae as
long as broad (127) *ruparia* Cam.
- Abdomen much less thickly punctured
and pubescent, especially behind
(Sub-gen. *Acrotona* Rey). 18.
18. Head black; abdomen reddish-testaceous,
intermediate and posterior tibiae each
with two distinct setae . . . (128) *rufiventris* Cam.
- Head testaceous; 6th abdominal segment
blackish; intermediate and posterior
tibiae without distinct setae . . . *annuliventris* Kr.

Exatheta Cam.

1. Puncturation of the thorax very fine,
not asperate; ground sculpture
wanting (129) *cingulata* Cam.

- Punctuation of the thorax fine, asperate;
ground-sculpture visible . . . (130) *consors* Cam.

Mimatheta Cam.

1. Black, moderately shining; elytra obscure testaceous, more or less infusate at the sides; first three joints of the antennae mouth-parts and legs, testaceous. Length 2 mm. . . (131) *fungicola* Cam.

Mimacrotona Cam.

1. Rufo-testaceous, moderately shining; the head, 5th, 6th and base of the 7th abdominal segments blackish; the elytra more or less infusate; first three joints of the antennae, mouth-parts and legs testaceous. Length 1.2 mm. . . (132) *cingulata* Cam.

Paratheta Cam.

1. Black, rather shining; the elytra castaneous-brown; first three joints of the antennae and legs reddish-testaceous. Length 2 mm. . . (133) *carnivora* Cam.

Fenyesia Cam.

1. Black, shining, robust, convex; last two joints of the antennae and legs testaceous, the femora infusate. Length 2 mm. . . (134) *nigra* Cam.

Termitoptochus Silv.

1. ♀: Reddish, elongate, laevigate; abdomen inflated, obtriangular, reflexed forwards. . . *indicus* Silv.

Myrmedonota Cam.

1. Black or pitchy-black, shining; the first two visible segments of the abdomen testaceous-yellow; first two joints of the antennae and the legs testaceous, the apices of the femora and the tibiae more or less pitchy. Length 3 mm. . (135) *cingulata* Cam.

Myrmedonia Er.

1. 2nd joint of the antennae much shorter than the 3rd; 3rd and following joints compressed. Size larger; species reddish-brown *indorum* Fauv.
- 2nd joint of the antennae but little shorter than the 3rd; 3rd and following joints not compressed. Size smaller; species shining black, the base of the elytra and first four visible abdominal segments reddish-testaceous (136) *apicalis* Cam.

Schistogenia Kr.

1. Reddish-brown, opaque, coarsely and rugosely punctured *crenicollis* Kr.

Myrmedonella Cam.

1. Bright rufo-testaceous, shining; the elytra pitchy-red. Antennae, mouth-parts and legs reddish-testaceous. Length 1.8 mm. (137) *rufa* Cam.

Tetrasticta Kr.

1. Black, shining, base of the antennae and of the abdomen and legs testaceous; elytra brown *polita* Kr.

Paraleochara Cam.

1. Shining castaneous, elytra pitchy-black; abdomen reddish-testaceous, the 6th and 7th segments black; first three joints of the antennae, mouth-parts and legs testaceous (138) *fungivora* Cam.

Hoplandria Kr.

1. Pitchy-brown, shining, narrowed behind; elytra darker, base of the abdomen lighter, first four and apex of the last joints of the antennae, mouth-parts and legs testaceous (139) *frugivora* Cam.

Aleochara Grav.

1. Mesosternum simple, the antennae stout, the penultimate joints three times as broad as long (Sub-gen. *Heterochara*

Rey). Black, shining, the elytra red, with a large lateral spot black; base of the antennae and legs testaceous.

Length 3 mm. var. *maculipennis* Kr. (*croceipennis* Motsch.).

- Mesosternum keeled. 2.
- 2. Elytra sinuate internal to the postero-external angles. Species pitchy, the elytra red, with triangular scutellary marking and the sides dark; abdomen very thickly punctured in front. . *puberula* Klug.
- Elytra not sinuate. 3.
- 3. Penultimate joints of the antennae three times broader than long. Species black, the first three joints of the antennae pitchy-testaceous; legs reddish-testaceous . . . *nigra* Kr.
- Penultimate joints of the antennae moderately transverse. 4.
- 4. Thorax with all the margins narrowly but distinctly testaceous; first two joints of the antennae clear testaceous . . *asiatica* Kr.
- Thorax at most with the lateral margins obscurely reddish-testaceous; first two joints of the antennae red . . *viatica* Faun.

LIST OF THE STAPHYLINIDAE OF SINGAPORE *

I. Subfam. OXYTELINAE.

I. Tribe PIESTINI.

Sub-tribe ELEUSII.

Genus *Eleusis* Cast.

fusciceps Kr. Arch. Naturgesch., xxv, 1859, i, p. 184.

humilis Er. Gen. Spec. Staph., p. 839.

kraatzii Fauv. Ann. Mus. Civ. Gen., xii, 1878, p. 207.

lunigera Fauv. Rev. d'Ent., xxiii, 1904, p. 84.

* Species of which the type form is not recorded from Singapore are placed in brackets.

Sub-tribe *LEPTOCHIRI*.

Genus **Leptochirus** Germ.

Sub-gen. *Strongylochirus* Bernh.

laevis Cast. Hist. Nat., i, 1840, p. 186.

Genus **Borolinus** Bernh.

(*minutus* Cast. Hist. Nat., i, 1840, p. 186.)

var. *cruentus* Fauv. Rev. d'Ent., xiv, 1895, p. 181.

Genus **Priochirus** Sharp.

Sub-gen. *Triacanthus* Bernh.

(*tridens* Motsch. Bull. Mosc., xxx, 1857, ii, p. 502.)

var. *insularis* Bernh. D. E. Z., 1903, p. 139.

Sub-gen. *Cephalomerus* Bernh.

hoplites Fauv. Rev. d'Ent., xiv, 1895, p. 182.

pygmaeus Kr. Arch. Naturgesch., xxv, 1859, i, p. 191.

Sub-tribe *LISPINI*.

Genus **Ancaeus** Fauv.

exiguus Er. Gen. Spec. Staph., p. 830.

singularis n. sp. Trans. Ent. Soc., 1918, p. 58.

Genus **Holosus** Motsch.

plicatus Bernh. W. Z. B., liv, 1904, p. 14.

tachyporiformis Motsch. Bull. Mosc., xxx, 1857, ii, p. 498.

Genus **Lispinus** Er.

coarcticollis Kr. Arch. Naturgesch., xxv, 1859, p. 186.

impressicollis Motsch. Bull. Mosc., xxx, 1857, ii, p. 495.

minutus n. sp. Trans. Ent. Soc., 1918, p. 60.

setosus, n. sp. Trans. Ent. Soc., 1918, p. 59.

sharpi, n. sp. Trans. Ent. Soc., 1920, p. 278.

tenuicornis Kr. Arch. Naturgesch., xxv, 1859, i, p. 187.

II. Tribe *OMALIINI*.

Genus **Phloeonomus** Heer.

Sub-gen. *Phloeonomus* s.str. Ganglb.

discahis Cam. Trans. Ent. Soc., 1913, p. 525.

obscurus Kr. Arch. Naturgesch., xxv, 1859, i, p. 181.

III. Tribe OXYTELINI.

Sub-tribe OXYTELI.

Genus **Trogophloeus** Mannh.Sub-gen. *Trogophloeus* s.str.*orientalis*, n. sp. Trans. Ent. Soc., 1918, p. 61.*silvestris*, n. sp. Trans. Ent. Soc., 1918, p. 61.Sub-gen. *Taenosoma* Mannh.*halophiloides*, n. sp. Trans. Ent. Soc., 1918, p. 62.*littoralis*, n. sp. Trans. Ent. Soc., 1918, p. 63.*lucens*, n. sp. Trans. Ent. Soc., 1918, p. 63.*rufotestaceus*, n. sp. Trans. Ent. Soc., 1918, p. 64.Genus **Oxytelus** Grav.Sub-gen. *Caccoporus* Thoms.*bengalensis* Er. Gen. Spec. Staph., p. 789.*ferrugineus* Kr. Arch. Naturgesch., xxv, 1859, p. 173.*nigriceps* Kr. Arch. Naturgesch., xxv, 1859, i, p. 171.*exasperatus* Kr. Arch. Naturgesch., xxv, 1859, i, p. 175.*frugicola*, n. sp. Trans. Ent. Soc., 1918, p. 67.*granadillae*, n. sp. Trans. Ent. Soc., 1918, p. 66.*kraatzi*, n. n.*pulcher* Kr. Arch. Naturgesch., xxv, 1859, i, p. 173.*latiusculus* Kr. Arch. Naturgesch., xxv, 1859, i, p. 176.*nitidifrons* Woll. Ann. Mag. Nat. Hist., (4) viii, 1871, p. 411.*obscurus*, n. sp. Trans. Ent. Soc., 1918, p. 67.*pygmaeus* Kr. Arch. Naturgesch., xxv, 1859, i, p. 176.*raffrayi* Fauv. Rev. d'Ent., xxiv, 1905, p. 117.*thoracicus* Motsch. Bull. Mosc., 1857, iv, p. 504.Genus **Bledius** Mannerh.*bellicosus* Fauv. Rev. d'Ent., xxiii, 1904, p. 111.Sub-tribe *THINOBII*.Genus **Thinobius** (s.str.) Muls. & Rey.*marinus* Cam. E. M. M., 1917, p. 155.

IV. Tribe OSORIINI.

Sub-tribe *OSORII*.Genus **Paragonus** Fauv.*heteroceros* Fauv. Rev. d'Ent., xxiv, 1905, p. 134.

Genus **Mimogonus** Fauv.

fumator Fauv. Rev. d'Ent., viii, 1889, p. 246.

Genus **Holotrochus** Er.

nitidus, n. sp. Trans. Ent. Soc., 1918, p. 68.

Genus **Osorius** Latr.

rugifrons Er. Gen. Spec. Staph., p. 756.

II. Sub-fam. **MEGALOPSINAE.**

I. Tribe MEGALOPSINI.

Genus **Megalops** Er.

? sp. The specimen having escaped.

III. Sub-fam. **STENINAE.**

I. Tribe STENINI.

Genus **Stenus** Latr.

Sub-gen. *Hypostenus* Rey.

(?) *bivulneratus* Motsch. Bull. Mosc., 1857, ii, p. 514.

castaneus, n. sp. Trans. Ent. Soc., 1918, p. 69.

fortepunctatus, n. sp. Trans. Ent. Soc., 1918, p. 68.

monmeros Fauv. Rev. d'Ent., xiv, 1895, p. 214.

IV. Sub-fam. **EVAESTHETINAE.**

I. Tribe STENAESTHETINI.

Genus **Stenaesthetus** Sharp.

sunioides Sharp. Trans. Ent. Soc., 1874, p. 80.

II. Tribe EVAESTHETINI.

Genus **Edaphus** J. Lec.

dentiventris Fauv. Rev. d'Ent., xxiv, 1905, p. 137.

TRANS. ENT. SOC. LOND, 1920.—PARTS III, IV, V. (APR.'21) D D

V. Sub-fam. **PAEDERINAE**.

I. Tribe PINOPHILINI.

Sub-tribe *PINOPHILI*.Genus **Pinophilus** Grav.

borneensis Fauv. Rev. d'Ent., xiv, 1895, p. 221.

orientalis Cam. Trans. Ent. Soc., 1920, p. 278.

Neopinophilus, n. gen. Trans. Ent. Soc., 1920, p. 279.

notabilis, n. sp. Trans. Ent. Soc., 1918, p. 70 (*Pinophilus*).

Sub-tribe *PROCIRRI*.Genus **Eucirrus**.

miricornis Fauv. Rev. d'Ent., xiv, 1895, p. 216.

Genus **Palaminus** Er.

bryanti, n. sp. Trans. Ent. Soc., 1920, p. 280.

parvus, n. sp. Trans. Ent. Soc., 1918, p. 71.

II. Tribe PAEDERINI.

Genus **Paederus** F.

tamulus Er. Gen. Spec. Staph., p. 661.

Genus **Astenus** Steph.

castaneus, n. sp. Trans. Ent. Soc., 1920, p. 281.

gracilentus Fauv. Ann. Mus. Civ. Gen., xv, 1879-80, p. 83.

gracilis Kr. Arch. Naturgesch., xxv, 1859, i, p. 147.

orientalis, n. sp. Trans. Ent. Soc., 1918, p. 71.

Genus **Stilicopsis** Sachse.

breviceps Fauv. Rev. d'Ent., xxiv, 1905, p. 138.

obliqua, n. sp. Trans. Ent. Soc., 1918, p. 72.

persimilis, n. sp. Trans. Ent. Soc., 1918, p. 72.

Genus **Stilicus** Latr.

ocularis Fauv. Rev. d'Ent., xiv, 1895, p. 226.

Genus **Psilotrachelus** Kr.

crassus Kr. Arch. Naturgesch., xxv, 1859, i, p. 124.

Genus **Thinocharis** Kr.

- carinicollis* Kr. Arch. Naturgesch., xxv, 1859, i, p. 143.
nigricans, n. sp. Trans. Ent. Soc., 1918, p. 73.
pygmaea Kr. Arch. Naturgesch., xxv, 1859, i, p. 143.

Genus **Acanthoglossa** Kr.

- hirta* Kr. Arch. Naturgesch., xxv, 1859, i, p. 144.

Genus **Medon** Steph.

Sub-gen. *Medon* s.str.

- opacellus* Fauv. Rev. d'Ent., xiv, 1895, p. 231.
orientalis, n. sp. Trans. Ent. Soc., 1920, p. 281.
rubicundus, n. sp. Trans. Ent. Soc., 1918, p. 73.

Sub-gen. *Hypomedon* Cas.

- debilicornis* Woll. Cat. Col. Mad., 1857, p. 194.
fasciatus, n. sp. Trans. Ent. Soc., 1918, p. 74.
granulatus, n. sp. Trans. Ent. Soc., 1918, p. 75.
lucens, n. sp. Trans. Ent. Soc., 1918, p. 75.

Sub-gen. *Lithocharis* Boisd. & Lacord.

- ochraceus* Grav. Col. Mic. Brunsv., 1802, p. 59.
uvidus Kr. Arch. Naturgesch., xxv, 1859, i, p. 138.

Sub-gen. *Isocheilus* Sharp.

- staphylinoides* Kr. Arch. Naturgesch., xxv, 1859, i, p. 134.

Sub-gen. *Charichirus* Sharp.

- chinensis* Boh. Eugen. Resa, 1858, Ins. p. 32.
terminalis, n. sp. Trans. Ent. Soc., 1920, p. 282.

Parascopaeus, n. sub-gen.

- nitidus*, n. sp. Trans. Ent. Soc., 1918, p. 76.

Genus **Scopaeus** Er.

- limbatus* Kr. Arch. Naturgesch., xxv, 1859, i, p. 130.
micros Kr. Arch. Naturgesch., xxv, 1859, i, p. 132.
niger, n. sp. Trans. Ent. Soc., 1918, p. 77.
pallidulus, Kr. Arch. Naturgesch., xxv, 1859, i, p. 131.
puncticeps Kr. Arch. Naturgesch., xxv, 1859, i, p. 132.
testaceus Motsch. Bull. Mosc., 1858, ii, p. 642.

Genus **Calliderma** Motsch.

nitens, n. sp. Trans. Ent. Soc., 1918, p. 79.

rufum, n. sp. Trans. Ent. Soc., 1918, p. 78.

rugicolle, n. sp. Trans. Ent. Soc., 1918, p. 80.

Genus **Cryptobium** Mannerh.

filum Kr. Arch. Naturgesch., xxv, 1859, i, p. 119.

foveatum, n. sp. Trans. Ent. Soc., 1918, p. 81.

VI. Sub-fam. **STAPHYLININAE**.I. Tribe **XANTHOLININI**.Genus **Oligolinus** Casey.

parvus, n. sp. Trans. Ent. Soc., 1918, p. 81.

Genus **Leptacinus** Er.

tricolor Kr. Arch. Naturgesch., xxv, 1859, i, p. 110.

Genus **Somoleptus** Sharp.

linearis, n. sp. Trans. Ent. Soc., 1918, p. 82.

Genus **Eulissus** Mannerh.

lateralis, n. sp. Trans. Ent. Soc., 1918, p. 83.

Genus **Thyreocephalus** Guer.

annulatus Fauv. Rev. d'Ent., xiv, 1895, p. 241.

Genus **Diochus** Er.

pulchellus, n. sp. Trans. Ent. Soc., 1918, p. 84.

II. Tribe **STAPHYLININI**.I. Sub-tribe **STAPHYLINI**.

Holisomimus, n. gen. Trans. Ent. Soc., 1920, p. 283.

cingulatus, n. sp. Trans. Ent. Soc., 1918, p. 85 (*Holisus*).

parvus, n. sp. Trans. Ent. Soc., 1918, p. 85 (*Holisus*).

Genus **Actobius** Fauv.

laticeps, n. sp. Trans. Ent. Soc., 1918, p. 86.

Genus **Philonthus** Curtis.

- belonuchoides*, n. sp. Trans. Ent. Soc., 1918, p. 88.
castaneipennis, n. sp. Trans. Ent. Soc., 1918, p. 87.
circumductus Fauv. Rev. d'Ent., xiv, 1895, p. 263.
crassicornis Fauv. Rev. d'Ent., xiv, 1895, p. 264.
delicatulus Boh. Eugen. Resa, 1858, Ins. p. 29.
flavocinctus Motsch. Bull. Mosc., xxxi, 1858, p. 663.
gemellus Kr. Arch. Naturgesch., xxv, 1859, i, p. 91.
geminus Kr. Arch. Naturgesch., xxv, 1859, i, p. 87.
longiceps Fauv. Ann. Mus. Civ. Gen., xv, p. 104.
maritimus Motsch. (*Gabrius*). Bull. Mosc., xxxi, 1858, ii,
p. 661.
notabilis Kr. Arch. Naturgesch., xxv, 1859, i, p. 79.
pulchellus Kr. (*Gabrius*). Arch. Naturgesch., xxv, i, p. 92.
sulcatus, n. sp. Trans. Ent. Soc., 1918, p. 87.

Genus **Orthidus** Muls. & Rey.

- cupreipennis*, n. sp. Trans. Ent. Soc., 1918, p. 89.

Genus **Cafius** Steph.

- corallicola* Fairm. Rev. Zool., 1849, p. 289.
nauticus Fairm. Rev. Zool., 1849, p. 288.

Genus **Hesperus** Fauv.

- laevigatus* Fauv. Rev. d'Ent., xiv, 1895, p. 259.

Genus **Belonuchus** Nordm.

- aeneipennis* Fauv. Rev. d'Ent., xiv, 1895, p. 268.
mutator Fauv. Ann. Mus. Civ. Gen., xv, p. 106.

III. Tribe QUESIINI.

Genus **Acylophorus** Nordm.

- rotundicollis*, n. sp. Trans. Ent. Soc., 1918, p. 90.

VII. Sub-fam. **PYGOSTENINAE**.

Genus **Delibius** Fauv.

- longicornis* Fauv. Rev. d'Ent., xviii, 1899, p. 13.

VIII. Sub-fam. **TACHYPORINAE.**I. Tribe **MEGARTHROPSINI.****Megarthritis**, n. gen.

decorata, n. sp. Trans. Ent. Soc., 1918, p. 232.

II. Tribe **TACHYPORINI.**Genus **Atanygnathus** Jacobson.

terminalis Er. Kaf. Mark. Brand., p. 418.

Genus **Conosoma** Kr.

abdominale, n. sp. Trans. Ent. Soc., 1918, p. 235.

ceylanense Kr. Arch. Naturgesch., xxv, 1859, i, p. 62.

championi, n. sp. Trans. Ent. Soc., 1918, p. 236.

flavoguttatum, n. sp. Trans. Ent. Soc., 1918, p. 234.

malayanum, n. sp. Trans. Ent. Soc., 1920, p. 283.

nigromaculatum, n. sp. Trans. Ent. Soc., 1918, p. 237.

perplexum, n. sp. Trans. Ent. Soc., 1918, p. 237.

rufobrunneum, n. sp. Trans. Ent. Soc., 1918, p. 234.

rufotestaceum, n. sp. Trans. Ent. Soc., 1918, p. 238.

robustum, n. sp. Trans. Ent. Soc., 1918, p. 233.

suave Fauv. Rev. d'Ent., xiv, 1895, p. 284.

walkeri, n. sp. Trans. Ent. Soc., 1918, p. 236.

Genus **Tachinomorphus** Kr.

ceylonicus Bernh. D. E. Z., 1902, p. 24.

fulvipes Er. Gen. Spec. Staph., p. 921.

Genus **Coproporus** Kr.

atomus Kr. Arch. Naturgesch., xxv, 1859, i, p. 58.

brunneicollis Motsch. Bull. Mosc., xxxi, 1858, iii, p. 220.

fasciipennis Kr. Arch. Naturgesch., xxv, 1859, i, p. 59.

flavipennis, n. sp. Trans. Ent. Soc., 1918, p. 239.

melanarius Er. Gen. Spec. Staph., p. 252.

minimus Motsch. Bull. Mosc., xxxi, 1858, ii, p. 220.

parvulus, n. sp. Trans. Ent. Soc., 1918, p. 240.

rufiventris, n. sp. Trans. Ent. Soc., 1918, p. 238.

secretus Bernh. Col. Rundsch., 1917 (7/9), p. 4.

varians, n. sp. Trans. Ent. Soc., 1920, p. 284.

Genus **Leucoparyphus** Kr.

silphoides L. Syst. Nat., i, 2, 1735, p. 684.

Mimocyptus, n. gen.

globulus, n. sp. Trans. Ent. Soc., 1918, p. 241.

IX. Sub-fam. **ADIMOPSINAE**.

Adimopsis, n. gen.

rufobrunnea, n. sp. Trans. Ent. Soc., 1918, p. 243.

X. Sub-fam. **ALEOCHARINAE**.

I. Tribe GYMNUSINI.

Genus **Leucocraspedum** Kr.

nigrum, n. sp. Trans. Ent. Soc., 1918, p. 243.

II. Tribe MYLLAENINI.

Genus **Myllaena** Er.

faberensis, n. sp. Trans. Ent. Soc., 1918, p. 244.

III. Tribe PRONOMAEINI.

Genus **Pronomaea** Er.

leontopolitana, n. sp. Trans. Ent. Soc., 1918, p. 245.

IV. Tribe DIGLOTTINI.

Genus **Diglotta** Champ.

testaceipennis, n. sp. Trans. Ent. Soc., 1918, p. 245.

V. Tribe OLIGOTINI.

Genus **Oligota** Mannerh.

forticornis, n. sp. Trans. Ent. Soc., 1920, p. 212.

moultoni, n. sp. Trans. Ent. Soc., 1920, p. 212.

VI. Tribe BOLITOCCHARINI.

Pseudoligota, n. gen.

robusta, n. sp. Trans. Ent. Soc., 1920, p. 215.

varians, n. sp. Trans. Ent. Soc., 1920, p. 214.

Genus **Gyrophaena** Mannerh.

appendiculata Motsch. Bull. Mosc., 1858, iii, p. 228.

laminata Kr. Arch. Naturgesch., xxv, 1859, i, p. 45.

Sub-gen. *Gyrophæna* s.str.

- bidens*, n. sp. Trans. Ent. Soc., 1920, p. 217.
crenulata, n. sp. Trans. Ent. Soc., 1920, p. 216.
cristata, n. sp. Trans. Ent. Soc. 1920, p. 217.
dubia, n. sp. Trans. Ent. Soc., 1920, p. 218.
granulosa, n. sp. Trans. Ent. Soc., 1920, p. 216.
irregularis, n. sp. Trans. Ent. Soc., 1920, p. 218.
moultoni, n. sp. Trans. Ent. Soc., 1920, p. 219.
tridentata, n. sp. Trans. Ent. Soc., 1920, p. 215.

Sub-gen. *Phænogyra* Rey.

- metallica*, n. sp. Trans. Ent. Soc., 1920, p. 220.

Sternotropa, n. gen.

- nigra*, n. sp. Trans. Ent. Soc., 1920, p. 221.
ruficollis, n. sp. Trans. Ent. Soc., 1920, p. 221.

Adelarthra, n. gen.

- barbara*, n. sp. Trans. Ent. Soc., 1920, p. 223.

Hetairotermes.*

* Nom. nov. for *Termophila* Lea, nom. praeoc.

- agilis*, n. sp. Trans. Ent. Soc., 1920, p. 223.
piceus, n. sp. Trans. Ent. Soc., 1920, p. 224.

Pseudatheta, n. gen.

- elegans*, n. sp. Trans. Ent. Soc., 1920, p. 225.

Pelekoglossa, n. gen.

- cingulata*, n. sp. Trans. Ent. Soc., 1920, p. 227.

Genus **Placusa**, Er.

- bispina*, n. sp. Trans. Ent. Soc., 1920, p. 228.
conura, n. sp. Trans. Ent. Soc., 1920, p. 227.
lobata, n. sp. Trans. Ent. Soc., 1920, p. 228.
notabilis, n. sp. Trans. Ent. Soc., 1920, p. 229.

Pseudoplacusa, n. gen.

- rufiventris*, n. sp. Trans. Ent. Soc., 1920, p. 230.

Chledophila, n. gen.

- annularis*, n. sp. Trans. Ent. Soc., 1920, p. 232.

Neosilusa, n. gen.

- ceylonica* Kr. Arch. Naturgesch., xxv, 1859, i, p. 10
(*Stenusa*).
moultoni, n. sp. Trans. Ent. Soc., 1920, p. 233.

Ousilusa, n. gen.

- castanea*, n. sp. Trans. Ent. Soc., 1920, p. 235.
myrmecobia, n. sp. Trans. Ent. Soc., 1920, p. 235.

Prosilusa, n. gen.

- rufa*, n. sp. Trans. Ent. Soc., 1920, p. 237.

Deralia, n. gen.

- fuscipennis*, n. sp. Trans. Ent. Soc., 1920, p. 238.

Pseudophaena, n. gen.

- castanea*, n. sp. Trans. Ent. Soc., 1920, p. 240.

Genus **Coenonica** Kr.

- angusticollis*, n. sp. Trans. Ent. Soc., 1920, p. 240.
puncticollis Kr. Linn. Ent., xi, p. 47.
stricticollis, n. sp. Trans. Ent. Soc., 1920, p. 241.

Mimomalota, n. gen.

- bispina*, n. sp. Trans. Ent. Soc., 1920, p. 243.
testacea, n. sp. Trans. Ent. Soc., 1920, p. 243.

Neomalota, n. gen.

- cingulata*, n. sp. Trans. Ent. Soc., 1920, p. 245.

Lampromalota, n. gen.

- brunneicollis*, n. sp. Trans. Ent. Soc., 1920, p. 246.

Genus **Homalota** Mannerh.

- bidens*, n. sp. Trans. Ent. Soc., 1920, p. 247.
cingulata, n. sp. Trans. Ent. Soc., 1920, p. 248.
denticulata, n. sp. Trans. Ent. Soc., 1920, p. 250.
fuscipennis, n. sp. Trans. Ent. Soc., 1920, p. 248.
nitescens, n. sp. Trans. Ent. Soc., 1920, p. 247.
platygaster, Kr. Arch. Naturgesch., xxv, 1859, i, p. 33.
serrata, n. sp. Trans. Ent. Soc., 1920, p. 249.
tuberculicollis Kr. Arch. Naturgesch., xxv, 1859, p. 33.
variventris Kr. Arch. Naturgesch., xxv, 1859, p. 34.

Genus **Thectura** Thoms.

brunneicollis, n. sp. Trans. Ent. Soc., 1920, p. 250.

Genus **Heterota** Rey.

arenaria, n. sp. Trans. Ent. Soc., 1920, p. 251.

Paractocharis, n. gen.

fucicola, n. sp. Ent. Mo. Mag., 1917, p. 154.

VII. Tribe MYRMEDONIINI.

Genus **Falagria** Mannerh.Sub-gen. *Falagria* s.str.

brevicornis, n. sp. Trans. Ent. Soc., 1920, p. 252.

dimidiata Motsch. Bull. Mosc., 1858, ii, p. 260.

flavipennis, n. sp. Trans. Ent. Soc., 1920, p. 253.

pygmaea Kr. Arch. Naturgesch., xxv, 1859, i, p. 7.

tenuicornis, n. sp. Trans. Ent. Soc., 1920, p. 252.

Sub-gen. *Cardiola* Rey.

vestita Boh. Eugen. Resa, 1858, Ins., p. 25.

Genus **Amaurodera** Fauv.

veluticollis Motsch. Bull. Mosc., 1858, ii, p. 261.

Eusteniamorpha, n. gen.

rufa, n. sp. Trans. Ent. Soc., 1920, p. 254.

Genus **Pelioptera** Kr.

micans Kr. Linnaea Ent., xi, p. 55.

opaca Kr. Linnaea Ent., xi, p. 56.

Genus **Atheta** Thoms.Sub-gen. *Glossola* Fowler.

moultoni, n. sp. Trans. Ent. Soc., 1920, p. 255.

Sub-gen. *Metaxyia* Rey.

alophila, n. sp. Trans. Ent. Soc., 1920, p. 256.

Genus (?) **Dralica** Rey.

picea, n. sp. Trans. Ent. Soc., 1920, p. 256.

Sub-gen. *Microdota* Rey.

- inutilis* Kr. Arch. Naturgesch., xxv, 1859, i, p. 35.
malayana, n. sp. Trans. Ent. Soc., 1920, p. 257.
melata, n. sp. Trans. Ent. Soc., 1920, p. 257.
purpurascens, n. sp. Trans. Ent. Soc., 1920, p. 259.
putridula Kr. Arch. Naturgesch., xxv, 1859, i, p. 35.
vulgaris, n. sp. Trans. Ent. Soc., 1920, p. 258.

Sub-gen. *Atheta* s.str.

- dilutipennis* Motsch. Bull. Mosc., 1858, ii, p. 252.
miriventris, n. sp. Trans. Ent. Soc., 1920, p. 259.

Sub-gen. *Dimetrota* Rey.

- carpophila*, n. sp. Trans. Ent. Soc., 1920, p. 260.
mycetophaga, n. sp. Trans. Ent. Soc., 1920, p. 262.
xylophila, n. sp. Trans. Ent. Soc., 1920, p. 261.

Sub-gen. *Datomicra* Rey.

- mycetophila*, n. sp. Trans. Ent. Soc., 1920, p. 263.
onthophila, n. sp. Trans. Ent. Soc., 1920, p. 262.

Sub-gen. *Colpodota* Rey.

- ruparia*, n. sp. Trans. Ent. Soc., 1920, p. 264.

Sub-gen. *Acrotona* Rey.

- annuliventris* Kr. Arch. Naturgesch., xxv, 1859, i, p. 40.
rufiventris, n. sp. Trans. Ent. Soc., 1920, p. 264.

Exatheta, n. gen.

- cingulata*, n. sp. Trans. Ent. Soc., 1920, p. 266.
consors, n. sp. Trans. Ent. Soc., 1920, p. 266.

Mimatheta, n. gen.

- fungicola*, n. sp. Trans. Ent. Soc., 1920, p. 267.

Mimacrotona, n. gen.

- cingulata*, n. sp. Trans. Ent. Soc., 1920, p. 269.

Paratheta, n. gen.

- carnivora*, n. sp. Trans. Ent. Soc., 1920, p. 270.

Fenyesia, n. gen.

- nigra*, n. sp. Trans. Ent. Soc., 1920, p. 271.

Genus **Termitoptochus**, Silv.*indicus* Silv. Bol. Lab. Port., 5, 1909, p. 39.**Myrmedonota**, n. gen.*cingulata*, n. sp. Trans. Ent. Soc., 1920, p. 272.Genus **Myrmedonia** Er.*apicalis*, n. sp. Trans. Ent. Soc., 1920, p. 273.*indorum* Fauv. Rev. d'Ent., xxii, p. 162.Genus **Schistogenia** Kr.*crenicollis* Kr. Linnaea Ent., xi, p. 39.

ALEOCHARINI.

Myrmedonella, n. gen.*rufa*, n. sp. Trans. Ent. Soc., 1920, p. 275.Genus **Tetrasticta** Kr.*polita* Kr. Linn. Ent., 1857, p. 55.**Paraleochara**, n. gen.*fungivora*, n. sp. Trans. Ent. Soc., 1920, p. 276.Genus **Hoplandria** Kr.*frugivora*, n. sp. Trans. Ent. Soc., 1920, p. 277.Genus **Aleochara** Grav.Sub-gen. *Xenochara* Rey.*puberula* Klug. Ins. Madag., p. 139.Sub-gen. *Heterochara* Rey.*(croceipennis* Motsch. Bull. Mosc., 1858, ii, p. 238.)*v. maculipennis* Kr. Arch. Naturgesch., xxv, 1859, i, p. 17.Sub-gen. *Polychara* Rey.*asiatica* Kr. Arch. Naturgesch., xxv, 1859, i, p. 15.*nigra* Kr. Arch. Naturgesch., xxv, 1859, i, p. 13.*viatica* Fauv. Rev. d'Ent., xxiii, p. 67.

CORRIGENDA.

1918, p. 65. Delete description of *Aploderus testaceus*, n. sp., which is *Oxytelus thoracicus* Motsch., Bull. Mosc., 1857, iv, p. 504.

1918, p. 68, line 28, for *Tesnus* read *Hypostenus*.

1920, p. 215, line 5, for *robustus* read *robusta*.

Cameron, Malcolm. 1921. "XVI. New species of Staphylinidae from Singapore. Part IV (Conclusion)." *Transactions of the Entomological Society of London* 68, 347–413. <https://doi.org/10.1111/j.1365-2311.1921.tb00228.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/55146>

DOI: <https://doi.org/10.1111/j.1365-2311.1921.tb00228.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/57040>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.